The background of the cover is a dark blue field filled with numerous spermatozoa. Each sperm is stained with a blue dye, likely DAPI, to visualize the nucleus. Superimposed on the blue nuclei are small, bright spots of red, green, and yellow fluorescence, which represent the localization of specific DNA sequences (probed with FISH) within the sperm's genome. The sperm are oriented in various directions, some showing the characteristic curved shape of the head and tail.

Universidad de Salamanca

Instituto de Biología Molecular y Celular de Cáncer

Análisis funcional de los genes
***Usp26* y *Hsf2bp* como responsables**
de infertilidad humana

Tesis Doctoral
Natalia Felipe Medina
Noviembre, 2020

Análisis funcional de los genes *Usp26* y *Hsf2bp* como responsables de infertilidad humana

Tesis Doctoral

Natalia Felipe Medina

Director: Dr. Alberto Martín Pendás

Co-director: Dra. Elena Llano Cuadra

Universidad de Salamanca

Instituto de Biología Molecular y Celular del Cáncer

Salamanca, 2020

El Dr. Alberto Martín Pendás, Investigador principal del Instituto de Biología Molecular y Celular del Cáncer (IBMCC) y la Dra. Elena Llano Cuadra, Profesora Titular del Departamento de Fisiología y Farmacología de la Universidad de Salamanca,

Certifican que Natalia Felipe Medina ha realizado bajo su dirección el trabajo de tesis doctoral titulado:

**Análisis funcional de los genes *Usp26* y *Hsf2bp*
como responsables de infertilidad humana**

Revisado el presente trabajo, consideran que reúne todos los méritos necesarios para su presentación y defensa, con el fin de optar al grado de Doctor por el programa “Biociencias: Biología y Clínica del Cáncer y Medicina Traslacional” de la Universidad de Salamanca.

Y para que así conste firman el presente certificado en Salamanca a 26 de octubre de 2020

Dr. Alberto Martín Pendás

Dra. Elena Llano Cuadra

AGRADECIMIENTOS

UNIÓN EUROPEA
Fondo Social Europeo

La presente tesis ha sido financiada por:

- Beca predoctoral (EDU/310/2015) de la Junta de Castilla y León/Fondo Social Europeo.
- Fundación Investigación del Cáncer de la Universidad de Salamanca. Escalera de Excelencia de la Consejería de Educación de la Junta de Castilla y León (CLC-2017-01).
- Ministerio de Ciencia e Innovación (BFU_2014-59307-R y BFU2017-89408-R).

ÍNDICE

RESUMEN	17
ABREVIATURAS.....	21
INTRODUCCIÓN	25
1. Infertilidad humana	27
1.1. Bases genéticas de la infertilidad	28
2. Gametogénesis	30
2.1. Especificación de las células germinales primordiales.....	30
2.2. Oogénesis	32
2.3. Espermatogénesis	34
3. Meiosis	36
3.1. Profase I.....	36
3.2. Divisiones meióticas	39
4. Recombinación meiótica	40
4.1. Inicio de la recombinación	41
4.2. Reparación de las DSBs	42
4.2.1. <i>Procesamiento de las DSBs.....</i>	42
4.2.2. <i>Invasión del cromosoma homólogo</i>	43
4.2.3. <i>Procesamiento de los intermediarios de recombinación</i>	44
4.2.4. <i>Resolución de los intermediarios de recombinación</i>	45
OBJETIVOS	47
MATERIAL Y MÉTODOS	51
1. Modelos murinos	53
1.1. Bienestar animal.....	53
1.2. Generación de ratones modificados genéticamente mediante CRISPR/Cas9.....	53
1.2.1. <i>Diseño de sgRNAs</i>	53
1.2.2. <i>Obtención de sgRNAs, Cas9 y ssODNs.....</i>	54
1.2.3. <i>Microinyección del sistema CRISPR/Cas9 en cigotos.....</i>	56
1.2.4. <i>Caracterización de los fundadores y establecimiento final de las líneas.....</i>	56
1.3. Modelos murinos generados como herramienta en esta tesis	57
1.4. Otros modelos	59
2. Técnicas de Biología Molecular.....	59
2.1. Extracción de DNA genómico	59
2.1.1. <i>Extracción por lisis alcalina.....</i>	59
2.1.2. <i>Extracción con fenol/cloroformo</i>	59

2.2.	Genotipado por PCR	59
2.3.	Generación de vectores de expresión	60
2.3.1.	<i>Extracción de RNA</i>	60
2.3.2.	<i>Transcripción reversa-PCR (RT-PCR)</i>	60
2.3.3.	<i>Preparación de insertos y clonning</i>	60
2.3.4.	<i>Transformación de células competentes DH5α</i>	61
2.3.5.	<i>Extracción de DNA plasmídico de un cultivo bacteriano por lisis alcalina</i>	61
2.3.6.	<i>Análisis de restricción</i>	61
2.3.7.	<i>Extracción de DNA plasmídico por lisis alcalina seguido de purificación por columna de afinidad</i>	61
2.4.	Análisis de proteínas	62
2.4.1.	<i>Extracción de proteínas de testículo de ratón</i>	62
2.4.2.	<i>Inmunoprecipitación de proteínas endógenas de testículo</i>	62
2.4.3.	<i>Crosslink de los anticuerpos a bolas de Sefarosa</i>	63
2.4.4.	<i>Co-inmunoprecipitación de proteínas sobrepresadas</i>	63
2.4.5.	<i>Western Blot</i>	64
2.4.6.	<i>Análisis por espectrometría de masas (MS/MS)</i>	65
2.5.	Ensayo de doble híbrido y screening.....	66
2.6.	Transcripción-traducción <i>in vitro</i>	66
2.7.	Ensayo de unión al DNA	66
3.	Técnicas citológicas	67
3.1.	Análisis histopatológico	67
3.2.	Esparcido de espermatozoides (<i>spreading</i>)	67
3.3.	Aplastado de túbulos seminíferos (<i>squashing</i>)	68
3.4.	Esparcido de oocitos (<i>spreading</i>)	68
3.5.	Inmunofluorescencia	69
3.6.	Adquisición y análisis de imágenes	71
3.7.	Microscopía de super-resolución	72
3.8.	Electroporación <i>in vivo</i> de testículos.....	72
4.	Cultivos celulares	72
4.1.	Líneas celulares y condiciones de cultivo.....	72
4.2.	Transfección de líneas celulares.....	73
5.	Análisis estadístico	73
	RESULTADOS.....	75
1.	USP26 no es esencial para la fertilidad en ratón	77

1.1.	Generación de los modelos murinos mutantes para <i>Usp26</i>	78
1.2.	Los ratones deficientes en <i>USP26</i> son fértiles	80
1.3.	Los espermatoцитos deficientes en <i>USP26</i> no muestran defectos en sinapsis, dinámica de las DSBs ni recombinación meiótica	83
2.	Una variante en <i>HSF2BP</i> causa POI a través de su nuevo interactor <i>BRME1</i>	86
2.1.	La variante <i>S167L</i> en el gen <i>HSF2BP</i> es una firme candidata a producir POI en humanos	86
2.2.	Generación de los modelos murinos mutantes para <i>Hsf2bp</i>	88
2.3.	<i>HSF2BP</i> es esencial para la fertilidad en ratón y la variante <i>S167L</i> produce una reducción parcial de la misma	90
2.4.	La ausencia de <i>HSF2BP</i> produce fuertes defectos en sinapsis y la variante <i>S167L</i> provoca una pérdida parcial de función	92
2.5.	La variante <i>S167L</i> produce una reducción de la expresión/estabilidad de la proteína	94
2.6.	<i>HSF2BP</i> es esencial para la recombinación meiótica y la variante <i>S167L</i> produce defectos severos en este proceso.....	95
2.7.	Los defectos meióticos producidos por la variante <i>S167L</i> no están mediados por <i>BRCA2</i>	100
2.8.	<i>BRME1</i> , un nuevo interactor de <i>HSF2BP</i>	101
2.9.	<i>BRME1</i> es un nuevo componente de los RNs que co-localiza con <i>HSF2BP</i> .	102
2.10.	<i>BRME1</i> interacciona con <i>HSF2BP</i> a través de su extremo C-terminal	104
2.11.	<i>HSF2BP</i> es necesaria para la localización/estabilidad de <i>BRME1</i>	107
2.12.	<i>BRME1</i> y <i>HSF2BP</i> intervienen en los eventos tempranos tras la formación de DSBs.....	108
2.13.	<i>BRME1</i> y <i>HSF2BP</i> no tienen capacidad de unión al DNA	109
2.14.	<i>BRME1</i> es esencial para la fertilidad en ratón	109
2.15.	La ausencia de <i>BRME1</i> produce defectos en la recombinación meiótica ...	112
2.16.	<i>BRME1</i> es necesaria para la localización/estabilidad de <i>HSF2BP</i>	115
2.17.	<i>BRME1</i> y <i>HSF2BP</i> forman un complejo multimérico con <i>PALB2</i> y <i>BRCA2</i> ...	117
	DISCUSIÓN.....	121
1.	<i>USP26</i> no es esencial para la fertilidad en ratón	123
1.1.	La ausencia de <i>USP26</i> no afecta a la meiosis ni la fertilidad en ratón	123
1.2.	Asociación entre <i>USP26</i> e infertilidad humana.....	123
2.	Una variante en <i>HSF2BP</i> causa POI a través de su nuevo interactor <i>BRME1</i> ...	125
2.1.	La variante <i>S167L</i> en <i>HSF2BP</i> afecta a la fertilidad en mujeres	125
2.2.	Los mutantes de <i>Hsf2bp</i> y <i>Brme1</i> presentan dimorfismo sexual	126
2.3.	<i>HSF2BP</i> y <i>BRME1</i> forman un complejo funcional esencial para la recombinación.....	127

2.4. La variante HSF2BP-S167L se comporta como un alelo hipomorfo en ratón	129
3. Relevancia del estudio	131
CONCLUSIONES.....	133
ANEXO	137
1. Material suplementario	139
BIBLIOGRAFÍA	147
PUBLICACIONES	165

RESUMEN

La infertilidad humana es un problema de salud pública emergente que afecta al 10-15% de las parejas en edad reproductiva desconociéndose su etiología en gran parte de los casos. Se estima que las alteraciones genéticas subyacen a la mitad de estos casos y cientos de genes candidatos han sido identificados mediante el estudio de la gametogénesis en modelos animales. Sin embargo, son pocos los casos en los que se ha podido establecer una relación causal directa entre estos genes y el desarrollo de infertilidad.

En la presente tesis, hemos llevado a cabo el análisis funcional de *Usp26* y *Hsf2bp*, dos genes candidatos de infertilidad humana. USP26 es una enzima desubiquitinante con un patrón de expresión restringido a la línea germinal masculina. En la última década numerosos estudios han reportado la presencia de polimorfismos en *USP26* asociados con trastornos de infertilidad masculina. Sin embargo, también se han descrito polimorfismos en varones que presentan una espermatogénesis normal. Teniendo en cuenta esta controversia y la función desconocida de USP26 en la gametogénesis, en este trabajo hemos abordado el análisis funcional de USP26 mediante la generación de un modelo animal de pérdida de función. Nuestros resultados describen la primera evidencia in vivo de que USP26 no es esencial para la gametogénesis y la fertilidad del ratón.

HSF2BP es una proteína descrita inicialmente como interactor del *Heat Shock Factor 2* (HSF2) y cuya función es desconocida. La secuenciación del exoma de una familia con 3 casos de insuficiencia ovárica primaria identificó una variante en *HSF2BP* (S167L) como mejor candidata causal de la infertilidad. Mediante la caracterización de un modelo murino nulo para *Hsf2bp* y un modelo humanizado portador de la variante S167L hemos demostrado que HSF2BP es una proteína esencial para la recombinación meiótica y que esta variante produce una pérdida parcial de su expresión y/o función. La ausencia de HSF2BP o la presencia de su variante patogénica producen defectos en la localización de las recombinasas RAD51/DMC1 y en la formación de los COs los cuales provocan infertilidad y subfertilidad, respectivamente. Mecánicamente, hemos identificado a C19ORF57/BRME1 como un potente interactor y estabilizador de HSF2BP cuya ausencia produce un fenotipo muy similar al de los mutantes nulos para *Hsf2bp*. De acuerdo con ello, la presencia de la variante causante de POI en el heterodímero HSF2BP/BRME1 provoca una reducción de su expresión que es la responsable última de los defectos meióticos descritos los cuales señalizan a través de un macro-complejo formado por HSF2BP/BRME1/BRCA2/RAD51/PALB1/RPA. Por tanto, en este trabajo hemos identificado un nuevo complejo esencial para la recombinación meiótica cuya caracterización proporciona nueva información sobre el mecanismo molecular que subyace a la patogenicidad de la variante S167L en la insuficiencia ovárica primaria.

ABREVIATURAS

Ab: anticuerpo (*Antibody*)

AEs: elementos axiales (*Axial Elements*)

bp: pares de bases (*base pair*)

CE: elemento central (*Central Element*)

co-IP: co-inmunoprecipitación

CO: *crossover*

CRISPR: *Clustered Regularly Interspaced Short Palindromic Repeats*

crRNA: crispr RNA

dHJ: uniones dobles de Holliday (*double Holliday Junctions*)

dpc: días post-coito

dpp: días post-parto

DSBs: roturas de doble hebra (*Double Strand Breaks*)

dsDNA: DNA de doble hebra (*double-stranded DNA*)

FSH: hormona estimulante de folículos (*Follicle Stimulating Hormone*)

H&E: tinción de hematoxilina-eosina

IF: Inmunofluorescencia

kDa: kiloDalton

KO: mutante nulo (*knockout*)

LEs: elementos laterales (*Lateral Elements*)

LH: hormona luteinizante (*Luteinizing Hormone*)

MSCI: Inactivación de los cromosomas sexuales meióticos (*Meiotic Sex Chromosome Inactivation*)

NCO: *non-crossover*

NHEJ: reparación de extremos no homólogos (*Non-Homologous End Joining*)

PAR: región pseudoautosómica (*PseudoAutosomal Region*)

PAS: tinción *Periodic Acid-Schiff*

PGCs: células germinales primordiales (*Primordial Germ Cells*)

POI: insuficiencia ovárica prematura (*Primary Ovarian Insufficiency*)

RA: ácido retinoico (*Retinoic Acid*)

RNs: nódulos de recombinación (*Recombination Nodules*)

SC: complejo sinaptonémico (*Synaptonemal Complex*)

sgRNA: RNA de guía única (*single guide RNA*)

sHJ: uniones simples de Holliday (*single Holliday Junctions*)

SSCs: células madre espermato goniales (*Spermatogonial Stem Cells*)

ssDNA: DNA de hebra simple (*single-stranded DNA*)

ssODN: oligonucleótido donador de hebra simple (*single-stranded donor oligonucleotide*)

TF: filamentos transversales (*Transverse Filaments*)

TnT: sistema de Transcripción/Traducción *in vitro*

tracrRNA: RNA transactivador (*trans-activating crRNA*)

TUNEL: *TdT mediated dUTP Nick End Labelling*, marcaje de apoptosis

u.a.: unidades arbitrarias

WB: *Western Blot*

WT: silvestre (*Wild-Type*)

Y2H: ensayo de doble híbrido en levaduras (*Yeast Two Hybrid*)

INTRODUCCIÓN

1. Infertilidad humana

El término infertilidad hace referencia a la incapacidad de concebir de una pareja. De forma más precisa, tanto el Comité Internacional de Monitorización de las Técnicas de Reproducción Asistida (ICMART) como la Organización Mundial para la Salud (OMS) definen la infertilidad como “una enfermedad del sistema reproductivo caracterizada por la imposibilidad de conseguir un embarazo clínico tras 12 o más meses de relaciones sexuales regulares sin protección”. Sin embargo, no existe un consenso absoluto en la definición de infertilidad lo que dificulta el correcto diagnóstico y la determinación de su prevalencia (Gurunath et al. 2011). Pese a ello, la infertilidad se considera un problema de salud pública importante ya que se estima que afecta al 10-15% de las parejas a nivel mundial. Esta puede deberse a factores masculinos, femeninos o a una combinación de ambos, representando cada uno de ellos aproximadamente el 25% de los casos mientras que en el resto se desconoce su origen (Shah et al. 2003; Isaksson and Tiitinen 2004).

Las causas de la **infertilidad masculina** engloban cualquier condición que afecte a la cantidad y/o calidad del espermatozoide. Entre ellas se encuentran las alteraciones en el transporte de los espermatozoides, la disfunción del eje hipotálamo-hipofisario (también denominado hipogonadismo secundario) o, la causa más comúnmente identificada, la disfunción o fallo testicular (hipogonadismo primario). Sin embargo, en la mitad de los casos la etiología se desconoce, diagnosticándose como infertilidad idiopática (Jose-Miller et al. 2007). Las principales alteraciones espermáticas identificadas en los pacientes son la oligozoospermia o reducción de la concentración de espermatozoides, la astenozoospermia o disminución de su movilidad, la teratozoospermia o incremento en el porcentaje de espermatozoides con morfología aberrante y, por último, la azoospermia o ausencia total de espermatozoides en el eyaculado. Dentro de este último subgrupo distinguimos la azoospermia obstructiva, debida a una obstrucción de los conductos seminales, y la azoospermia no obstructiva o secretora, en la cual la ausencia de espermatozoides se debe a un fallo en la espermatogénesis. La azoospermia afecta al 1% de la población masculina y al 10-15% de los varones infértiles, siendo la forma más frecuente la azoospermia no obstructiva (Lindsay 2015).

En el caso de la **infertilidad femenina** las causas incluyen la obstrucción de las trompas de Falopio, las afecciones en el útero, la endometriosis y la disfunción ovárica. El cese de la función del ovario ocurre de forma fisiológica como consecuencia de la edad (menopausia), en cambio, la disfunción ovárica hace referencia a un cese prematuro de dicha función que puede aparecer como consecuencia de factores ambientales como el consumo de tabaco, tratamientos quimioterápicos, trastornos endocrinos o alteraciones ginecológicas como son el síndrome de ovario poliquístico y la insuficiencia ovárica primaria (Jose-Miller et al. 2007). Actualmente, insuficiencia ovárica primaria (POI, *Primary ovarian insufficiency*) es el término preferencial para denominar lo que antes se refería como fallo ovárico prematuro o menopausia prematura. Clínicamente, la POI se caracteriza por la pérdida de función ovárica antes de los 40 años presentando amenorrea (ausencia de menstruación) durante 4 o más meses y niveles de FSH (*Follicle-Stimulating Hormone*) en el rango típico de la menopausia. Esta condición afecta al 1-3% de las mujeres menores de 40 años y es el resultado de un agotamiento prematuro de la reserva folicular, aunque es importante destacar que el 5-10% de estas mujeres son capaces de concebir después del diagnóstico con ayuda de técnicas de reproducción asistida (Nelson 2009; Golezar et al. 2019).

1.1. Bases genéticas de la infertilidad

Como ha quedado patente, las causas subyacentes a la infertilidad pueden ser muy diversas pudiendo estar asociadas tanto a factores fisiológicos y genéticos como ambientales e incluso sociales. Establecer de forma precisa la contribución de los factores genéticos en la reducción de la fertilidad resulta complejo si tenemos en cuenta que la mayoría de las causas descritas, si no todas, probablemente tienen un componente genético. Actualmente se estima que los factores genéticos se encuentran detrás del 50% de los casos de infertilidad siendo cada vez más los estudios dirigidos a su identificación. Estos estudios abarcan desde los primeros estadios de diferenciación de las células germinales hasta la maduración de los gametos y la fecundación, siendo especialmente relevantes los aspectos relacionados con la progresión de la meiosis. Tales esfuerzos han permitido identificar genes fundamentales para el mantenimiento de la fertilidad, pero además han puesto de manifiesto la enorme complejidad de estos procesos debido a su naturaleza multifactorial y el elevado número de genes implicados (Shah et al. 2003; Schimenti and Handel 2018).

Los defectos genéticos asociados con la infertilidad se pueden clasificar en: alteraciones cromosómicas, alteraciones de un solo gen o monogénicas, alteraciones poligénicas y alteraciones epigenéticas. Algunas de estas alteraciones se han descrito únicamente en hombres o en mujeres mientras que otras están presentes en los dos sexos, pudiendo afectar a la fertilidad en ambos o solo en uno de ellos (Zorrilla and Yatsenko 2013).

El 15-30% de las infertilidades masculinas son de origen genético siendo los defectos más frecuentes las alteraciones cromosómicas las cuales son detectadas en el 20% de los pacientes (Ferlin et al. 2007). Dentro de ellas podemos distinguir las que afectan a cromosomas completos, como ocurre en el Síndrome de Klinefelter (cariotipo 47,XXY) o el Síndrome 47,XYY (Martin 2008), y las alteraciones cromosómicas estructurales que incluyen deleciones, duplicaciones, translocaciones o inversiones de fragmentos cromosómicos de tamaño variable. Entre estas, destacan las microdeleciones del cromosoma Y que afectan a la región conocida como región del factor de azoospermia (AZF) la cual contiene genes esenciales para la espermatogénesis (Reijo et al. 1996; Matzuk and Lamb 2008). Dentro de este grupo también se encuentran las variaciones en el número de copias (CNVs, copy number variations) que afectan a pequeñas regiones (0,05 a 3 Mb) que alteran la dosis génica (Tüttelmann et al. 2011). También se han asociado con infertilidad masculina alteraciones epigenéticas como los defectos en la metilación del DNA y en las modificaciones post-transcripcionales (Yatsenko et al. 2013). Por último, en algunos pacientes se ha podido demostrar la asociación de mutaciones en un único gen con el desarrollo de infertilidad masculina. Este es el caso, entre otros, de las mutaciones en CFTR fuertemente asociadas con el desarrollo de aplasia congénita bilateral de los conductos deferentes que produce azoospermia obstructiva (Grangeia et al. 2007; O'Flynn O'Brien et al. 2010; Zorrilla and Yatsenko 2013).

El conocimiento de las bases genéticas de la infertilidad femenina ha avanzado más lentamente que en el caso de la masculina. Esto se debe a la mayor facilidad para obtener material de estudio en el caso de los hombres, dado que la espermatogénesis se produce de manera continua durante toda la vida adulta. Al contrario, en las mujeres la ovogénesis comienza de manera sincronizada durante el desarrollo embrionario y posteriormente se detiene generando una reserva de oocitos limitada. Pese a este menor conocimiento,

se sabe que las dos principales causas de infertilidad femenina, la disfunción ovárica y la endometriosis, tienen una importante predisposición familiar lo que sugiere un origen genético que en muchos casos ya ha sido demostrado (Mallepaly et al. 2017). En el caso concreto de la POI, las causas genéticas se encuentran detrás de aproximadamente el 20% de los casos (Rossetti et al. 2017).

La alteración cromosómica asociada con infertilidad mejor documentada en mujeres es el síndrome de Turner (o monosomía del X; 45,X0) (Abir et al. 2001). También se han descrito alteraciones cromosómicas estructurales, principalmente translocaciones recíprocas entre el cromosoma X y un autosoma (X:A) y CNVs que afectan a genes relacionados con POI (McGuire et al. 2011; Moysés-Oliveira et al. 2015). De igual manera que en la infertilidad masculina, se ha demostrado la asociación directa de alteraciones monogénicas con el desarrollo de POI como es el caso de las mutaciones en los *FMR1* (causante del síndrome X-frágil) y *BMP15*, ambos asociados al cromosoma X, o en genes autosómicos como *FOXL2* (Harris et al. 2002; Di Pasquale et al. 2004; Pastore and Johnson 2014). Por último, existen trastornos complejos como la endometriosis o el síndrome de ovario poliquístico los cuales presentan un origen multifactorial asociado a alteraciones genéticas, metabólicas, endocrinas e incluso ambientales (Zorrilla and Yatsenko 2013; Mallepaly et al. 2017).

La importancia de identificar las bases genéticas de la infertilidad no radica únicamente en el correcto diagnóstico de las parejas afectadas. En la era de las técnicas de reproducción asistida, definir estos defectos genéticos cobra más sentido que nunca. La utilización de estas técnicas conlleva sobrepasar las barreras de la selección natural generando un embrión viable en parejas genéticamente infértiles. En estos casos, diagnosticar si existe una causa genética subyacente tiene una clara importancia clínica, ya que podría tener implicaciones para la salud (general y reproductiva) tanto del paciente como del futuro hijo. Este conocimiento, además, puede tener valor pronóstico y en un futuro próximo ayudará al establecimiento de terapias personalizadas permitiendo dar soluciones más efectivas a los pacientes (O'Flynn O'Brien et al. 2010; Krausz and Riera-Escamilla 2018).

En la práctica, el estudio de las causas genéticas de la infertilidad se basa fundamentalmente en el trabajo con modelos animales, dada la imposibilidad de llevar a cabo estos estudios en humanos. En los últimos años se han identificado un gran número de factores genéticos asociados con infertilidad gracias al análisis de modelos murinos y a los estudios de asociación del genoma (GWAS, Genome-wide association study) (Matzuk and Lamb 2008; Hu et al. 2011; Gomez-H et al. 2016). Sin embargo, son pocos los casos en los cuales se puede establecer una relación causal directa con la infertilidad humana. Si bien es cierto que las alteraciones genéticas causantes de infertilidad tienen pocas probabilidades de propagarse en la población, algunas pueden identificarse mediante el estudio de pedigrees familiares, especialmente en familias con cierto grado de consanguinidad. En este sentido, en la última década la secuenciación del exoma completo (WES, Whole Exome Sequencing) de familias afectadas ha permitido identificar varios genes con una relación causal directa con la infertilidad en humanos. Este es el caso de los genes meióticos *SYCE1*, *STAG3*, *SPO11*, *MSH4*, *DMC1*, *BRCA2* o *MEIOB* (Caburet et al. 2014; De Vries et al. 2014; Carlosama et al. 2017; He et al. 2018; Caburet et al. 2019; Tran and Schimenti 2019; Caburet et al. 2020).

Pese a todos estos avances, la infertilidad idiopática sigue siendo el diagnóstico más frecuente entre las parejas afectadas por esta patología y actualmente se cree que una alta proporción de estos casos están

asociados con causas genéticas (Mallepaly et al. 2017; Krausz and Riera-Escamilla 2018). En este sentido, el estudio de la gametogénesis y los procesos reproductivos resulta fundamental para seguir ampliando este conocimiento y poder proporcionar mejores soluciones a las parejas afectadas.

2. Gametogénesis

La gametogénesis es el proceso biológico por el cual se obtienen los gametos a partir de las células germinales. Los gametos, oocitos y espermatozoides, son células sexuales haploides altamente especializadas que presentan características únicas, no solo desde el punto de vista genético, sino también epigenético y morfológico. La reducción de la ploidía es fundamental para que, tras la fusión de los gametos materno y paterno, se genere un cigoto diploide capaz de dar lugar a un nuevo individuo. Para ello, las células germinales llevan a cabo un tipo especial de división celular reduccional, la meiosis. Todo esto hace de la gametogénesis uno de los programas de diferenciación más complejos y estrictamente regulados de la biología del desarrollo.

2.1. Especificación de las células germinales primordiales

La gametogénesis comienza durante el desarrollo embrionario temprano con la aparición de unas células madre indiferenciadas denominadas células germinales primordiales (PGCs, Primordial Germ Cells). El estudio *in vivo* de la especificación de las PGCs en humanos se encuentra muy limitado por cuestiones técnicas y éticas, por lo que su conocimiento se basa en la extrapolación de los datos obtenidos en ratón o del estudio de células madre pluripotentes humanas diferenciadas *in vitro* a células similares a PGCs (hPGCLs, human PGC-like cells) (Irie et al. 2015; Sasaki et al. 2015). Estos estudios han contribuido a establecer los mecanismos de determinación de las PGCs en ambas especies poniendo de manifiesto sus similitudes y diferencias (Mitsunaga and Shioda 2018).

La especificación de las PGCs en ratón (mPGCs, mouse PGCs) comienza en el día embrionario (E) 6,0-6,5 con la aparición de un pequeño clúster de precursores en la región más proximal del epiblasto. En estas células la expresión de WNT3 junto con la señalización extrínseca mediada por BMP4 inducen la expresión del factor de transcripción *T* también conocido como *Brachyury* (Lawson et al. 1999; Ohinata et al. 2009; Aramaki et al. 2013). Este factor desencadena una cascada de genes encargados de orquestar la determinación hacia mPGC como *Blimp1*, *Prdm14*, *Tfap2c* y *Stella (Dppa3)*, reprimiendo la diferenciación somática y manteniendo el programa de pluripotencia característico de las mPGCs (expresión de genes como *Sox2*, *Nanog* y *Oct4*) (Saitou et al. 2002; Ohinata et al. 2005; Yamaji et al. 2008; Weber et al. 2010; Saitou and Miyauchi 2016). Este proceso genera una población de unas 40 mPGCs en la base del alantoides en el día E7,25 (Figura I).

Desarrollo embrionario

Figura I. Representación esquemática de los principales procesos que ocurren durante la gametogénesis, desde la especificación de las PGCs hasta la obtención de los gametos. MII: segunda división meiótica. MII*: la segunda división del primer corpúsculo polar no siempre sucede. Adaptado de (McMurray 2010).

En humanos este proceso comienza probablemente en el momento de la gastrulación, aproximadamente en el día E17-18, aunque el momento exacto se desconoce (Tang et al. 2016). El origen de las células germinales primordiales humanas (hPGCs) tampoco ha sido demostrado. Clásicamente, se ha aceptado que su origen podría estar en el epiblasto posterior de igual manera que en ratón, pero estudios recientes en primates (*Macaca fascicularis*) indican que la especificación podría ocurrir en el amnios temprano. Ambas posibilidades son factibles postulándose un posible origen dual de las hPGCs (Sasaki et al. 2016; Kobayashi and Surani 2018). En relación con el mecanismo molecular, existen procesos conservados como la participación de BLIMP1 y TFA2PC, así como la importancia de los factores de pluripotencia NANOG y OCT4, siendo el principal mecanismo diferenciador la función crucial de SOX17 (junto a BLIMP1 y TFA2PC) en la especificación hacia hPGC (Irie et al. 2015). En relación con los factores de pluripotencia, las hPGCs no expresan *SOX2* y sí *KLF4*, mientras que en mPGCs sucede al revés (Durcova-Hills et al. 2008; Campolo et al. 2013; Hackett et al. 2017). La acción conjunta de estos factores dará lugar a una población de hPGCs que se observan por primera vez durante la cuarta semana (~E24) en la pared del saco vitelino cerca del alantoides, ubicación equivalente a la de las mPGCs en el E7,25 (Figura I).

Tras su especificación, las PGCs migran hacia las crestas genitales, futuras gónadas, colonizándolas en torno al E37 en humano (E11,5 en ratón, Figura I). Durante la migración las PGCs proliferan activamente y eliminan la memoria somática mediante una reprogramación epigenética global que incluye la demetilación del DNA, la eliminación de la impronta genética, la reactivación del cromosoma X y la reorganización de las modificaciones de la cromatina (Tang et al. 2016). De esta manera se reprimen genes somáticos y se activan genes específicos de células germinales que preparan a las PGCs para iniciar la gametogénesis. En la sexta semana del desarrollo embrionario (E11,5 en ratón) comienza la diferenciación sexual de las gónadas (Figura I). En los embriones masculinos (XY) la expresión del gen *Sry*, codificado en el cromosoma Y, desencadena la diferenciación masculina hacia células de Sertoli y la formación de los testículos (Sinclair et al. 1990;

Miyawaki et al. 2020). En los femeninos (XX), la ausencia de este gen provoca una cascada de expresión génica específica que da lugar a la diferenciación hacia células de la granulosa y el desarrollo ovárico (Mclaren 1998; Griswold 2016). A partir de este momento, las PGCs dejarán de ser consideradas como tal siguiendo rutas de diferenciación distintas en función de si son XY o XX (Saitou and Yamaji 2012).

2.2. Oogénesis

En los embriones femeninos (XX) la oogénesis comienza con la proliferación de las PGCs y su diferenciación a oogonias en las crestas genitales. El número de divisiones mitóticas que ocurren es limitado (~20 divisiones) y se producen con una citocinesis incompleta que da lugar a la formación de nidos de oogonias rodeados por células somáticas (Figura V) (Jagarlamudi and Rajkovic 2012). En torno a la novena semana (E13,5 en ratón) estas oogonias entran en meiosis convirtiéndose en oocitos primarios. La entrada en meiosis se produce gracias a la represión de los genes de pluripotencia y a la diferenciación promovida por el ácido retinoico (RA, *Retinoic Acid*) producido en las gónadas. El RA induce la expresión del gen *Stra8* (Stimulated by Retinoic Acid 8) que a su vez promueve la expresión de genes esenciales para las primeras etapas de la meiosis, entre ellos los que codifican para proteínas del complejo sinaptonémico (SYCP1, SYCP2 y SYCP3) y proteínas de la maquinaria de recombinación meiótica (SPO11 y DMC1) (Baltus et al. 2006; Wear et al. 2016). Además, recientemente se ha demostrado que, en paralelo con la inducción de *Stra8*, el RA induce la expresión de la subunidad alpha-kleisina del complejo de cohesinas *Rec8*, esencial para los estadios tempranos de la meiosis, postulándose que el RA promueve la entrada en meiosis a través de al menos dos rutas independientes (Koubova et al. 2014).

Figura II. Representación esquemática de los principales procesos que ocurren durante la ovogénesis, desde la migración de las PGCs a la ovulación. PGCs: células germinales primordiales.

Una vez iniciada la meiosis, los oocitos primarios pasarán por los distintos estadios de la profase I quedando bloqueados al final de la misma en un estadio de quiescencia conocido como dictiata. Así, en el momento del nacimiento todos los oocitos se encuentran ya detenidos en este estadio (Figura I) (Slizynski 1957). En paralelo a la profase I se produce la rotura de los nidos de oocitos los cuales se rodean de una capa de células de la pregranulosa (células somáticas planas), dando lugar a los folículos primordiales (Figura II). Durante estos procesos ocurre una apoptosis masiva de oocitos y solo aquellos que sobreviven formarán un folículo (Pepling and Spradling 2001). Aunque diferentes estudios han descrito la presencia de células en el ovario con capacidad de generar nuevos oocitos cuando son trasplantadas bajo ciertas condiciones experimentales muy específicas (Johnson et al. 2004; Niikura et al. 2009), no existen evidencias de que *in vivo* estas células sean capaces de contribuir activamente al aumento de la reserva folicular (Lei and Spradling 2013; Zhang et al. 2014; Guo et al. 2016), aceptándose por tanto que el conjunto de folículos primordiales generado durante la etapa embrionaria representa la reserva ovárica disponible para toda la vida reproductiva femenina (De Felici et al. 2005; Wang et al. 2017a).

Los folículos primordiales se mantienen quiescentes hasta su activación, momento en el que pasan a considerarse folículos en crecimiento y comienza la foliculogénesis. Esta activación se produce por la influencia de numerosos factores y ocurre de forma continua desde el nacimiento hasta la menopausia. El primer paso tras la activación es la transición de folículo primordial a folículo primario que se caracteriza por cambios morfológicos en las células de la granulosa (Figura II). Estas células cambian de forma plana a forma cuboidal y se dividen mitóticamente formando múltiples capas que rodean al oocito. Además, durante esta transición los oocitos aumentan de tamaño (Pedersen and Peters 1968). En el desarrollo hacia folículo secundario, el oocito continúa creciendo, las células de la granulosa proliferando y aparece una capa externa adicional de células formando la teca (Figura II). Estos estadios del desarrollo folicular son independientes de las gonadotropinas FSH y LH (*Luteinizing hormone*). Sin embargo, los siguientes estadios son dependientes de las mismas (McGee and Hsueh 2000; Edson et al. 2009) y por ello los folículos activados antes de la pubertad son eliminados (atresia folicular) debido a la ausencia de las señales gonadotrópicas necesarias para su progresión (Peters 1969; McGee et al. 1998). Tras la pubertad y de manera cíclica se produce un aumento de FSH que desencadena los eventos posteriores. En primer lugar, en los folículos se forman cavidades llenas de fluido que acaban uniéndose en una sola denominada *antrum*, dando lugar a los denominados folículos antrales. La mayor parte de estos folículos sufrirán atresia y solo una parte de ellos (folículos dominantes) continuarán el desarrollo hacia el estadio preovulatorio (Figura II). El número de folículos capaces de progresar en la maduración varía con la especie, siendo normalmente solo uno en humanos (entre 2 y 12 en ratón). Los folículos dominantes responden al aumento de LH reanudando la meiosis y expulsando el oocito a los oviductos (ovulación). Una vez que el oocito ha sido liberado, las células de la granulosa y de la teca finalizan su diferenciación formando el cuerpo lúteo (Figura II).

Tras el aumento de la LH, la reanudación de la meiosis puede identificarse morfológicamente por la disolución de la envuelta nuclear del oocito, proceso conocido como ruptura de la vesícula germinal (GVBD, *germinal vesicle breakdown*). En este momento se lleva a cabo la primera división meiótica con una división asimétrica del citoplasma que da lugar a la extrusión del primer corpúsculo polar (Figura I). El inicio de la segunda división meiótica se produce de forma inmediata aunque su finalización y la liberación del segundo

corpúsculo polar solo se producirá si existe fertilización (Figura I). Por tanto, las células liberadas en la ovulación no son óvulos si no oocitos bloqueados en la segunda división meiótica (McGee and Hsueh 2000; Edson et al. 2009; Rimon-Dahari et al. 2016).

2.3. Espermatogénesis

En los embriones masculinos (XY), una vez que las PGCs alcanzan las crestas genitales e interaccionan con las células somáticas (Sertoli, Leydig y mioides), la determinación sexual finaliza y se forman los testículos embrionarios. En ellos, las PGCs junto a las células de Sertoli formarán los cordones seminíferos, futuros túbulos seminíferos (Cool and Capel 2009). En este punto, las PGCs adquieren características morfológicas diferenciales pasando a llamarse gonocitos o proespermatogonias. Estas células llevan a cabo un elevado número de divisiones mitóticas para a continuación entrar en un bloqueo mitótico en G_0/G_1 (~E49 en humanos; ~E13,5 en ratón) en el que se mantendrán hasta después del nacimiento (McLaren and Southee 1997; De Rooij 1998). Hacia la semana 8-12 post-parto (5 días post-parto -dpp- en ratón) estas proespermatogonias retoman la proliferación y una parte migra a la base de los túbulos seminíferos donde encontrarán el nicho apropiado para convertirse en células madre espermatogoniales (SSCs, *Spermatogonial Stem Cells*) las cuales mantienen la espermatogénesis de forma continua desde la pubertad (de Rooij 1973; Spradling et al. 2001; Nikolic et al. 2016; Griswold 2016). Estas SSCs, también conocidas como espermatogonias A_{single} (A_s) en ratón, son capaces de auto-renovarse para mantener el pool de células madre, pero también de comenzar la diferenciación hacia espermatozoide. Para ello, en primer lugar, las SSCs llevan a cabo una mitosis con citocinesis incompleta que da lugar a una cadena de espermatogonias conectadas (A_{aligned} ; A_{al}) que por divisiones sucesivas crece alcanzando las 8, 16 y, en ocasiones hasta 32 espermatogonias en ratón. Hasta este punto, las espermatogonias todavía se consideran indiferenciadas. Mediante sucesivas divisiones mitóticas estas espermatogonias avanzarán hacia los estadios más diferenciados de espermatogonia A (A_1 , A_2 , A_3 y A_4), espermatogonia intermedia (In) y finalmente espermatogonia B. Todas estas divisiones de amplificación también se producen con una citocinesis incompleta dando lugar a largas cadenas sincitiales de células conectadas por conexiones intercelulares (Weber and Russell 1987). A continuación, las espermatogonias B se dividirán e iniciarán la meiosis dando lugar a los espermátocitos primarios. Finalmente, tras dos rondas de división meiótica se formarán las espermátidas redondas haploides que se diferenciarán a espermatozoide maduro mediante un proceso denominado espermiogénesis. Gracias a esta sucesión de divisiones mitóticas y meióticas, a partir de un *pool* reducido de células madre se obtiene un número muy elevado de espermatozoides (de Rooij 2001; Oatley and Brinster 2008; Fayomi and Orwig 2018). En humanos, el proceso es similar pero en este caso las espermatogonias A se denominan $A_{\text{dark}}/A_{\text{pale}}$ y el número de divisiones de amplificación desde las SSCs hasta entrar en meiosis es menor, lo cual se ve compensado en parte por un mayor *pool* de SSCs (Fayomi and Orwig 2018).

La espermatogénesis tiene lugar de manera continua e ininterrumpida en los túbulos seminíferos tanto de humano como de ratón. El proceso de maduración completo desde espermatogonia indiferenciada hasta espermatozoide maduro tiene una duración de ~35 días en ratón (72 días en humano). Sin embargo, cada 8,6 días (16 días en humano) el proceso comienza de nuevo solapándose con el anterior. Las células que inician el nuevo ciclo lo hacen en la base de los túbulos desplazando a las más avanzadas en la diferenciación (procedentes de ciclos anteriores) hacia el lumen de los túbulos. En este desplazamiento los espermátocitos

deben atravesar las uniones estrechas existentes entre las células de Sertoli (barrera hemato-testicular) sin alterar su integridad lo que requiere una importante reorganización de las mismas (Smith and Braun 2012). Como consecuencia del solapamiento de los ciclos, en los túbulos seminíferos coexisten agrupaciones de tipos celulares en diferentes estadios de desarrollo. Este proceso ocurre de forma altamente coordinada y repetida en un programa conocido como ciclo del epitelio seminífero (Figura III). En ratón, el análisis de cortes histológicos de túbulos seminíferos ha permitido caracterizar 12 asociaciones celulares diferentes que determinan los estadios del ciclo epitelial seminífero (estadios I-XII). Esta clasificación resulta muy útil en la determinación del estadio en el que se produce el bloqueo en los mutantes meióticos (Oakberg 1956; Clermont 1972; Muciaccia et al. 2013; Ernst et al. 2019).

Figura III. Representación esquemática del ciclo del epitelio seminífero en ratón. Se representan las células presentes en cada estadio, desde la base al lumen de los túbulos seminíferos. A, In y B: espermatogonia A, Intermedia y B respectivamente. pL, L, Z, P, D: espermatocitos en preLeptotena, Leptotena, Zigotena, Paquitena y Diplotena, respectivamente. M: divisiones meióticas. 1-8: espermatidas redondas. 9-16: espermatidas alargadas.

Al igual que en la oogenesis, el inicio de la meiosis masculina en el ratón está mediado por la acción del RA (Griswold et al. 2012). Antes del nacimiento, la actividad de la enzima CYP26B1 en las células de Sertoli inmaduras degrada el RA impidiendo su acumulación y, por tanto, la entrada prematura en meiosis. Tras el nacimiento, se reprime la expresión del gen *Cyp26b1* en las gónadas masculinas y en respuesta a la acumulación de RA las espermatogonias comienzan a expresar *Stra8*, induciendo el inicio de la meiosis de forma regular gracias al aumento cíclico del RA (Koubova et al. 2006; Bowles et al. 2006; Griswold 2016). Clásicamente se ha aceptado que el balance entre la síntesis y la degradación del RA durante el desarrollo del sistema reproductivo determina la inducción de *Stra8* en el momento apropiado en cada sexo lo que produce la entrada en meiosis. Sin embargo, este modelo ha sido cuestionado dado que los ratones deficientes para la síntesis de RA (*Raldh2*^{-/-}) presentan expresión de *Stra8* en los ovarios fetales y, además,

la inhibición de CYP26B1 en testículo induce la expresión de *Stra8* pese a la ausencia de RA (Kumar et al. 2011). En base a estos resultados, se ha propuesto la existencia de algún mecanismo alternativo, todavía desconocido, en la inducción de la entrada en meiosis (Kumar et al. 2011; Griswold et al. 2012).

Por último, cabe destacar que durante la espermatogénesis el RA es producido además de por las células de Sertoli por los espermatocitos primarios en estadio de paquitena y diplotena y es esencial no solo para el inicio de la meiosis, sino también para la diferenciación de las espermatogonias (transición A-A1), la formación de la barrera hematotesticular, la espermiogénesis y la espermiación. Otro metabolito fundamental para la espermatogénesis es la testosterona producida por las células de Leydig y cuya función está relacionada con la regulación del final de la meiosis, el establecimiento de la barrera hematotesticular y la espermiación (Raverdeau et al. 2012; Jauregui et al. 2018).

3. Meiosis

La meiosis es un tipo de división celular especializada característica de los organismos con reproducción sexual. A diferencia de la mitosis donde a partir de una célula diploide ($2n$) se obtienen dos células idénticas a ella, la meiosis es una división reduccional en la cual se producen dos rondas de segregación cromosómica tras una única replicación del DNA generando los gametos haploides (n). Como hemos comentado anteriormente, esta reducción de la ploidía es esencial durante la gametogénesis para asegurar que, tras la fertilización, se obtenga el cigoto diploide ($2n$) capaz de dar lugar a un nuevo individuo. Durante la primera división meiótica se va a producir la segregación de los cromosomas homólogos (materno y paterno) y en la segunda la segregación de las cromátidas hermanas. Mientras que la segunda división meiótica se asemeja a la mitosis, la primera división reduccional es una adquisición exclusiva de la meiosis. Por ello se cree que, muy probablemente, la meiosis evolucionó a partir de la mitosis adquiriendo sus procesos característicos y diferenciales. Estos incluyen el apareamiento/sinapsis de cromosomas homólogos, la recombinación entre cromátidas no hermanas, la ausencia de segregación de las cromátidas hermanas durante la primera división meiótica y, por último, la ausencia de replicación del DNA previa a la segunda división meiótica (Wilkins and Holliday 2009).

3.1. Profase I

Tras la replicación del DNA comienza la profase I meiótica, la etapa de mayor duración de la meiosis y en la que se producen muchos de sus eventos característicos como son la formación de roturas de doble hebra (DSBs, Double Strand Breaks) programadas y su reparación por recombinación homóloga, el apareamiento y sinapsis de los cromosomas homólogos para formar los bivalentes, así como el remodelamiento de estos bivalentes. El objetivo principal de esta fase es establecer uniones físicas (quiasmas) entre los homólogos a través de la reparación por recombinación homóloga de las roturas del DNA usando como molde el cromosoma homólogo. Estas uniones son esenciales para asegurar la correcta segregación durante la primera división meiótica y evitar la formación de gametos aneuploides.

La profase I puede dividirse en cinco fases distinguibles citológicamente atendiendo al comportamiento de los cromosomas y el grado de ensamblaje del complejo sinaptonémico (SC, Synaptonemal Complex) (Figura VI). El SC es un complejo proteico específico de la meiosis que se forma entre los cromosomas

homólogos manteniéndolos físicamente unidos (sinapsis) y actuando como andamiaje sobre el cual ocurre la recombinación meiótica. El SC presenta una estructura tripartita compuesta por dos elementos axiales paralelos (AEs, Axial Elements) y un elemento central (CE, Central Element) unidos por filamentos transversales (TFs, Transverse Filaments). Una vez establecida la sinapsis los AEs pasan a denominarse elementos laterales (LEs, Lateral Elements). Esta estructura se encuentra altamente conservada en los organismos con reproducción sexual (Gillies 1975; Fraune et al. 2015). En mamíferos se han identificado 8 proteínas del SC hasta la fecha: SYCP2 y SYCP3 en los LEs, SYCP1 en los TFs y SYCE1/2/3, TEX12 y SIX6OS1 en el CE; esta última recientemente identificada y caracterizada en nuestro laboratorio (Meuwissen et al. 1992; Lammers et al. 1994; Offenberg et al. 1998; Costa et al. 2005; Hamer et al. 2006; Schramm et al. 2011; Gomez-H et al. 2016). Mediante inmunofluorescencia con anticuerpos frente a estas proteínas (principalmente de los AEs/LEs) es posible seguir la progresión de la meiosis e identificar con facilidad los distintos estadios de la profase I: leptotena, zigotena, paquitena, diplotena y diacinesis (Figura VI) (Bolcun-Filas and Schimenti 2012; Bolcun-Filas and Handel 2018).

Figura IV. Representación esquemática de los eventos de la profase I en el contexto del SC y la evolución de los RNs a lo largo de la misma. Adaptado de (Bolcun-Filas and Handel 2018).

Al inicio de la profase I, en leptotena, el DNA comienza a condensarse en forma de bucles y los AEs empiezan a ensamblarse simultáneamente con los complejos de cohesinas, formando los ejes cromosómicos (Figura IV). Estos ejes son los encargados de mantener las cromátidas hermanas unidas y participan en la formación de los *loops* en la cromatina sirviendo como puntos de anclaje. Los complejos de cohesinas son estructuras proteicas en forma de anillo que abrazan las cromátidas hermanas manteniéndolas unidas

(cohesión) y son esenciales para el ensamblaje de los AEs, la reparación de DSBs y la recombinación meiótica (Bannister et al. 2004; Novak et al. 2008; Herrán et al. 2011; Llano et al. 2012). En este estadio, los AEs y los complejos de cohesinas se ensamblan en pequeños fragmentos que van alargándose mientras las células progresan hacia el siguiente estadio. Además, durante leptotena se generan DSBs programadas en el DNA de los meiocitos por acción de la topoisomerasa SPO11 (Figura IV) (Keeney et al. 1997; Baudat et al. 2000).

La formación de las DSBs determina el comienzo de la recombinación meiótica (explicada en detalle en la sección 4). Para que esta se produzca los cromosomas deben reconocer a su homólogo y aparearse para a continuación, establecer una unión íntima entre sí, la sinapsis. Aunque existe apareamiento previo a la formación de DSBs, en la mayoría de los organismos, incluido el ratón, la formación de las DSBs es fundamental para el apareamiento y la sinapsis (Boateng et al. 2013). De hecho, en ausencia de SPO11 los cromosomas no aparean correctamente y se produce una sinapsis aberrante entre cromosomas no homólogos (Baudat et al. 2000). Además, en leptotena, se producen fuertes movimientos cromosómicos y una remodelación de la envuelta nuclear que da lugar a la formación de una estructura en ramillete denominada *bouquet* durante la transición hacia zigotena. Esta estructura es el resultado de la unión de los cromosomas a través de sus telómeros a una región muy delimitada de la envuelta nuclear (Scherthan et al. 1996; Lee et al. 2015). Todos estos eventos facilitan el apareamiento de los cromosomas homólogos y el comienzo de la sinapsis.

La sinapsis comienza en las regiones subteloméricas y marca el inicio de la zigotena. Durante este estadio la sinapsis se va extendiendo gradualmente a lo largo de los ejes cromosómicos a modo de cremallera. Para ello, entre los AEs paralelos (LEs tras la sinapsis) se ensambla el CE que se ancla a los AEs a través de numerosos TFs, formando la estructura tripartita característica del SC (Figura IV) (Heyting 1996). El avance de la sinapsis, además, se ve facilitado por la carga de proteínas de recombinación que forman los nódulos de recombinación (RNs, Recombination Nodules) tempranos (Figura IV). Diferentes proteínas de recombinación se irán asociando a los RNs dando lugar a los RNs de transición y determinando, hacia el final de zigotena, cuáles seguirán la vía de los *crossovers* (CO) y cuáles la de los *non-crossovers* (NCO).

El estadio de paquitena se caracteriza por una sinapsis completa de los cromosomas homólogos y una cromatina altamente condensada (Figura IV). Hacia el final de paquitena todas las DSBs han sido reparadas y se forman los RNs tardíos que median la resolución de los intermediarios de recombinación como CO o NCO. La excepción a la sinapsis completa de los homólogos en esta fase se observa en los cromosomas sexuales X e Y en machos. En mamíferos, estos cromosomas son muy distintos en estructura y contenido génico y solo muestran homología en una pequeña región conocida como región pseudoautosómica (PAR, PseudoAutosomal Region). Esta región es la única capaz de sinapsar (Figura IV) y en ella es donde tiene lugar la formación del quiasma que asegura la correcta segregación de los cromosomas X e Y. Como consecuencia de la asinapsis de la mayor parte de los cromosomas sexuales se produce su inactivación transcripcional o silenciamiento (MSCI, Meiotic Sex Chromosome Inactivation). En leptotena la fosforilación de la histona H2AX (γ H2AX) tiene un papel fundamental en los eventos tempranos de la reparación de DSBs (ver sección 4.2.1). Además, en paquitena existe otra oleada de fosforilación de H2AX (mediada por ATR y BRCA1) restringida a la cromatina de los cromosomas no sinapsados y que es la responsable del inicio la MSCI

(Mahadevaiah et al. 2001; Turner et al. 2005). Durante el inicio de la MSCI, los cromosomas X e Y sufren modificaciones post-traduccionales como la ubiquitinación de la histona H2A, la desacetilación de H3 y H4, la dimetilación de H3 y la sumoilación de histonas y otros sustratos aún por identificar (Baarends et al. 1999; Rogers et al. 2004; Khalil et al. 2004; Vigodner and Morris 2005). Todas estas modificaciones podrían participar en el mantenimiento de la MSCI ya que se mantienen asociadas a los cromosomas sexuales durante las divisiones meióticas y la espermiogénesis, a pesar de que en este estadio H2AX ya ha sido defosforilada, y ponen de manifiesto la importancia de la regulación de las modificaciones post-traduccionales en la progresión de la meiosis. Citológicamente, la MSCI produce una elevada condensación de la cromatina que da lugar a una estructura compacta, perfectamente distinguible en el núcleo, conocida como *XY body* o *sex body* (Figura IV) (Cloutier and Turner 2010).

Por último, en diplotena el CE del SC empieza a desensamblarse (Figura IV) y se produce un remodelamiento de los bivalentes. Este proceso, que se prolonga durante la diacinesis, incluye un aumento de la condensación del DNA y la localización en los cromosomas de proteínas implicadas en la regulación de la liberación de los complejos de cohesinas durante la transición metafase-anafase I (Marston 2015). Así, al final de la profase I los cromosomas están unidos únicamente por los quiasmas y preparados para iniciar la primera división meiótica (Figura IV).

3.2. Divisiones meióticas

Al finalizar la profase I meiótica se van a producir dos divisiones meióticas consecutivas que culminarán con la formación de los gametos haploides. La primera es una división reduccional donde segregan los cromosomas homólogos. La segunda es una división ecuacional, similar a la mitosis, donde segregan las cromátidas hermanas. Esta segregación en dos pasos es posible gracias a una estricta regulación espacio-temporal de la liberación de los complejos de cohesinas, así como por el cambio de la geometría de los cinetocoros en relación con su orientación a los polos del huso (mono-orientación vs bi-orientación de los cinetocoros hermanos). En la transición metafase-anafase I solamente se van a eliminar los complejos de los brazos cromosómicos manteniéndose los de los centrómeros lo que permite la segregación de los homólogos. Los complejos de cohesinas centroméricos van a mantener las cromátidas hermanas unidas hasta la transición metafase-anafase II cuando serán eliminados permitiendo la segregación de las mismas (Llano et al. 2008; Bolcun-Filas and Handel 2018). Los errores en la segregación cromosómica durante estos procesos conllevan la aparición de aneuploidías en los gametos que pueden producir anomalías genéticas en la descendencia (como el Síndrome de Down), abortos e infertilidad (Petronczki et al. 2003).

Esta liberación secuencial está mediada por dos oleadas de activación de Separasa, una proteasa que corta la subunidad alpha-kleisina de los complejos de cohesinas (REC8 o RAD21L, Buonomo et al. 2000; Kitajima et al. 2003; Kudo et al. 2009). Para asegurar la correcta segregación durante las divisiones meióticas, la activación de Separasa está estrictamente regulada por dos mecanismos de inhibición mediados por Securina y el complejo CDK1/Ciclina B1 (Stemmann et al. 2001; Hellmuth et al. 2015). En la primera división meiótica, cuando los bivalentes se encuentran alineados en la placa metafásica y los cinetocoros hermanos de cada homólogo orientados hacia el mismo polo (orientación sintética o mono-orientación), se satisface el *checkpoint* de ensamblaje del huso (SAC, Spindle Assembly Checkpoint) dando lugar a la activación de un

macrocomplejo proteico con actividad E3 ubiquitin ligasa denominado APC/C (Anaphase promoting complex/cyclosome). Este complejo va a ubiquitinar los inhibidores de Separasa marcándolos para su degradación por el proteasoma y permitiendo la activación de Separasa (Herbert et al. 2003). Una vez activa, Separasa va a cortar los complejos de cohesinas de los brazos cromosómicos, permitiendo la resolución de los quiasmas y la segregación de los homólogos. Durante esta primera activación de Separasa los complejos de cohesinas centroméricos deben estar protegidos para no ser cortados, evitando la segregación de las cromátidas hermanas. Las encargadas de esta protección son una familia de proteínas altamente conservadas denominadas shugoshinas. En mamíferos, existen dos parálogos de estas proteínas: SGOL1, esencial para el mantenimiento de la cohesión centromérica en mitosis (Kitajima et al. 2006) y SGOL2, encargada de esta protección en meiosis. En el ratón la ausencia de SGOL2 provoca infertilidad debido a una separación prematura de las cromátidas hermanas durante anafase I que da lugar a la formación de gametos aneuploides (Lee et al. 2008b; Llano et al. 2008).

Durante la segunda división meiótica los cromosomas se alinean en la placa metafásica con cada cinetocoro hermano orientado hacia un polo (orientación anfitélica). La tensión ejercida sobre los centrómeros produce una redistribución de SGOL2 del centrómero hacia los cinetocoros dejando los complejos de cohesinas centroméricos accesibles para su corte durante la segunda activación de Separasa en la transición metafase-anafase II (Gómez et al. 2007; Lee et al. 2008b). Como resultado de este proceso se generan los gametos haploides con una única copia de cada cromosoma.

En mamíferos el número de gametos viables obtenidos tras la meiosis difiere entre machos y hembras. En machos, a partir de cada espermatogonia que entra en meiosis se producen 4 gametos haploides que se diferenciarán hacia espermatozoide maduro durante la espermiogénesis y espermiación. Sin embargo, en hembras solo se forma un oocito funcional tras la meiosis. Esto se debe a una elevada asimetría en las divisiones citoplasmáticas donde el futuro óvulo (oocito fertilizable) retiene la mayor parte del citoplasma mientras que los corpúsculos polares reciben una mínima cantidad del mismo (Figura I) (Bolcun-Filas and Handel 2018).

4. Recombinación meiótica

La recombinación meiótica es el proceso por el cual los cromosomas homólogos intercambian material genético incrementando la diversidad genética en los organismos con reproducción sexual. Durante la profase I, este intercambio entre los cromosomas homólogos paterno y materno genera las nuevas combinaciones de alelos que se transmitirán a la siguiente generación. Por tanto, este proceso juega un papel fundamental en la evolución de los genomas proporcionando el material de partida sobre el que actuará la selección natural (Coop and Przeworski 2007). Sin embargo, el principal objetivo de la recombinación meiótica es establecer las conexiones físicas entre los homólogos necesarias para el apareamiento, la sinapsis y para la correcta segregación durante la primera división meiótica (Hassold and Hunt 2001).

La recombinación meiótica puede dar lugar a la formación de dos productos finales, los COs y los NCO. En los NCO el intercambio de material genético entre homólogos (conocido como conversión génica) es

unidireccional, afecta a pequeños fragmentos y, por tanto, su aportación a la generación de diversidad genética es reducida (Figura V). Al contrario, en los COs se produce un intercambio recíproco de grandes segmentos de material genético entre los cromosomas parentales contribuyendo en gran medida a la generación de diversidad genética (Figura V). Además, como resultado de los COs se establecen las uniones físicas (quiasmas) que permiten la correcta segregación de los homólogos. Los quiasmas, junto con la cohesión de los brazos cromosómicos en las regiones distales a ellos, permiten la biorientación de los homólogos y evitan la segregación prematura ya que se oponen a la fuerza ejercida por el huso permitiendo la estabilización de todas las uniones antes de la segregación (Miyazaki and Orr-Weaver 1994; Buonomo et al. 2000). Por su parte, los NCOs son necesarios para el apareamiento/sinapsis entre homólogos durante las primeras etapas de la profase de mamíferos (Baudat et al. 2013). Por lo tanto, la recombinación representa un proceso crucial durante la meiosis del cual dependen eventos tan importantes como el apareamiento, la sinapsis y la correcta segregación cromosómica.

4.1. Inicio de la recombinación

La recombinación comienza con la formación programada de DSBs a lo largo de todo el genoma. Estas DSBs se forman en sitios específicos denominados *hotspots* (puntos calientes) que, en ratón y humano, están determinados por la histona metiltransferasa PRDM9 (Figura V) (Baudat et al. 2010; Myers et al. 2010; Parvanov et al. 2010). Al comienzo de leptotena, PRDM9 se une a secuencias específicas del DNA a través de su dominio zinc finger y cataliza la trimetilación de las lisinas 4 y 36 de la histona H3 (H3K4me3 y H3K36me3) en los nucleosomas adyacentes (Hayashi et al. 2005; Grey et al. 2011; Powers et al. 2016). Estas marcas epigenéticas promueven el reclutamiento de la maquinaria de formación de DSBs aunque el mecanismo preciso se desconoce.

La formación de las DSBs meióticas está catalizada por SPO11, una proteína con alta similitud con la subunidad A de la DNA topoisomerasa VI (TopoVIA). En humano y ratón existen dos isoformas de SPO11, SPO11 α y SPO11 β , siendo SPO11 β la responsable de la introducción de la mayoría de las DSBs meióticas en leptotena. La expresión de SPO11 α comienza en zigotena y es esencial para el apareamiento y la recombinación de los cromosomas XY en ratón (Bellani et al. 2010; Kauppi et al. 2011). Recientemente, se ha identificado una familia conservada de proteínas con alta similitud estructural con la subunidad B de las topoisomerasas TopoVI (TopoVIB). En ratón, TOPOVIBL (TOPOVIB-Like) interacciona con SPO11 β y es necesaria para la formación de las DSBs, demostrando que la inducción de DSBs en meiosis está catalizada por un complejo formado al menos por SPO11 y TOPOVIBL (Figura V) (Robert et al. 2016). Además, existen proteínas adicionales que participan en la formación de DSBs. En ratón, MEI4, REC114 e IHO1 (Interactor of HORMAD1) forman un complejo, reclutado a los ejes cromosómicos a través de HORMAD1, que interacciona con SPO11 a través de proteínas intermediarias (Figura V). Este complejo participa en el reclutamiento de SPO11 (y probablemente TOPOVIBL) a los ejes cromosómicos donde se forman las DSBs (Kumar et al. 2010; Kumar et al. 2015; Stanzione et al. 2016).

La cantidad de DSBs meióticas que se generan es difícil de cuantificar, especialmente por su carácter transitorio. De forma general se ha utilizado el número de foci formados por las recombinasas RAD51 y DMC1 (ver más adelante) como la mejor estimación del número de DSBs, encontrándose en humano y ratón entre

200 y 400 foci por célula. La reducción en este número de DSBs provoca defectos en sinapsis, lo cual sugiere que es necesario un número mínimo de DSBs para el correcto apareamiento y sinapsis entre homólogos (Kauppi et al. 2013). Actualmente, mediante el empleo de ChIPseq se puede determinar con precisión el número y posición exacta de las DSBs en el genoma (Smagulova et al. 2011; Khil et al. 2012; Lange et al. 2016).

4.2. Reparación de las DSBs

El número de DSBs que se forman en cada célula es muy elevado y altamente variable (~32-37% de variación). Sin embargo, solamente 1 de cada 10 DSBs son procesadas como COs y su número final es relativamente constante en todas las células (~12% de variación) (Cole et al. 2012). Esto pone de manifiesto la importancia de una correcta regulación de la reparación de las DSBs que mantenga la homeostasis en la formación de COs y asegure la formación de al menos un CO por par de homólogos, permitiendo la correcta segregación durante la primera división meiótica (Baudat et al. 2013). La reparación de las DSBs se produce por la acción de múltiples proteínas de recombinación que se ensamblan formando los RNs. Estos núcleos están localizados en el contexto de los ejes cromosómicos y el SC donde se detectan como foci discretos. Los componentes de los RNs van cambiando secuencialmente a lo largo del tiempo y en base a esta composición se clasifican en RNs tempranos, intermedios (o de transición) y tardíos (Figura V) (Bolcun-Filas and Handel 2018).

4.2.1. Procesamiento de las DSBs

En leptotena, una de las primeras señales detectadas tras la formación de las DSBs es la fosforilación de la histona H2AX en la serina 139 (γ H2AX) por la quinasa ATM (ataxia telangiectasia mutated), señal que activa la cascada de reparación posterior (Mahadevaiah et al. 2001; Bellani et al. 2005). El primer paso en el procesamiento de las DSBs es la liberación de los complejos de SPO11 del DNA mediante el corte endonucleolítico mediado por el complejo MRN (MRE11-RAD50-NBS1) (Neale et al. 2005; Zhang et al. 2020a). A continuación, tiene lugar una resección 5' $\rightarrow$ 3' que genera los extremos 3' de ssDNA (DNA de hebra simple, single-stranded DNA) que llevarán a cabo la búsqueda de las regiones de homología y la invasión durante la recombinación (Figura V). En *Saccharomyces cerevisiae* esta resección está mediada por Exo1 (Zakharyevich et al. 2010) y regulada por Tel1 (ortólogo de ATM). En ratón, mientras que la función de ATM está conservada, la contribución de EXO1 en este proceso es mínima, lo que sugiere la existencia de proteínas adicionales que actúen de forma redundante con EXO1 en la resección (Yamada et al. 2020).

Tras la resección de las DSBs, la proteína RPA (Replication Protein A) se une a los extremos 3' libres protegiéndolos de la degradación y evitando la formación de estructuras secundarias (Figura V) (Chen et al. 2013). RPA es un complejo heterotrimérico de unión a ssDNA formado por RPA1, RPA2 y RPA3 en el que RPA1 posee la mayor afinidad por el ssDNA gracias a que posee 3 dominios de unión conocidos como OB-folds (*Oligonucleotide-binding folds*) (Ribeiro et al. 2016). Seguidamente, RPA es desplazado rápidamente por las recombinasas RAD51 (Radiation sensitive 51) y DMC1 (disrupted meiotic cDNA 1; homólogo de RAD51 específico de meiosis), formando los RNs tempranos los cuales son detectables en leptotena como 200-400 foci discretos sobre los ejes cromosómicos (Figuras IV y V) (Barlow et al. 1997; Pittman et al. 1998; Yoshida

et al. 1998). El intercambio de RPA por RAD51 y DMC1 es un paso crítico durante la reparación de las DSBs que se cree que está mediado principalmente por BRCA2 (Breast cancer susceptibility 2) dada su interacción con RAD51, DMC1 y el DNA (Davies et al. 2001; Yang et al. 2002; Thorslund et al. 2007). De acuerdo con esto, la deficiencia de BRCA2 en ratón provoca defectos en recombinación meiótica debido a una importante reducción en la carga de RAD51 y DMC1 (Sharan et al. 2004). Recientemente, mediante aproximaciones bioquímicas se ha descrito que BRCA2 interacciona con DSS1 quien, a su vez, interacciona con RPA reduciendo su afinidad por el ssDNA y postulándose como un factor crucial en el remplazamiento de RPA por RAD51 durante la recombinación homóloga (Zhao et al. 2015). Sin embargo, su función en meiosis no ha sido abordada. También se ha descrito interacción de RAD51 con BRCA1 en células mitóticas y meióticas (Scully et al. 1997) y su deficiencia en ratón produce defectos en la recombinación meiótica como consecuencia de la reducción de la carga de RAD51 (Xu et al. 2003). Por último, la ausencia de TEX15 (Testis expressed sequence 15) en ratón suprime la carga de las recombinasas en la espermatogénesis (Yang et al. 2008) siendo, por tanto, al menos tres los factores conocidos que participan en la carga de las recombinasas tras la formación de DSBs (Figura V).

Figura V. Representación esquemática de la ruta de recombinación meiótica en mamíferos. dHJ: uniones dobles de Holliday. sHJ: uniones simples de Holliday. SDSA: anillamiento de hebra dependiente de síntesis.

4.2.2. Invasión del cromosoma homólogo

Los nucleofilamentos cubiertos por RAD51 y DMC1 son los encargados de la búsqueda de homología y la invasión del cromosoma homólogo produciendo el desplazamiento de la hebra y generando una estructura conocida como *D-loop* (Figura V). Mediante el empleo de ChIP-Seq *in vivo* se han localizado con precisión los sitios de unión de RAD51, DMC1 y RPA determinándose que, en realidad, RAD51 y DMC1 tienen una

localización no solapante en los extremos de ssDNA durante la meiosis. DMC1 se une en las regiones próximas a la rotura y RAD51 se encuentra localizado en las más alejadas (Figura V). Además, en las regiones de los *D-loops* donde RPA se carga de nuevo, DMC1 co-localiza mejor con RPA que RAD51 sugiriendo que DMC1 sería la recombinasa *bona fide* que llevaría a cabo la invasión del cromosoma homólogo (Hinch et al. 2020).

La búsqueda de homología y la invasión mediada por RAD51 y DMC1 facilita que los homólogos se encuentren e interaccionen. Estas interacciones son posteriormente estabilizadas por la carga de las proteínas del SC durante la sinapsis en zigotena. A medida que los homólogos van sinapsando, RAD51 y DMC1 son reemplazadas por los componentes de los RNs intermedios, encargados del procesamiento de los intermediarios de recombinación hacia la vía de los COs o NCOs (Figura IV). Así, tras la invasión de la hebra y la formación de los *D-loops*, se produce la elongación del extremo 3' usando una hebra del cromosoma homólogo como molde. La hebra que no actúa como molde se conoce como hebra desplazada y en los *D-loops* se encuentra en forma de ssDNA (Figura V). Además, en el segundo extremo de las DSBs, es decir, aquel que ha sido reseccionado pero no ha llevado a cabo la invasión, también existe una región de ssDNA. Ambas regiones son protegidas por RPA junto con otras dos proteínas recientemente descritas que forman complejo entre sí e interaccionan con RPA, que son MEIOB (Meiosis specific with OB domains) y SPATA22 (Spermatogenesis Associated 22). Este trímero además de participar en la protección del ssDNA tras la resección, media la captura del segundo extremo por parte del homólogo, evento fundamental para la formación de las dobles uniones de Holliday (dHJ, Double Holliday Junctions) en la vía de los COs (Figura V) (La Salle et al. 2012; Luo et al. 2013; Xu et al. 2017).

4.2.3. Procesamiento de los intermediarios de recombinación

Además de RPA, MEIOB y SPATA22, un elevado número de proteínas están implicadas en la estabilización/procesamiento de los intermediarios de recombinación y en la regulación del destino final de las DSBs (CO o NCO). Entre ellas destacan BLM, TEX11, CNTD1, RNF212, HEI10 (también conocido como CCNB1IP1) y el complejo MSH4/MHS5 (complejo MutS γ , Figura V). Mientras que los pasos iniciales de la recombinación meiótica y la formación de COs están bien caracterizados, el conocimiento de los pasos intermedios está limitado en gran parte por la naturaleza dinámica los mismos y el carácter transitorio de los intermediarios de recombinación que se forman (Baudat et al. 2013; Hinch et al. 2020). En este ámbito, se ha postulado recientemente que la sumoilación y ubiquitinación tienen funciones antagónicas esenciales en la regulación y control de estos procesos (Rao et al. 2017). Estas modificaciones post-traduccionales se localizan en el contexto del SC en zigotena y paquitena donde son introducidas por E3 ligasas como RNF212 (E3-SUMO ligasa) y HEI10 (E3 Ubiquitin-ligasa) (Toby et al. 2003; Reynolds et al. 2013). Se ha propuesto que estas proteínas establecen un balance entre SUMO y Ubiquitina encargado de regular la renovación de proteínas de los RNs intermedios como el complejo MutS γ y TEX11 modulando la tasa de recombinación. Mientras RNF212 estabilizaría los RNs promoviendo su maduración hacia COs, la acción de HEI10 y el proteasoma los desestabilizaría promoviendo los NCOs (Qiao et al. 2014; Rao et al. 2017). Por tanto, la correcta regulación de la ubiquitinación y la sumoilación sería esencial para el éxito de la recombinación meiótica.

4.2.4. Resolución de los intermediarios de recombinación

Por último, el evento final de la recombinación meiótica es la resolución de los intermediarios de recombinación como CO o NCO. En la vía preferencial de formación de NCO el *D-loop* vuelve a anillar con su cromátida hermana y la hebra que invade conecta de nuevo con el segundo extremo perteneciente a su mismo cromosoma (Figura V). Este proceso se conoce como anillamiento de hebra dependiente de síntesis (SDSA, Synthesis Dependent Strand Annealing) y se produce en la mayor parte de las DSBs. En ocasiones, los NCO también pueden producirse por una disolución de las dHJ (Youds and Boulton 2011; Baudat et al. 2013). Los COs se generan por la resolución de las uniones de Holliday pudiendo producirse por dos mecanismos. En el primero y mayoritario, los COs se generan por la resolución de dHJ estando sujetos a un fenómeno de interferencia positiva (COs clase I). En ratón, esta vía está mediada por las proteínas *mismatch repair* MLH1/MLH3 (complejo MutL γ) y EXO1 (Figura V) (Edelmann et al. 1996; Baker et al. 1996; Lipkin et al. 2002; Wei et al. 2003). El segundo mecanismo supone la formación y resolución de uniones de Holliday simples (sHJ, Single Holliday Junctions) y estos CO no se ven sujetos al fenómeno interferencia (COs clase II). La resolución por esta vía está mediada por MUS81 y solo ocurre en el 10% de los COs (Figura V) (Holloway et al. 2008). Finalmente, en paquitena, todas las DSBs han sido reparadas y los intermediarios de recombinación han sido resueltos asegurando la existencia de al menos un CO por bivalente. Como se ha comentado anteriormente, la correcta regulación espacio-temporal de todos los elementos de esta ruta es esencial para el mantenimiento de la homeostasis de los COs que asegure una la correcta segregación durante metafase I y evite la aparición de aneuploidías, abortos e infertilidad (Petronczki et al. 2003; Baudat et al. 2013).

OBJETIVOS

Descifrar las causas de la infertilidad humana es de gran importancia dado que una elevada fracción de sus diagnósticos son idiopáticos con una base genética aún desconocida. Dada la imposibilidad de estudiar la gametogénesis en humanos, el empleo del ratón como modelo experimental representa la mejor aproximación para profundizar en este proceso. El presente trabajo aborda la identificación y caracterización funcional de dos genes candidatos de infertilidad humana mediante la generación y el análisis de modelos murinos. Esta aproximación, no solo permite establecer relaciones causales directas entre las variantes génicas y la infertilidad, sino que además profundiza en los procesos biológicos que gobiernan la gametogénesis. En base a ello, los objetivos concretos que nos planteamos en este trabajo fueron los siguientes:

1. Análisis funcional de la desubiquitinasa USP26 en la gametogénesis de ratón y su relación con la infertilidad masculina.
2. Identificación y caracterización de la causa genética subyacente a la insuficiencia ovárica primaria presente en un pedigrí familiar.
 - a. Validación de la variante S167L en *HSF2BP* como causante de la infertilidad familiar.
 - b. Disección genética del mecanismo molecular de actuación de HSF2BP.

MATERIAL Y MÉTODOS

1. Modelos murinos

1.1. Bienestar animal

Los ratones fueron estabulados siguiendo las regulaciones establecidas por la Unión Europea para el Servicio de Experimentación Animal (SEA) de la Universidad de Salamanca (USAL). Así, la estabulación se ha llevado a cabo en una instalación libre de patógenos específicos (*specific-pathogen free*, SPF) y con temperatura controlada, utilizando jaulas con ventilación individual, dieta estándar y un ciclo de luz-oscuridad de 12 horas. Los protocolos utilizados en el trabajo con ratones han sido aprobados por el Comité Ético de Experimentación Animal de la USAL. Siguiendo el principio de “Las tres Rs: reemplazar, reducir y refinar”, todos los esfuerzos se han dirigido a reducir el número de animales empleados, minimizar el sufrimiento y mejorar el bienestar de estos.

1.2. Generación de ratones modificados genéticamente mediante CRISPR/Cas9

Durante el desarrollo de esta tesis la tecnología CRISPR/Cas9 ha surgido como una potente alternativa a los métodos clásicos de generación de ratones modificados genéticamente como el *gene targeting*. Esta tecnología de edición del genoma tiene dos componentes principales: por un lado, una endonucleasa (Cas9), encargada de generar roturas dirigidas en el DNA. El otro componente son los RNAs guía (guías o sgRNAs) encargados de dirigir a la Cas9 de manera específica a su diana en el genoma. Una vez generadas, estas roturas pueden repararse mediante reparación de extremos no homólogos (NHEJ). Este es un mecanismo con alta tendencia a cometer errores, pudiendo generar inserciones o deleciones inespecíficas que den lugar a alelos mutantes. De manera alternativa, si añadimos al sistema un molde para la reparación, podemos generar inserciones, deleciones o mutaciones puntuales muy concretas. Este molde normalmente consiste en un DNA donador de hebra simple (*single-stranded donor oligonucleotide-ssODN*) que contiene las mutaciones a introducir flanqueadas por brazos de homología colindantes a la zona diana para favorecer la reparación por recombinación homóloga, un mecanismo de reparación de alta fidelidad (Yang et al. 2014). Los modelos *Usp26*^{21/21}, *Usp26*^{Cat/Cat}, *Hsf2bp*^{S167L/S167L}, *Hsf2bp*^{-/-}, *Brme1*^{Δ142-472/Δ142-472}, *Brme1*^{-/-}, *Spo11*^{-/-}, *Rnf212*^{-/-} and *Hei10*^{-/-} empleados en esta tesis han sido generados utilizando esta tecnología.

1.2.1. Diseño de sgRNAs

Los sgRNAs o RNAs guía realmente están formados por dos componentes, ambos imprescindibles: los RNAs CRISPR (crRNAs) y los crRNAs trans-activadores (tracrRNA). Los crRNAs son los responsables de la especificidad y los tracrRNAs intervienen en el procesamiento de los crRNAs actuando como andamiaje para la unión de ambos RNAs a la Cas9 y de este complejo al DNA para producir el corte dirigido (Wang et al. 2016). Por lo tanto, la parte a diseñar en cada modelo será el crRNA mientras que la secuencia del tracrRNA será siempre la misma. Desde la primera descripción del sistema CRISPR/Cas9 como herramienta de biología molecular (Jinek et al. 2012) hasta el día de hoy, el sistema ha sufrido una gran evolución, apareciendo diferentes herramientas de diseño que han ido mejorando las ya existentes. En este sentido, para el diseño de nuestros crRNAs, hemos utilizado dos herramientas:

- En primer lugar, el servidor crispr.mit.edu desarrollado por el Zhang Lab (Massachusetts Institute of Technology, MIT). Posteriormente, ellos mismos cerraron el servidor y recomendaron utilizar las nuevas y mejoradas fuentes.
- Desde ese momento el diseño se ha realizado usando el servidor de IDT (https://eu.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM).

Los crRNAs utilizados para los distintos modelos están agrupados en la Tabla I.

Tabla I. Lista de los crRNAs utilizados para la generación de los diferentes modelos de ratón. generados.

Modelo	crRNA ID	Secuencia
<i>Usp26</i>^{21/21}	sgRNA1	5'- TGGCCGAATACTACCTTGAG -3'
	sgRNA2	5'- GTGAGCCTAGCACAGTGCAC-3'
<i>Usp26</i>^{Cat/Cat}	sgRNA3	5'- AGGTGTTTCCCACATTAGGC-3'
	sgRNA4	5'- ATTAATGATTTATTCAACCA-3'
<i>Hsf2bp</i>^{S167L/S167L}	sgRNA1	5'-TCACAAAACCTCTCCATCGTC-3'
	sgRNA2	5'-ATTGGATGGGGATGTCAAGG-3'
<i>Hsf2bp</i>^{-/-}	sgRNA1	5'-TCACAAAACCTCTCCATCGTC-3'
	sgRNA2	5'-ATTGGATGGGGATGTCAAGG-3'
<i>Brme1</i>^{Δ142-472 / Δ142-472}	sgRNA3	5'-AACCTCAGGGACTCTCTCTG-3'
	sgRNA4	5'-GAAGTCTAGTTCCATTGCTG-3'
<i>Brme1</i>^{-/-}	sgRNA3	5'-AACCTCAGGGACTCTCTCTG-3'
<i>Spo11</i>^{-/-}	sgRNA5	5'-TATGTCTCTATGCAGATGCA-3'
	sgRNA6	5'- AACTGACAGCCAGCTCTTT-3'
<i>Rnf212</i>^{-/-}	sgRNA7	5'- ACCCACGTGAGACTCGCGCG-3'
	sgRNA8	5'- CCTCAAAGGTCCGCGTATTC-3'
<i>Hei10</i>^{-/-}	sgRNA9	5'- GAAAGGGTACTGTTGCAAGC-3'

1.2.2. Obtención de sgRNAs, Cas9 y ssODNs.

También la manera de obtener los distintos componentes del sistema CRISPR/Cas9 ha ido evolucionando con el tiempo. En nuestro caso, inicialmente los sgRNAs se produjeron mediante anillamiento de oligonucleótidos complementarios que codifican para los crRNAs (Figura VI).

Figura VI. Esquema representativo de los oligonucleótidos anillados para generar el crRNA. En negrita aparecen los extremos compatibles para clonar el fragmento en el vector pX330 digerido con BbsI y en azul las bases correspondientes a la secuencia específica de cada crRNA. La Guanina inicial en rojo (y su Citosina complementaria) aumenta la eficiencia de transcripción del RNA por parte de la RNA Pol I. De no existir en la propia secuencia del crRNA es recomendable añadirla. No afecta a la especificidad del sgRNA.

- T7-sgRNA-S: TAATACGACTCACTATAGGNNNNNNNNNNNNNNNNNNNNNNN donde la zona subrayada corresponde al promotor T7, común para todos los diseños, y las N la secuencia correspondiente a cada crRNA.
- T7-sgRNA-AS: AAAAGCACCGACTCGGTGCC.

1.2.3. Microinyección del sistema CRISPR/Cas9 en cigotos

Para la microinyección se preparó una mezcla de los diferentes componentes en las siguientes concentraciones:

- Microinyección de RNAs procedentes de transcripción *in vitro*: 50-100 ng/μl de Cas9, 30ng/μl de cada sgRNA (cr+tracrRNA sintetizados como una única molécula) y 30 ng/μl de ssODN en los casos que se requiriera.
- Microinyección de RNAs y proteína sintéticos: 30 ng/μl de Cas9 recombinante, 20ng/μl de cada sgRNA anillado y 10ng/μl de ssODN en los casos que se requiriera.

Esta mezcla fue microinyectada en cigotos B6/CBA F2 (híbridos entre cepas C57BL/6J y CBA/J) como se ha descrito previamente (Singh et al. 2015). Este procedimiento fue llevado a cabo por la Unidad de Transgénicos de la Universidad de Salamanca.

Tabla II. Lista de los ssODNs utilizados para la generación de los diferentes modelos de ratón generados. Las letras en rojo representan las mutaciones puntuales deseadas en los diferentes modelos. En negrita se resaltan las mutaciones introducidas para eliminar las secuencias PAM. Con esto, sin alterar la secuencia proteica, se evita que el sistema CRISPR/Cas9 vuelva a cortar una vez que se han introducido las mutaciones deseadas.

Modelo	Secuencia ssODN
Hsf2bp ^{S167L/S167L}	5'CTTTGGAAAGATGTGACAGTTCTATCTTTTTATCTTTCAGGACAAAGCATTGAAGTT TTCAACATAACTGG AC AGACGATGGAGAGTTTTGTGAAGT T ATTGGATGGGGATGT CAAGGA AG TTGATTCTGATGAAAATCAATTTGTCTTTGCACTGGCTGGAATTGTAACA AGTAGGTAACTTTTAGATACAGCGCT3'
Brme1 ^{Δ142-472/} Δ142-472	5'CTTCAGAGTGCTTGCTTATTGAAGGCCAGGACTGAATCTCTTTTTCCACAGGAAAC AAGGCCAGAGCTGGGAGCCCTCAAAGCAGCCAGCCAGCCACAGGCAATGGAACTAG ACTTCTGCCTGACAGCCAGATACAGGATGCCCTGGATGCCACTAACATGGAGCAGGT AAGAGCTTTCTGTACTCAAATGTACACCC3'
Spo11 ^{-/-}	5'GTTTCCTGCGGTATGTGTTCTCTGCCGTGGTCTGTGTTGTACCGTCCAGGAGCAA TGCTCATTCTGTGTTGAGCTTGCATCTGCATAGAGACATAT TCTT CACTGACAGCCAGC TCTTTGG CA ACCAGGCTGCGGTGGACAGCGCCATCGATGACATTTCTGTATGCTGAA AGTGCCCAGGAGGAGTCTGCACGTGG3'

1.2.4. Caracterización de los fundadores y establecimiento final de las líneas

Los ratones fundadores con la edición deseada se identificaron mediante PCR con primers flanqueando la región editada (los primers de genotipado están recogidos en la Tabla III). Los productos de PCR se secuenciaron directamente o se subclonaron en el plásmido pBlueScript (Stratagene) para posteriormente secuenciarlos mediante secuenciación Sanger. Los fundadores seleccionados se cruzaron con ratones silvestres (WT, *Wild-Type*) C57BL/6J para eliminar los off-targets no deseados. Posteriormente, los descendientes heterocigotos fueron secuenciados de nuevo y cruzados para generar los homocigotos editados. El genotipado se llevó a cabo mediante análisis en geles de agarosa y secuenciación Sanger, en los casos necesarios (mutantes puntuales), de los productos de PCR obtenidos de DNA genómico extraído de biopsias de cola (ver secciones 2.1.1 y 2.2).

Tabla III. Primers y tamaño de amplicón esperado en el genotipado por PCR de cada modelo.

Modelo	Primer	Secuencia	Tamaño (bp)
Usp26^{21/21}	F1	5'- TCCTTCAAACCTATGGCGAAGA -3'	Alelo WT: 411
	R1	5'- AAGGTTGATGAGGATGGAAGCA -3'	Alelo 21: 307
Usp26^{Cat/Cat}	F2	5'- TTCTCCAGTTCTGTTGAAACACAC -3'	Alelo WT: 259
	R2	5'- TTTCAAAACAAGCAGTTGCATCA -3'	Alelo Cat: 173
Hsf2bp^{-/-} Hsf2bp^{S167L/S167L}	F1	5'-TTCTTTGAAAGATGTGACAGTTC-3'	Alelos WT and S167L: 383
	R1	5'-ACCTGGGTTTCCTTTAGATCAGTTA-3'	Alelo KO: 355
Brme1^{Δ142-472/Δ142-472}	F2	5'-GAAAGTTCTTCAGAGTGCTTGCT-3'	Alelo WT: 1228
	R2	5'-AGCCCTATCTGTACCTAAAG-3'	Alelo Δ: 235
	F3	5'-CCCAGCAGATGCCTCTCTTAT-3'	Alelo WT: 150
	R3	5'-CTCAGCAGAGTTCCAATGCAG-3'	Alelo Δ: No amplifica
Brme1^{-/-}	F2	5'-GAAAGTTCTTCAGAGTGCTTGCT-3'	Alelo WT: 646
	R3	5'-CTCAGCAGAGTTCCAATGCAG-3'	Alelo KO: 606
Spo11^{-/-}	F4	5'- AGAGCCCCCAGTGCTCTTAAC-3'	Alelo WT: 416
	R4	5'- GGCAGACCCCTCTACCTCTGT-3'	Alelo KO: 417
Rnf212^{-/-}	F5	5'- TTTCTTTGCCTCCGTACTTTTGG-3'	Alelo WT: 557
	R5	5'- CCCAGGCTTTACTTCAACAACAA -3'	Alelo KO: 415
Hei10^{-/-}	F6	5'- CTGCCTGTTCTCACATCTTC-3'	Alelo WT: 100
	R6	5'- AGCTTTCAGAAAGGGTACTG -3'	Alelo KO: 96

1.3. Modelos murinos generados como herramienta en esta tesis

Los modelos nulos para *Rnf212*, *Hei10* y *Spo11* ya han sido generados y caracterizados en trabajos anteriores y no son objeto de estudio de la presente tesis. Sin embargo, teniendo en cuenta la experiencia de nuestro laboratorio en la generación de modelos mutantes mediante CRISPR/Cas9, decidimos generar nuestros propios modelos *Rnf212^{-/-}*, *Hei10^{-/-}* y *Spo11^{-/-}* por dos motivos. El primero es tratar de trabajar siempre con animales que tengan el mismo fondo genético y el segundo evitar los procedimientos de rederivación indispensables para la introducción de nuevos animales en el animalario del SEA de la Universidad de Salamanca. Los mutantes para *Rnf212*, *Hei10* y *Spo11* se describen por primera vez en este trabajo (Figura VIII) y han sido validados comparando sus fenotipos (arresto meiótico, defectos en sinapsis, alteraciones moleculares, etc.) con los de los mutantes ya publicados (Baudat et al. 2000; Qiao et al. 2014).

Figura VIII. Generación y caracterización de los mutantes *Spo11*^{-/-}, *Rnf212*^{-/-} y *Hei10*^{-/-}. (a-c) Representación esquemática de los locus murinos *Spo11*, *Rnf212* y *Hei10* y la estrategia de edición genómica seguida en cada caso. Cuadros grises: exones codificantes; cuadros blancos: no codificantes. Las líneas gruesas (secuencias codificantes) y finas (no codificantes) localizadas bajo los exones indican la transcripción esperada desde el alelo WT y los alelos nulos. ATG, codón de inicio, TGA/TAA y *, codones de stop. Flechas: primers de genotipado (F y R). (a) El cromatograma muestra la secuencia de la región editada en ratones WT y *Spo11*^{-/-}. Panel de la derecha: doble IF de SYCP3 y SYCP1 mostrando un bloqueo en zigotena-like con sinapsis entre no homólogos en los ratones *Spo11*^{-/-}, fenotipo previamente descrito para estos mutantes (Baudat et al. 2000). (b-c) Paneles de la derecha: doble IFs de SYCP3 con RNF212 o HEI10 en los mutantes (b) *Rnf212*^{-/-} y (c) *Hei10*^{-/-} mostrando la ausencia de la proteína mutada en cada caso (demostración de que son alelos nulos) y una localización aberrante de (b) HEI10 y (c) RNF212 como se ha descrito previamente para estos mutantes (Qiao et al. 2014).

1.4. Otros modelos

El modelo mutante para *Rec8* ha sido descrito previamente (Bannister et al. 2004). Los modelos *knockout* para *Six6os1* and *Psmas8* han sido desarrollados durante la presente tesis y han sido descritos en las siguientes publicaciones (Gomez-H et al. 2016; Gomez-H et al. 2019).

2. Técnicas de Biología Molecular

2.1. Extracción de DNA genómico

2.1.1. Extracción por lisis alcalina

Se tomaron fragmentos de colas de unos 2 mm de longitud de ratones de 21 días post-parto (dpp) y se incubaron en 600 μ l de NaOH 50 mM durante 30 minutos a 95°C. A continuación, los fragmentos se disgregaron mediante *vortex* y se neutralizó el NaOH con 50 μ l de Tris-HCl 1M pH 8,8. Este DNA genómico se utilizó como molde para el genotipado de ratones por PCR.

2.1.2. Extracción con fenol/cloroformo

Este método de extracción permite obtener una mayor pureza de DNA eliminando un alto porcentaje de proteínas y polisacáridos. Para llevarlo a cabo se incubaron los fragmentos de cola de ratón en 500 μ l de *buffer* de lisis (0,1 M Tris-HCl pH 7,4, 0,1 M EDTA, 0,5% SDS) suplementado con 0,6 mg/ml de Proteinasa K (Roche) a 55°C durante 12-20 horas. Al DNA solubilizado se le añadieron 200 μ l de fenol y 200 μ l de cloroformo:alcohol isoamílico (24:1), se mezcló por inversión y se centrifugó 5 min a 13000 rpm. La fase acuosa superior que contiene el DNA se recogió y el DNA se precipitó añadiendo 200 μ l de acetato amónico 7,5 M y 700 μ l de isopropanol (0,4 y 1,4 volúmenes, respectivamente). A continuación, el DNA se lavó con 500 μ l de etanol 70%. Una vez retirado el etanol, se dejó evaporar completamente y el DNA se resuspendió en TE 1x.

2.2. Genotipado por PCR

El genotipado de ratones se llevó a cabo mediante amplificación por PCR. Los *primers* se diseñaron flanqueando la región editada del gen de interés. La puesta a punto de las condiciones de reacción se hizo en base al protocolo de la casa comercial de la Taq DNA polimerasa (NZYTaq II DNA polymerase, NZYTech). 1,5 μ l de DNA (< 0,1 μ g), extraído mediante lisis alcalina de biopsias de cola, se utilizó como molde en una mezcla que contenía 0,4 μ M de cada *primer*, 200 μ M de dNTPs (desoxiribonucleótidos trifosfato), *Buffer* polimerasa 1x y 0,75 unidades de Taq polimerasa en un volumen final de 25 μ l. La reacción se llevó a cabo en un termociclador Veriti (Thermo Fisher). Las condiciones de PCR varían en función del tamaño del amplicón y de los *primers* utilizados (ver Tabla III). Sin embargo, existen ciertas condiciones comunes: desnaturalización inicial de 2 min a 94°C; 35 ciclos de i) desnaturalización: 20 seg a 94°C, ii) anillamiento: 20 seg a 55-68°C en función de los *primers* utilizados, iii) elongación: a 72°C y tiempo de 1 min por kb a amplificar; una elongación final de 5 min a 72°C. Los productos de PCR se analizaron por electroforesis en geles de agarosa y se secuenciaron en los casos necesarios (mutantes puntuales).

2.3. Generación de vectores de expresión

Con el fin de obtener proteínas fusionadas con diferentes epítomos (Flag, EGFP, HA o Myc) para los diferentes experimentos en los que se requieren, se clonaron los cDNAs completos o truncados de las proteínas de interés en diferentes vectores de expresión eucariota.

2.3.1. Extracción de RNA

Se colocaron 100 mg del tejido de partida en un tubo Eppendorf de 2 ml con 750 µl de GIT (tiocianato de guanidinio 4M, citrato de sodio 25mM, sodio lauril sarcosinato 0,5% (w/v) y β-mercaptoetanol 0,1M). El tejido fue disgregado con un homogeneizador Polytron (IKA T10 basic, UltraTurrax). A continuación, se añadieron 0,1 volúmenes de acetato de sodio 2 M pH 4,0, 1 volumen de fenol equilibrado con H₂O, 0,4 volúmenes de cloroformo:alcohol isoamílico (24:1) mezclando fuertemente con *vortex* tras cada solución añadida. Esta mezcla se incubó 15 min a temperatura ambiente (RT, *room temperature*) y se centrifugó a 13000 rpm durante 15 min a 4°C. La fase acuosa superior que contiene el RNA se transfirió a un nuevo tubo y el RNA se precipitó añadiendo 1 volumen de isopropanol y centrifugando de nuevo 15 min a 13000 rpm y 4°C. El pellet de RNA se lavó dos veces con etanol 70% y se dejó secar al aire. Finalmente, el RNA se resuspendió en H₂O MilliQ y se cuantificó midiendo la absorbancia a 260 nm con NanoDrop (Thermo Scientific) o Qubit (Thermo Scientific).

2.3.2. Transcripción reversa-PCR (RT-PCR)

El cDNA total de testículo fue producido mediante retrotranscripción a partir de 5 µg de RNA total usando oligonucleótido(dT) y siguiendo el protocolo del Kit SuperScript® II Reverse Transcriptase (Invitrogen). Este cDNA se utilizó como molde para amplificar por PCR los cDNAs de interés usando oligonucleótidos específicos y las polimerasas Phusion High-Fidelity (ThermoFisher) o Expand Long polymerase (Roche) para los amplicones de gran tamaño. Para ello se añadieron 2 µl de cDNA total en un mix formado por 0,4 µM de cada *primer*, 200 µM dNTPs, 1x *buffer* polimerasa y la polimerasa adecuada (0,8 U Phusion, 3,5 U Expand Long) en un volumen total de 50 µl. Las condiciones estándar de PCR fueron: una desnaturalización inicial de 2 min a 94°C seguido de 35 ciclos de: i) desnaturalización: 20 seg a 94°C, ii) anillamiento: 20 seg a 55-68°C en función de los *primers* utilizados, iii) elongación: 1 min por kb a amplificar a 72°C (Phusion polymerase) o 68°C (Expand Long polymerase); una elongación final de 5 min a 72/68°C. Los productos de PCR se analizaron por electroforesis en geles de agarosa.

2.3.3. Preparación de insertos y *clonning*

Los cDNAs amplificados como se ha explicado en el apartado anterior se purificaron en columna (NZYGelpure, MB011, NZYTech) y se fosforilaron usando la T4 polinucleótido quinasa (Takara). En los casos necesarios (PCRs realizadas con Expand Long, que no deja extremos romos), los insertos se repararon usando la T4 DNA polimerasa (Takara) siguiendo las instrucciones del fabricante. Por último, estos insertos se clonaron en diferentes vectores de expresión en mamíferos (pcDNA3, pcDNA3 2xFlag, pEGFP-C1, pEGFP-N1, pcDNA3 2xHA, pcDNA3.1 Myc-His (-)) mediante ligación usando la enzima T4 ligasa (Takara) durante 3 horas a RT. Esta ligación posteriormente se transformó en células competentes de *E.coli*.

2.3.4. Transformación de células competentes DH5α

La propagación del plásmido obtenido como producto de la ligación se realizó en células competentes de *E.coli* DH5α por choque térmico. Para ello, se descongeló un vial de células competentes en hielo y se añadieron 4 µl de la ligación. El vial se incubó durante 30 minutos en hielo, seguido de 33 segundos en un baño a 42°C y 1 minuto en hielo. Tras esto, las bacterias se dejaron recuperar durante 1 hora (resistencia a Ampicilina) o 3 h (resistencia a Kanamicina) en 1 ml de medio LB líquido sin antibiótico a 37°C en agitación para permitir la expresión del gen de resistencia al antibiótico. Por último, se sembraron en una placa de LB agar que contenía el antibiótico de selección correspondiente y se incubaron a 37°C durante al menos 15 horas. En el caso de transformación de plásmidos puros, no procedentes de ligación, se incubaron 30 µl de células competentes con 1 µl de plásmido durante 15-20 minutos en hielo, seguido de 33 segundos en un baño a 42°C y 1 minuto en hielo. Tras esto, cuando la resistencia era a Ampicilina, las bacterias se sembraron directamente en una placa de LB agar. Pero en los casos en que el plásmido lleva resistencia a Kanamicina, se recuperaron durante 1 hora en 1 ml de medio LB líquido a 37°C en agitación, antes de ser sembradas en placa.

2.3.5. Extracción de DNA plasmídico de un cultivo bacteriano por lisis alcalina

Las colonias de *E.coli* obtenidas en la placa de LB se inocularon en 1,5 ml de LB líquido con el antibiótico de selección y se incubaron a 37°C en agitación hasta el día siguiente. El cultivo se centrifugó 2 min a 13000 rpm y el pellet de bacterias obtenido se resuspendió en 200 µl de *buffer* P1 (50 mM de Tris HCl pH 8, 10 mM de EDTA y 100 µg/ml de RNasa). A continuación, se lisó añadiendo 200 µl de *buffer* P2 (0,2 M de NaOH 5M y 1% de SDS), mezclando por inversión e incubando 5 min a RT. Transcurridos los 5 minutos, se añadió 200 µl de *buffer* P3 (3 M de acetato de potasio y 11,5% de ácido acético glacial) para neutralizar la mezcla de lisis y precipitar el DNA bacteriano y las proteínas. Tras centrifugar 10 min a 13000 rpm, se recogió el sobrenadante, que contiene el DNA plasmídico, y se precipitó añadiendo 0,7 volúmenes de isopropanol, mezclando vigorosamente y centrifugando 5 min a 13000 rpm. El pellet de DNA se lavó con etanol al 70% y se resuspendió en TE 1x.

2.3.6. Análisis de restricción

Con el fin de comprobar qué colonias contenían el plásmido recombinante correcto, se llevó a cabo un análisis por digestión enzimática. Para ello, se digirió el DNA (1 µl) con 0,2 µl de cada una de las enzimas de restricción necesarias y el *buffer* correspondiente (1x) en 10 µl de volumen final. Esta mezcla se incubó durante 1 hora a la temperatura indicada para cada enzima. Finalmente se analizaron los fragmentos de restricción obtenidos por electroforesis en gel de agarosa y, en los casos correctos, se validó mediante secuenciación Sanger que el *cloning* estaba en pauta y no había errores.

2.3.7. Extracción de DNA plasmídico por lisis alcalina seguido de purificación por columna de afinidad

Una vez validado el *cloning* mediante secuenciación se obtuvo DNA plasmídico con alto grado de pureza mediante columna de intercambio aniónico. Para ello, previamente se sembró un cultivo de una colonia de *E. coli* transformada con el plásmido de interés, en 200 ml de LB suplementado con el antibiótico apropiado

según la resistencia del plásmido y se incubó a 37°C en agitación hasta obtener un cultivo saturado. Este se centrifugó a 7000 rpm 15 minutos y el pellet obtenido se resuspendió en 10 ml de *buffer* P1 (50 mM de Tris HCl pH 8, 10 mM de EDTA y 100 µg/ml de RNAsa). Las células se lisaron con 10 ml de *buffer* P2 (NaOH 5M y 1% de SDS), incubándolo 5 minutos a temperatura ambiente y tras ello, se neutralizó con 10 ml de *buffer* P3 (3 M de acetato de potasio y 11,5% de ácido acético glacial) y se centrifugó a 14000 rpm durante 10 minutos, quedando el DNA plasmídico en el sobrenadante. Por otro lado, se equilibró una columna (JETSTAR Column) con 30 ml de *buffer* E4 (NaCl 600 mM, ácido sódico 100 mM, Tritón X-100 0,15% y ácido acético hasta pH 5). Una vez equilibrada, el sobrenadante del lisado celular, previamente filtrado en una jeringa con lana de vidrio, se pasó por la columna. Tras ello, se lavó la columna con 60 ml de *buffer* E5 (NaCl 800 mM y ácido sódico 100 mM, ajustando con ácido acético hasta pH 5). El DNA se eluyó con 15 ml de *buffer* E6 (NaCl 1,5 M, ácido sódico 100 mM y ácido acético hasta pH 5), y se precipitó con 0,7 volúmenes de isopropanol. Se centrifugó a 15500 rpm durante 30 minutos a 4°C y el pellet se lavó con etanol al 70%, centrifugándolo 10 minutos a 15500 rpm. El pellet se dejó secar para que se evaporara el etanol y finalmente se resuspendió en 200 µl de 1x TE (10 mM de Tris pH 8 y 1 mM de EDTA). Para determinar la concentración de DNA, se midió su absorbancia en el espectrofotómetro NanoDrop.

2.4. Análisis de proteínas

2.4.1. Extracción de proteínas de testículo de ratón

Este tipo de extracción nos permite obtener una muestra de todas las proteínas expresadas en testículo para así poder analizar aquellas que nos interesen y determinar si, en los diferentes mutantes generados, estas se encuentran aumentadas, disminuidas o totalmente ausentes. Para ello, los testículos se extrajeron de la cavidad abdominal de los ratones y se retiró la túnica albugínea. Cada medio testículo se homogeneizó en 600 µl de *buffer* de lisis RIPA (50 mM Tris-HCl pH 7,5, 150 mM NaCl, 1% NP40, 0,5% desoxicolato de sodio, 0,1% SDS) suplementado con inhibidores de proteasas (Complete EDTA-free, Roche) con un homogeneizador Polytron (IKA T10 basic, UltraTurrax). Las muestras se incubaron 15 min en hielo para permitir la extracción proteica y, a continuación, se centrifugaron a 13000 rpm durante 30 min a 4°C. Finalmente, se cuantificó la concentración proteica del sobrenadante mediante un ensayo Bradford con el Kit DC Protein Assay (BioRad). Las proteínas así extraídas se utilizaron directamente para análisis por *Western Blot* (WB).

2.4.2. Inmunoprecipitación de proteínas endógenas de testículo

La inmunoprecipitación de proteínas endógenas permite identificar *in vivo* las proteínas que interaccionan con una proteína de interés. Para ello, se inmunoprecipita dicha proteína mediante el uso de anticuerpos (Abs, *antibodies*) específicos contra ella en unas condiciones que permiten mantener las interacciones endógenas. De esta manera al mismo tiempo que precipitamos nuestra proteína arrastraremos todas aquellas que interaccionen, directa o indirectamente, con ella y podremos identificarlas. La extracción de proteína para las inmunoprecipitaciones se llevó a cabo como se ha explicado en el apartado anterior salvo que en este caso se hizo en 1,5 ml de *buffer* de lisis Co-IP (50 mM Tris-HCl pH 8, 500 mM NaCl, 1 mM EDTA, 1% Triton X-100) suplementado con inhibidores de proteasas (Complete EDTA-free, Roche). 20 mg del extracto proteico se incubaron a 4°C en rotación toda la noche (ON, *over-night*) con

100 µl de bolas de Sefarosa acopladas a proteína G (GammaBind™ G Sepharose™, GE Healthcare) previamente unidas mediante *crosslink* (ver sección 2.4.3) al Ab contra la proteína de interés. De esta manera permitimos que la proteína de interés se una a las bolas a través del Ab específico y además retendremos también aquellas proteínas que interaccionen con ella. Tras la incubación, esta mezcla se transfirió a una columna (Plastic Small Column, CS-20, ABT) en la que quedan retenidas las bolas de Sefarosa así como las proteínas unidas a ellas. La columna se lavó 3 veces con 1 ml de *buffer* Co-IP y, a continuación, 3 veces con 1 ml de H₂O MilliQ para eliminar las sales y detergentes. Las proteínas retenidas se eluyeron con 100 µl de glicina 0,1 M pH 2,5-3, incubando 10 minutos a RT. Tras la primera elución se llevó a cabo una segunda de idéntica manera. Finalmente, las fracciones eluidas se neutralizaron con 5 µl de Tris-HCl pH 9,5 y ambas se juntaron para el análisis posterior por espectrometría de masas (MS/MS, ver sección 2.4.6). Al finalizar el procedimiento, las bolas unidas al Ab pueden ser reequilibradas y reutilizadas (ver la sección 2.4.3).

2.4.3. *Crosslink* de los anticuerpos a bolas de Sefarosa

100 µl de bolas de Sefarosa acopladas a proteína G (GammaBind™ G Sepharose™, GE Healthcare) se lavaron 2 veces con 500 µl de PBS 1x (Na⁺, no K⁺) recogiénolas por centrifugación (1 min a 10000 g). A continuación, se lavaron una vez más con 100 µl de *Dilution buffer* (PBS 1x Na⁺ con 1mg/ml de albúmina de suero fetal bovino (BSA, *Bovine Serum Albumin*)) rotando 10 min a 4°C. A continuación, se añadió el Ab (200 µg) sobre las bolas diluido en *Dilution buffer* (ratio 1:1, bolas:*Dilution buffer*) y se incubó toda la noche (mínimo 4 h) rotando a 4°C. A continuación, las bolas se recogieron por centrifugación y se lavaron con 10 volúmenes de *Dilution buffer* 5 min en rotación a 4°C, se centrifugaron de nuevo y se lavaron con 1 ml de PBS 1x (Na⁺). Para el *crosslink* del Ab las bolas se incubaron con 10 volúmenes (1 ml) de DMP 20 mM (*dimethyl pimelimidate*, Sigma-Aldrich) 30 min a temperatura ambiente rotando. El DMP se preparó justo en el momento de su utilización diluido en una solución formada por *Wash buffer* (0,2 M trietanolamina en PBS) y PBS 1x en proporción 1:1. Esta incubación seguida de un lavado con 500 µl de *Wash buffer* (5 min a RT en rotación) se repitió dos veces. Después de esto, para parar la reacción las bolas se incubaron dos veces con 500 µl de *Quenching buffer* (etanolamina 50 mM en PBS) durante 5 min a RT en rotación y finalmente se lavaron con 500 µl de PBS 1x. Para eliminar el exceso de Ab no unido a las bolas estas se lavaron 2 veces con glicina 1 M pH3 (10 min en rotación a RT). Por último, las bolas se lavaron 3 veces con 500 µl de PBS 1x (5 min en rotación a RT) y se guardaron hasta su uso en etanol 20% en PBS para evitar el crecimiento bacteriano.

2.4.4. Co-inmunoprecipitación de proteínas sobreexpresadas

Esta aproximación llevada a cabo en un sistema heterólogo permite validar *in vitro* interacciones directas entre dos o más proteínas candidatas. Estas proteínas, normalmente unidas a un epítipo, se obtuvieron mediante dos métodos distintos: (I) la expresión en células HEK293T transfectadas donde estarán presentes además de las proteínas sobreexpresadas todas aquellas endógenas de las células HEK293T o (II) un sistema de transcripción/traducción *in vitro* (TnT), donde solo estarán presentes las proteínas de interés y las del sistema TnT. En el primer caso, se transfectaron células HEK293T transitoriamente (ver sección 4.2) con al menos dos plásmidos de expresión que codificaran para las proteínas candidatas fusionadas con distintos epítopos (Flag, GFP, HA o Myc). 48 horas tras la transfección se extrajo la proteína total de las células en *buffer* de lisis D (50 mM Tris-HCl pH 7,4, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100) suplementado con inhibidores de proteasas (Complete EDTA-free, Roche). Para ello, las células se recogieron en 1 ml de

buffer de lisis D y se homogeneizaron con un homogeneizador Polytron (IKA T10 basic, UltraTurrax). Las muestras se incubaron 10 min en hielo para permitir la extracción proteica y, a continuación, se centrifugaron a 13000 rpm durante 15 min a 4°C. La concentración proteica del sobrenadante se cuantificó mediante ensayo Bradford con el Kit D_c Protein Assay (BioRad). Alternativamente, para obtener extractos más puros de las proteínas de interés, estas se produjeron mediante transcripción-traducción *in vitro* usando el Kit TnT® T7 *Coupled Reticulocyte Lysate Systems* (Promega, ver sección 2.6). 1 mg de proteína total (obtenida de extractos de HEK293T) o 15 µl de los extractos de TnT se bloquearon incubando con bolas de Sefarosa unidas a proteína G (GammaBind™ G Sepharose™, GE Healthcare) durante 1 hora a 4°C rotando. A continuación, el extracto bloqueado se incubó con el Ab correspondiente durante al menos 2 horas a 4°C rotando (ver Abs en Tabla V). Como control negativo los extractos proteicos se incubaron con IgGs (Inmunoglobulinas G; 5 µg/1 mg de proteína) producidas en la misma especie que el Ab específico utilizado. Los inmunocomplejos (formados por el Ab y las proteínas diana) se capturaron incubando en rotación con bolas de Sefarosa-proteína G toda la noche a 4°C. Al día siguiente, tras 5 lavados con *buffer* de lisis (centrifugando 1 min a 10000g y 4°C), las proteínas se eluyeron resuspendiendo las bolas en 20 µl de 2x *buffer* Laemmli (100 mM Tris-HCl pH 7, 4% SDS, 0,2% azul de bromofenol, 200 mM β-mercaptoetanol y 20% glicerol) e hirviéndolas 5 min a 100°C. La mitad de cada muestra se cargó en geles de acrilamida y las proteínas se detectaron mediante WB con los Abs indicados. Los Abs utilizados para las inmunoprecipitaciones y para el WB se describen en las Tablas IV y V.

En el caso de trabajar con proteínas que interaccionan débilmente, se añadió al medio de cultivo 1 mg/ml de ditiobis (*succinimidyl propionate*, Sigma-Aldrich) 10 min antes de lisar las células para estabilizar las interacciones lábiles. En este caso la extracción de proteína se llevó a cabo en *buffer* de lisis P (20 mM Tris-HCl pH 7,5, 100 mM NaCl, 20 mM β-glicerofosfatos, 5 mM MgCl₂, 1 mM NaF, 0,2% ND P-40 y 0,5 mM DTT) de igual manera que se ha explicado anteriormente.

2.4.5. Western Blot

Los extractos proteicos se resolvieron en geles desnaturalizantes siguiendo el protocolo descrito por Laemmli (Laemmli 1970). Así, las proteínas se separaron en geles de poliacrilamida (7-13%) que se corrieron en *buffer* SDS-Page 1x (250 mM Tris-HCl, 200 mM glicina, 0,05% SDS) a 120 V. Las proteínas se transfirieron a membranas de nitrocelulosa de 0,45 µm (GE Healthcare Amersham) en *buffer* de transferencia (25 mM Tris-HCl, 200 mM glicina, 20% metanol) a 65 V durante 1 hora. Una vez transferidas las proteínas se bloquearon las membranas con leche desnatada al 5% en TBST (TBS-Tween 20: 50 mM Tris-HCl pH 7,5, 150 mM NaCl, 0,1% Tween 20) durante 1 hora en agitación a RT. A continuación, las membranas se incubaron rotando con el Ab primario diluido en leche al 2,5% en TBST durante 1 hora a RT (ver Abs y diluciones en Tabla V). Tras la incubación las membranas se lavaron 3 veces con TBST (5 min en agitación) y se incubaron rotando con el Ab secundario conjugado con peroxidasa o con un fluorocromo incubándose en este caso en oscuridad (ver Abs secundarios en Tabla IV). Finalmente, para el revelado de la señal quimioluminiscente de la peroxidasa se incubaron las membranas con el sustrato Immobilon Western HRP (Millipore) durante 5 min y se expusieron a películas fotográficas el tiempo apropiado. En el caso de utilizar Abs secundarios conjugados con fluorocromos el revelado se llevó a cabo en con el sistema Odyssey Infrared Imaging.

Para la reincubación de membranas con nuevos Abs, se eliminaron el Ab primario y secundario incubando la membrana en *buffer de stripping* medio (0,2M glicine, 1% SDS, 0,01% Tween 20, pH 2,2) dos veces de 10 min en agitación a RT. A continuación, la membrana se lavó dos veces de 10 min con PBS, dos veces de 5 min en TBST y finalmente, se bloqueó e incubó de forma normal. Por otro lado, la membrana puede ser incubada sin eliminar los Abs (sin *stripping*), simplemente inactivando la actividad de la peroxidasa conjugada con el Ab secundario con un exceso de peróxido de hidrógeno (H₂O₂). Para ello, la membrana se incubó en agitación con H₂O₂ al 15% en PBS durante 15 min. A continuación, se lavó dos veces con TBST (5 min) y se bloqueó y reincubó de forma normal.

Tabla IV. Abs secundarios utilizados para WB. HRP: *Horseradish peroxidase*

Anticuerpo	Referencia	Especie (Host)	Dilución	Casa comercial
α-mouse HRP	NA931	Sheep	1:10000	GE Healthcare
	715-035-150	Donkey	1:5000	Jackson Immunoresearch
α-rabbit HRP	7074	Goat	1:3000	Cell Signaling
	711-035-152	Donkey	1:5000	Jackson Immunoresearch
α-goat HRP	A27014	Rabbit	1:10000	Thermo Scientific
	705-035-147	Donkey	1:5000	Jackson Immunoresearch
α-mouse DyLight™ 680	35518	Goat	1:10000	Thermo Scientific
α-rabbit DyLight™ 800	35571	Goat	1:10000	Thermo Scientific

2.4.6. Análisis por espectrometría de masas (MS/MS)

Esta técnica permite identificar todas las proteínas presentes en un extracto proteico. De esta manera, tras la inmunoprecipitación de una proteína de interés, en la que habremos arrastrado las proteínas que interaccionan con ella, podremos identificar los posibles interactores. Para ello, las proteínas inmunoprecipitadas como se ha descrito anteriormente (sección 2.4.2) se resolvieron en geles de poliacrilamida-SDS que se tiñeron con azul de Coomassie. Las bandas resultantes en estos geles se cortaron y se digirieron con tripsina. Los péptidos obtenidos se analizaron con un espectrómetro de masas LTQ-Orbitrap Velos. Los datos crudos (*Raw MS data*) se analizaron usando MaxQuant v1.6.2.6 (Cox and Mann 2008) con búsquedas frente al proteoma de *Mus musculus* (SwissProt UP000000589) y los contaminantes presentes en MaxQuant. Todos los FDRs (*False Discovery Rate*) fueron del 1%. Las modificaciones variables tenidas en cuenta fueron la oxidación de metioninas y la acetilación del N-terminal, mientras que en las modificaciones fijas solo consideró la carbamidometilación de cisteínas. El número máximo de modificaciones permitidas por péptido fue de 5. La intensidad de proteína se calculó usando el iBAQ (Schwanhüsser et al. 2011). Se eliminaron los posibles contaminantes, secuencias señuelo (*reverse decoy sequences*) y las proteínas identificadas por presentar un péptido con residuos modificados (*proteins identified by site*). Las proteínas con menos de dos péptidos únicos en ambos Abs utilizados (Ab1 y Ab2) no se consideraron para el análisis posterior. Las proteínas con menos de dos péptidos únicos en el control

(inmunoprecipitación con IgGs de la misma especie que los Abs utilizados) y más de dos con ambos Abs se seleccionaron como candidatos con alta credibilidad (grupo “Ab1 y Ab2”). Además, se seleccionó un grupo adicional de candidatos correspondiente a aquellas proteínas con dos o más péptidos únicos con uno de los dos Abs (Ab1 o Ab2) y que no mostraban péptidos únicos en el control (grupos “Ab1 solo” y “Ab2 solo”, respectivamente). El procedimiento experimental fue llevado a cabo por la Unidad de Proteómica del Centro de Investigación del Cáncer (CSIC/USAL). El filtrado inicial de los datos fue realizado por Rodrigo García-Valiente (Academisch Medisch Centrum Universiteit van Amsterdam).

2.5. Ensayo de doble híbrido y screening

El ensayo de doble híbrido (*Yeast Two Hybrid*, Y2H) permite detectar interacciones entre proteínas y así identificar posibles candidatos a interactuar con una proteína de interés. En nuestro caso, este ensayo se llevó a cabo utilizando el Kit Matchmaker Gold Yeast Two-Hybrid System (Clontech) siguiendo las instrucciones del fabricante. El cDNA murino de *Hsf2bp* se amplificó de cDNA de testículo y se clonó en el vector pGBKT7 como se ha explicado en la sección 2.3. HSF2BP se utilizó como cebo (*bait*) contra una librería de cDNA de testículo de ratón (*Mate & Plate cDNA library*, Clontech Laboratories Inc.) que representan las presas (*prey*). Los clones positivos, en los que se produce interacción, se identificaron en un primer momento en un medio de doble restricción (SD (*synthetic dropout*)/–Leu/–Trp/X- α -Gal/Aureobasidina A) y luego se sometieron a una selección más restrictiva en un medio de restricción cuádruple (SD/–Ade/–His/–Leu/–Trp/X- α -Gal/Aureobasidina A). De los clones positivos en ambos medios se aislaron se extrajeron los plásmidos presa y se transformaron en *Escherichia coli*. Para identificar definitivamente los candidatos, los plásmidos de dos colonias bacterianas independientes se aislaron, se secuenciaron mediante secuenciación Sanger y se analizaron utilizando la herramienta nucleotide BLAST.

2.6. Transcripción-traducción *in vitro*

Este sistema nos permite producir *in vitro* y en un solo paso extractos de proteínas (en nuestro caso marcadas con un epítipo) más puros que los obtenidos mediante transfección y extracción en células HEK293T. Para llevarlo a cabo, se utilizó el Kit *TnT® T7 Coupled Reticulocyte Lysate Systems* (Promega). Como molde se utilizaron plásmidos que codifican para las proteínas de interés bajo el promotor T7 y unidas a los epítopos deseados (pcDNA3 2xFlag o pcDNA3 2xHA). La reacción se llevó a cabo en un volumen final de 50 μ l con 25 μ l de lisado de reticulocitos de conejo, 2 μ l de *buffer* TnT, 1 μ l de RNA polimerasa, 0,5 μ l de aminoácidos menos Leu (1mM), 0,5 μ l de aminoácidos menos Met (1mM), 1 μ l de inhibidor de RNAsas 40 U/ μ l (RNasin®, Promega) y 1 μ g de DNA (obtenido por lisis alcalina convencional). La reacción se incubó durante 90 min a 30°C y se analizó por WB antes de utilizarlo para las posteriores aplicaciones.

2.7. Ensayo de unión al DNA

Este tipo de ensayo nos permite determinar si una proteína tiene capacidad de unión al DNA, ya sea de hebra simple (ssDNA) o de doble hebra (dsDNA) precipitando el DNA y determinando si arrastra a la proteína de interés. Estos experimentos se llevaron a cabo usando el protocolo previamente descrito por (Souquet et al. 2013) con ligeras modificaciones. Como ssDNA se utilizó un oligonucleótido biotinilado de 60 bp

(purificación HPLC, IDT) con la siguiente secuencia: F:5'-GATCTGCACGACGCACACCGGACGTA TCTGCTATCGCTCATGTCAACCGCTCAAGCTGC /3'BiotinTEG/. Para generar el dsDNA se utilizó el oligonucleótido reverso complementario: ss60-mer R (No biotinilado): 5'- GCAGCTTGAGC GGTGACATGAGCGATAGCAGATACGTCCGGTGTGCGTCGTGCAGATC-3'. El anillamiento para generar el dsDNA se llevó a cabo en *annealing buffer* (50 mM NaCl, 25 mM Tris-HCl, pH 7,5) con cantidades iguales de ambos oligonucleótidos mediante un paso de desnaturalización (5 min a 95°C) seguido de una rampa de descenso lento de temperatura hasta temperatura ambiente. A continuación, el DNA se inmovilizó en bolas unidas a estreptavidina (Dynabeads M-280 Streptavidin, Dynal) siguiendo las instrucciones del fabricante. La relación fue 0,2 pmol de DNA por 1 µg de bolas.

Por otro lado, los extractos proteicos que se incubarán con el DNA se obtuvieron mediante transcripción-traducción *in vitro* (TnT). 15 µl de la proteína (unida a Flag) obtenida en el ensayo se preincubaron en hielo durante 10 minutos en DBB modificado (DBB: 50 mM Tris HCl, 100mM NaCl, 10% (w/v) glicerol, *Complete Protease inhibitor*, 1mM 2-mercaptoetanol pH 7,4 modificado con 25 mM Tris-HCl, 1 mM EDTA y 5 mg/ml de BSA). Tras la preincubación se añadieron 500 µg de bolas con el DNA inmovilizado (ss- o dsDNA) y se incubaron 1 hora a 4°C con agitación suave. Tras esto las bolas se lavaron 3 veces (5 min rotando a RT) en 700 µl DBB modificado sin BSA, seguido de un lavado con *rinsing buffer* (DBB modificado suplementado con 150 mM NaCl). Finalmente, las proteínas unidas al DNA se eluyeron resuspendiendo las bolas en 30 µl de Laemmli *buffer* e hirviéndolas 5 min a 95°C y se analizaron por WB.

3. Técnicas citológicas

3.1. Análisis histopatológico

Para llevar a cabo el análisis histológico de tejidos de ratón tras sacrificar al animal se tomaron los órganos de interés y se fijaron durante 24 h a RT en formol al 10%. En el caso de los testículos el fijador utilizado fue Bouin. A continuación, los tejidos se incluyeron en bloques de parafina y se cortaron en secciones del grosor deseado según la aplicación, 5 µm para la mayoría de ellas. Las secciones se tiñeron con hematoxilina-eosina o con PAS (*Periodic Acid-Schiff*)-hematoxilina siguiendo los protocolos estándar. Para el marcaje con TUNEL (*TdT mediated dUTP Nick End Labelling*), las secciones se desparafinaron y las células apoptóticas se detectaron con el *In situ Cell Death Detection Kit* (Roche). Los núcleos se tiñeron con DAPI. La inclusión, el corte y la tinción de muestras fueron realizadas por el Servicio de Patología Molecular Comparada del Centro de Investigación del Cáncer (CSIC/USAL). Las muestras fueron analizadas usando un microscopio OLYMPUS BX51 y las imágenes fueron tomadas con una cámara OLYMPUS DP70.

3.2. Esparcido de espermatoцитos (*spreading*)

Para el estudio de espermatoцитos de ratón las células se fijaron sobre portaobjetos usando la técnica clásica *dry-down* con pequeñas modificaciones del protocolo descrito en (Peters et al. 1997). Esta técnica permite obtener preparaciones con células estalladas, en las que se pierde citoplasma, pero se visualizan muy bien los núcleos en dos dimensiones. Para obtener este tipo de preparaciones, en primer lugar, se eliminó la túnica albugínea de los testículos y se colocaron en una placa Petri con una gota de PBS 1x. Los

túbulos seminíferos se fueron desmenuzando con la ayuda de dos bisturís, recogiendo las células extraídas con PBS en un tubo de 15 ml (Falcon). Una vez extraídas todas las células se añadió PBS 1x a la suspensión hasta un volumen de 6 ml y se centrifugó 9 min a 1200 rpm. El pellet se lavó de nuevo con 6 ml de PBS 1x centrifugando 9 min a 1200 rpm. Tras esto, las células se resuspendieron en sacarosa 100 mM pH 8,4 (80-400 μ l) y se incubaron durante 5 min para someterlas a un choque hipotónico. A continuación, se extendieron 400 μ l de fijador F10b (1% Paraformaldehído (PFA), 0,15% Triton X-100, 0,05% PBS, 2,5 mM tetraborato de sodio, pH 9,2) sobre portas limpios, se añadieron 20 μ l de la suspensión celular en la esquina superior del porta y se dispersó suavemente mediante movimientos en zigzag. Estas preparaciones se incubaron durante 2 horas en una cámara cerrada para permitir la fijación de las células y, a continuación, se dejaron secar al aire. Para eliminar el fijador, las preparaciones se lavaron con Photo-Flo 0,08% (Kodak) y se secaron al aire. La calidad de los esparcidos se analizó con un microscopio invertido de contraste de fases Nikon Eclipse TS100 y estos se conservaron a 4°C en PBS 1x suplementado con Azida al 0,05% para prevenir el crecimiento bacteriano.

3.3. Aplastado de túbulos seminíferos (*squashing*)

Este método desarrollado por Parra et al. (2002) nos permite obtener una monocapa de células extraídas de los túbulos seminíferos pero que mantienen su conformación 3D. Para obtener estas preparaciones, se eliminó la túnica albugínea de los testículos y los túbulos se fijaron durante 10 min en un fijador con 2% formaldehído y 0,1% Triton X-100 en PBS 1x. Se tomó una pequeña fracción de los túbulos y pedazos de ellos se repartieron sobre un porta previamente tratado con poly-L-lisina (1mg/mgl, Sigma-Aldrich) y con una gota de fijador. Sobre ellos se colocó un cubreobjetos sobre el que se pasó un lápiz presionando y, posteriormente, se presionó con la mano para obtener una monocapa de espermatozoides. Estas preparaciones se introdujeron en nitrógeno líquido durante unos segundos e inmediatamente se eliminó el cubre con ayuda de un bisturí y se introdujeron en PBS 1x. Finalmente, las preparaciones se conservaron a 4°C en PBS 1x suplementado con 0,05% de azida.

3.4. Esparcido de oocitos (*spreading*)

Los oocitos en profase se obtuvieron de embriones hembra de entre 14,5 días post-coito (dpc, correspondiente a leptotena) y 19,5 dpc (justo antes del arresto en dictiata), dependiendo de la etapa que se quisiera analizar. Para ello las hembras preñadas se sacrificaron, se extrajeron los embriones del útero y se pusieron en PBS 1x. Se extrajeron los ovarios de los embriones y se colocaron en un pocillo (placa de 24 pocillos) con 200 μ l de medio M2 (Sigma-Aldrich). Sobre este pocillo se añadieron 10 μ l de colagenasa A/M2 (50 mg/mgl) y se incubaron 20-30 min a 37°C para favorecer la disgregación de los ovarios. A continuación, los ovarios se transfirieron a 200 μ l de *buffer* hipotónico (30 mM Tris-HCl pH 8,2, 50mM sacarosa, 17 mM citrato de sodio, 5 mM EDTA, 0,5 mM DTT) y se incubaron 45 min a RT. Durante este tiempo se prepararon los portaobjetos (6 por par de ovarios) dibujando un círculo de adhesivo Rubbercement en el centro para delimitar una pequeña área. Tras el tratamiento hipotónico, los ovarios se transfirieron a 60 μ l de sacarosa 100 mM pH 8,2 y se pipetearon vigorosamente (con una p100) hasta obtener una suspensión unicelular, lo cual se validó con un microscopio. Después de esto, se pusieron 40 μ l de fijador (1% (w/v) PFA, 5 mM borato de sodio, 0,15% Triton X-100, 3 mM DTT, pH 9,2) en el interior del círculo de adhesivo, se añadieron 10 μ l de

suspensión celular en el centro de la gota de fijador y preparación se fue moviendo en zigzag para esparcir la suspensión celular. Las preparaciones se incubaron en una cámara cerrada durante 2 horas y, posteriormente, se dejaron secar al aire. Para eliminar el fijador, las preparaciones se lavaron con Photo-Flo 0,08% (Kodak) y se secaron al aire. La calidad de los esparcidos se analizó con un microscopio invertido de contraste de fases Nikon Eclipse TS100 y estos se conservaron a 4°C en PBS 1x suplementado con Azida al 0,05% para prevenir el crecimiento bacteriano.

3.5. Inmunofluorescencia

Las preparaciones (esparcidos y aplastados) se incubaron con el Ab primario (Tabla V) diluido en PBS 1x durante 1 hora u ON a temperatura ambiente en una cámara húmeda. Tras esto, se lavaron 3 veces en PBS 1x durante 5 min y se incubaron 1 hora con el Ab secundario correspondiente conjugado con un fluorocromo diluido en PBS 1x (Tabla VI). Finalmente, las preparaciones se lavaron 3 veces de 5 min en PBS 1x y se montaron con medio de montaje Vectashield® (Vector Laboratories) y DAPI (4',6-Diamidino-2-Phenylindole, 10 µg/ml) para teñir el DNA. Con el fin de reducir la señal de fondo (background), especialmente en las inmunofluorescencias (IFs) de proteínas que forman foci, las preparaciones se bloquearon 15 min con *buffer* ADB 10% (10% Suero Fetal Bovino (FBS), 3% BSA y 0,05% Triton X-100 en PBS 1x) y los Abs se diluyeron en este mismo *buffer* para las incubaciones. En este caso los lavados se hicieron con PBST (PBS 1x con 0,05% Triton X-100).

Para las IFs de células en cultivo, las células se sembraron en placas de 3,5 cm sobre un cubre tratado con gelatina (0,5%, *Gelatin from cold water fish skin*, G7041, Sigma-Aldrich) para favorecer la adhesión celular. Para fijar las células se retiró el medio y se añadió en la misma placa 1 ml de PFA al 4% en PBS 1x. Se incubaron durante 7 min a 4°C y, posteriormente, se lavaron 3 veces con 2 ml PBS 1x. Una vez fijadas, los cubres se pegaron sobre portaobjetos y las células se permeabilizaron durante 4 min con *buffer* KB (0.1 M NaCl, 20 mM Tris-HCl pH 7,5, 0,1% BSA) con 0,2% Triton X-100 seguido de un lavado en PBS 1x. El bloqueo de las células se llevó a cabo con FBS al 7% en PBS 1x durante 30 min a RT (en cámara húmeda). Después, las células se incubaron 1 hora a RT con el Ab primario diluido en FBS al 7% en PBS 1x. Tras 3 lavados de 5 min en PBS, se incubó con el Ab secundario conjugado con un fluorocromo diluido en FBS al 7% en PBS 1x durante 1 hora. Finalmente, se lavaron 3 veces con PBS y se montaron con medio de montaje Vectashield® (Vector Laboratories) y DAPI (4',6-Diamidino-2-Phenylindole, 10 µg/ml) para teñir el DNA.

Material y métodos

Tabla V. Abs primarios utilizados para IF, WB e inmunoprecipitaciones (IP, µg de Ab por mg de proteína). * indica la dilución para IF de células en cultivo. El resto de las diluciones son para IF en esparcidos o aplastados.

Proteína detectada	Referencia	Especie (Host)	Tipo	Dilución			Casa comercial
				IF	WB	IP	
ACA	15-235	Human	IgG	1:15			Antibodies Inc.
BRME1	R1 (por encargo)	Rabbit	IgG	1:100	1:2000		Proteogenix
	R2 ((por encargo)	Rabbit	IgG	1:100	1:2000		Proteogenix
CDK2	sc-6248	Rabbit	IgG	1:20			Santa Cruz
DMC1	sc-22768	Rabbit	IgG	1:20			Santa Cruz
	R1 (por encargo)	Rabbit	IgG	1:500			Proteogenix
	R2 (por encargo)	Rabbit	IgG	1:500			Proteogenix
	ab11054	Mouse	IgG	1:50			Abcam
Flag	F1804	Mouse	IgG	1:50*	1:3000	5 µg	Sigma-Aldrich
	F7425	Rabbit	IgG	1:50*	1:2000	5 µg	Sigma-Aldrich
GFP	CSB-MA000051M0m	Mouse	IgG		1:3000	5 µg	Cusabio
	A-11122	Rabbit	IgG	1:50	1:3000	5 µg	Life Technologies
	sc-5385	Goat	IgG		1:3000		Santa Cruz
HA	αHA.11 101R	Mouse	IgG		1:2000	5 µg	Covance
	H6908	Rabbit	IgG		1:1000	5 µg	Sigma-Aldrich
HSF2BP	R1 (por encargo)	Rabbit	IgG	1:30	1:2000		Proteogenix
	R2 (por encargo)	Rabbit	IgG	1:30	1:2000		Proteogenix
HEI10	TA502700	Mouse	IgG	1:60			Origene
MLH1	51-1327GR	Mouse	IgG	1:20			BD Biosciences
RAD51	sc-8349	Rabbit	IgG	1:50			Santa-Cruz
	PC130	Rabbit	IgG	1:50			Calbiochem
RNF212		Goat	IgG	1:150			Dr. N. Hunter
RPA1	Molly-RPA	Rabbit	IgG	1:30			Dr. E. Marcon
RPA2	2208S	Rat	IgG	1:100	1:1000		Cell Signalling
SPATA22	16989-1-AP	Rabbit	IgG	1:60			Proteintech
SYCP1	Joe	Rabbit	Serum	1:200			
	K919	Rabbit	Serum	1:60			Dr. J.L. Barbero
	VAL G14/ VAL G13	Chicken	IgG	1:10			Inmunostep
SYCP3	sc-74569	Mouse	IgG	1:1000			Santa Cruz
	K921/K1037	Rabbit	Serum	1:500			Dr. J.L. Barbero
αUSP26 W-12	sc-51013	Goat	IgG	1:10	1:1000		Santa-Cruz
αUSP26-M		Rabbit	IgG	1:10	1:1000		Dr.Yi-Wen Lin
αTubulin	T9026	Mouse	IgG	1:100	1:10000		Sigma
β-Actin	Clone AC-15; A5441	Mouse	IgG		1:10000		Sigma-Aldrich
γH2AX (Ser139)	#05-636	Mouse	IgG	1:200			Millipore
	#07-164	Rabbit	IgG	1:400			Millipore
ChromePure mouse IgG	015-000-003	Mouse	IgG			5 µg	Jackson Immunoresearch
ChromePure rabbit IgG	011-000-003	Rabbit	IgG			5 µg	Jackson Immunoresearch
ChromePure goat IgG	011-000-003	Goat	IgG			5 µg	Jackson Immunoresearch

Tabla VI. Abs secundarios conjugados con fluorocromos utilizados para inmunofluorescencia. FITC: Fluoresceína, TRITC: Rodamina, AMCA: Aminomethylcoumarin acetate .

Anticuerpo	Fluorocromo	Referencia	Especie (Host)	Dilución	Casa comercial
α-chicken	TRITC	703-025-155	Donkey	1:100	Jackson Immunoresearch
	FITC	703-095-155	Donkey	1:100	Jackson Immunoresearch
α-goat	Rhodamine Red	705-295-147	Donkey	1:100	Jackson Immunoresearch
	FITC	705-095-147	Donkey	1:100	Jackson Immunoresearch
α-guinea pig	TRITC	706-025-148	Donkey	1:100	Jackson Immunoresearch
	FITC	706-095-148	Donkey	1:100	Jackson Immunoresearch
α-human	Texas Red	709-075-149	Donkey	1:100	Jackson Immunoresearch
α- mouse	TRITC	115-095-146	Goat	1:100	Jackson Immunoresearch
	TRITC	715-025-150	Donkey	1:100	Jackson Immunoresearch
	Alexa 555	A-32727	Goat	1:200	ThermoFisher
	FITC	115-095-146	Goat	1:100	Jackson Immunoresearch
	Alexa 488	A-11001	Goat	1:200	ThermoFisher
	AMCA	115-155-146	Goat	1:100	Jackson Immunoresearch
α-rabbit	TRITC	711-025-152	Donkey	1:100	Jackson Immunoresearch
	Alexa 555	A-31572	Donkey	1:200	ThermoFisher
	FITC	711-095-152	Donkey	1:100	Jackson Immunoresearch
	Alexa 488	A-32731	Goat	1:200	ThermoFisher
	Alexa 488 - Fab	111-547-003	Goat	1:100	Jackson Immunoresearch
	AMCA	711-155-152	Donkey	1:100	Jackson Immunoresearch
α-sheep	Texas Red	713-075-147	Donkey	1:100	Jackson Immunoresearch

3.6. Adquisición y análisis de imágenes

Los esparcidos se visualizaron a temperatura ambiente con un microscopio Leica DM6000 B con el objetivo de 63x. Las imágenes se capturaron con una cámara Hamamatsu ORCA-ER C4742-80. Las IFs de aplastados de túbulos seminíferos se analizaron con un microscopio Delta Vision con objetivos 100x. Todas las imágenes se procesaron con OPENLAB 4.0.3 y Adobe Photoshop CS6. Las preparaciones de los diferentes genotipos utilizadas para los análisis comparativos se prepararon en fresco y de forma paralela para todos ellos. Las IFs también se realizaron en paralelo y con la misma mezcla de Abs preparada en fresco. Las preparaciones no se congelaron para evitar diferencias en la señal de fondo y la inmunoreactividad. Todas las imágenes fueron adquiridas con exposiciones constantes que permitan la comparación directa. La cuantificación del número de foci y de la intensidad de fluorescencia se llevó a cabo con el software Fiji (ImageJ). Solo se cuantificaron los foci asociados a ejes cromosómicos y para la cuantificación de la intensidad de fluorescencia se sustrajo la señal de fondo. Para los análisis de co-localización, el mismo núcleo se cuantificó sin rotar (experimento) y tras rotar una de las imágenes 90 grados (control). Esta condición permite determinar la co-localización no específica (aleatoria).

3.7. Microscopía de super-resolución

Para generar imágenes de super-resolución se utilizó microscopía STED (*Stimulated emission depletion microscopy*, SP8, Leica). Las IFs se llevaron a cabo sobre esparcidos de espermatozoides de la forma previamente explicada. Sin embargo, los Abs secundarios utilizados para imágenes STED estaban conjugados con fluorocromos Alexa 555 y Alexa 488 (Tabla VI). Las preparaciones se montaron con medio de montaje ProLong Gold Antifade sin DAPI. Las imágenes se adquirieron y procesaron con el software LAS X (Leica).

3.8. Electroporación *in vivo* de testículos

Esta técnica descrita por primera vez por el Dr. Muramatsu (Jaroszeski et al. 2003) permite expresar en testículo cDNAs clonados en vectores de expresión de manera transitoria usando una cirugía menor. Para mejorar la eficiencia, la electroporación se llevó a cabo en ratones ICR a 16 dpp o en B6 a los 20-30 dpp. Tras anestesiar a los ratones por inhalación con isoflurano, los testículos se sacaron cuidadosamente de la cavidad abdominal y con ayuda de un capilar de vidrio se inyectaron 10 μ l de la solución de DNA en la *rete testis*. Esta solución de DNA contenía el vector de expresión a una concentración de 5 μ g/ μ l en HBS 1x (*HEPES buffered saline*: 20 mM HEPES, 140 mM NaCl, 5 mM KCl, 0,1% glucosa, 0,7 mM $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$) y estaba teñida con FastGreen (Sigma-Aldrich). El FastGreen da color verde y permite visualizar si la solución entra correctamente en el testículo. Se añade justo antes de la inyección en proporción de 1 μ l de FastGreen 0,1% por cada 10 μ l de DNA. Una hora tras la inyección, durante la cual el DNA penetra en los túbulos seminíferos, se procedió a la electroporación utilizando un electroporador CUY21 BEX (BEX Ltd). Para ello, se colocó el testículo entre el par de electrodos, se humedeció con una gota de PBS y se aplicaron 4 pulsos eléctricos (de 35V y 50 ms cada uno) en cada dirección. Finalmente, los testículos se introdujeron cuidadosamente en la cavidad abdominal y se suturó la incisión. Entre 24 y 72 horas tras la electroporación se sacrificaron los ratones y se hicieron esparcidos o aplastados de los testículos y se analizaron mediante IF.

4. Cultivos celulares

4.1. Líneas celulares y condiciones de cultivo

Durante el desarrollo de esta Tesis Doctoral se han utilizado las siguientes líneas celulares:

- HEK 293T: derivada de la línea tumoral de riñón embrionario HEK 293 (Human Embryonic Kidney) transformadas con el antígeno T del virus SV40.
- COS7: línea obtenida de riñón de mono verde africano transformada con el antígeno T del virus SV40 y con apariencia similar a la de los fibroblastos.
- U2OS: línea celular epitelial derivada de un osteosarcoma humano.

Todas estas líneas celulares se cultivaron en placas de cultivo (BD Falcon) en incubadores a 37°C y una atmósfera húmeda con un contenido de CO_2 del 7%. El medio de cultivo utilizado fue DMEM (Dulbecco's Modified Eagle Medium, GIBCO) suplementado con 10% de FBS y 1% de PSG (Penicilina-Estreptomicina-Glutamina, Gibco). Para tripsinizar las células, cuando alcanzaron confluencia o iban a ser sembradas, se lavaron con PBS 1x para eliminar los restos de FBS y, tras ello, se trataron con tripsina al 0,05% (GIBCO)

durante 2-5 min a 37°C. La tripsina se neutralizó añadiendo medio con FBS, y se disgregaron las células de forma mecánica con la pipeta. Finalmente, se contaron las células en una cámara de Neubauer, y se sembró el número de células deseado en función del experimento a realizar.

Para la conservación a largo plazo, las células se congelaron en medio de cultivo (DMEM) suplementado con 20% de FBS y 10% de DMSO (dimetilsulfóxido, Sigma-Aldrich), descendiendo lentamente la temperatura hasta -70°C en un recipiente de congelación con Isopropanol (MrFrosty™, Thermo Scientific™). Finalmente, las células se transfirieron a tanques de nitrógeno líquido (-180°C).

4.2. Transfección de líneas celulares

Para la transfección de las diferentes líneas se sembraron un día antes de la transfección a la densidad adecuada para alcanzar al día siguiente una confluencia del 70-80%. El número de células a sembrar para esto depende del tamaño de las células con las que se esté trabajando (Tabla VII). Estas se transfectaron con el plásmido/s de interés utilizando jetPEI (Polyplus-Transfection) según las recomendaciones del fabricante. Se preparó una primera solución con el DNA diluido en NaCl 150 mM en las proporciones adecuadas. Por otro lado, se diluyó el jetPEI en NaCl 150 mM. Esta última solución se añadió sobre la solución de DNA, se mezcló con *vortex* y se incubó durante 20-30 minutos a temperatura ambiente. Posteriormente se añadió la mezcla de transfección sobre las células. En el caso de las HEK 293T/COS7 se hizo gota a gota repartida por toda la placa de cultivo y se homogeneizó. Las células se incubaron 24 horas a 37°C y se reemplazó el medio de cultivo por medio fresco. En el caso de las U2OS la mezcla de transfección se mezcló con medio fresco (1/4 del volumen total de medio en la placa) y este se añadió sobre las células. En las transfecciones de U2OS se cambió el medio de transfección a las 4 h lavando dos veces las células y añadiendo medio fresco.

Tabla VII. Número de células a sembrar el día previo para conseguir una confluencia óptima para la transfección en función del tipo celular.

Línea celular	Placa	Nº de células
HEK293T	P3,5	0,5X10 ⁶
	P6	1,2 X10 ⁶
	P10	4x10 ⁶
U2OS	P3,5	0,27X10 ⁶

5. Análisis estadístico

Los datos numéricos de este trabajo han sido representados como media $\pm$ desviación estándar. Con el fin de comparar datos entre diferentes genotipos en distintas etapas, usamos el Análisis Welch *t*-test (*t*-test con varianzas desiguales). Esta prueba se consideró la apropiada ya que los datos de las cuantificaciones no están muy desviados, es decir, se aproximan razonablemente a una distribución normal y, en la mayoría de los casos, los grupos presentan varianzas desiguales. Se aplicó una prueba *two-sided* (dos colas) en todos los casos. Los asteriscos indican la significatividad estadística: *p valor <0,05, **p valor <0,01, ***p valor <0,001, ****p valor <0,0001 y sin asterisco: p valor > 0,05, no significativo. El análisis se realizó con GraphPad Prism

RESULTADOS

1. USP26 no es esencial para la fertilidad en ratón

La espermatogénesis, representa uno de los programas de diferenciación más complejos de la biología del desarrollo siendo necesaria una regulación espacio-temporal muy precisa de la transcripción, la traducción y la renovación proteica. En este sentido, la ubiquitinación resulta esencial para su correcto desarrollo ya que no solo se encarga de marcar los sustratos para su degradación por el proteasoma sino que, además, puede modificar su actividad, su estabilidad o su localización subcelular (He et al. 2016). De hecho, la importancia de la ubiquitinación en procesos específicos de la espermatogénesis como la MSCI y la determinación de los COs, así como en la fertilización y la calidad del espermatozoide, ya ha sido ampliamente reportada (Baarends et al. 1999; Baarends et al. 2000; Muratori et al. 2005; Yi et al. 2007; Qiao et al. 2014; Rao et al. 2017).

El gen *Usp26* (Ubiquitin-specific protease 26) se identificó por primera vez a partir de un cribado de genes expresados en espermatogonias de ratón (Wang et al. 2001). USP26 pertenece a una familia de cisteín-proteasas que eliminan largas cadenas de ubiquitina de sus sustratos siendo responsables del mantenimiento de la homeostasis de la ubiquitina en las células (Komander et al. 2009). En humanos, *USP26* ha despertado un gran interés como gen candidato de infertilidad masculina debido a su expresión altamente restringida a los testículos y por su localización en el cromosoma X, cromosoma enriquecido en genes específicos de testículo y para el cual los hombres son hemigóticos (Nishimune and Tanaka 2006). De forma más específica, tanto en ratón como en humano, la expresión de *USP26* se ha descrito en espermatogonia (tipos A y B), espermatoцитos en preleptotena, leptotena y zigotena, espermátidas redondas y en la barrera hemato-testicular (Wang et al. 2005; Lin et al. 2011). Además, en humanos también se ha descrito su expresión en células de Sertoli y Leyding (Wosnitzer et al. 2014).

El debate sobre la existencia de una asociación directa entre polimorfismos en *USP26* e infertilidad masculina ha suscitado gran controversia en las últimas dos décadas. Por un lado, existen estudios que asocian causalmente polimorfismos en *USP26* con casos de infertilidad masculina, desde Síndrome de sólo células de Sertoli hasta azoospermia no obstructiva o astenozoospermia (Paduch et al. 2005; Stouffs et al. 2005; Zhang et al. 2007; Lee et al. 2008a; Asadpor et al. 2013; Xia et al. 2014; Li et al. 2015; Li et al. 2020a; Arafat et al. 2020). Por el contrario, otros estudios han descrito la presencia de esos y otros polimorfismos en *USP26* con la misma frecuencia tanto en hombres fértiles como infértiles (Ravel et al. 2006; Stouffs et al. 2006; Christensen et al. 2008; Ribarski et al. 2009; Shi et al. 2011; Zhang et al. 2015; Luddi et al. 2016). Además, se ha reportado que la mayoría de los polimorfismos identificados no suprimen la actividad enzimática de USP26 (Zhang et al. 2015; Liu et al. 2018).

Con estos antecedentes y dada la ausencia de estudios funcionales, decidimos llevar a cabo el análisis funcional de USP26 mediante la generación de un modelo murino de pérdida de función.

1.1. Generación de los modelos murinos mutantes para *Usp26*

Con el fin de analizar la función de USP26 en la gametogénesis, generamos un modelo de pérdida de función dirigiendo el sistema CRISPR/Cas9 al único exón codificante de *Usp26*. El modelo generado presenta dos inserciones consecutivas de 85 y 19 bp, separadas entre ellas 5 bp. La primera inserción cambia la pauta y genera un codón de STOP prematuro en la posición 123 (proteína de 14 KDa) dando lugar a un alelo que, a priori, se predice nulo (nombrado como alelo²¹; Figura 1a-b). Para validar la pérdida de función del alelo analizamos la expresión de USP26 en testículos de ratón con dos anticuerpos (Ab, antibodies) diferentes, un α USP26 comercial (W-12) y el α USP26-M proporcionado por el Dr. Yi-Wen Lin (Lin et al. 2011). En primer lugar, llevamos a cabo un análisis por inmunofluorescencia (IF) en esparcidos de espermatocitos procedentes de ratones silvestres (WT, *Wild-Type* o *Usp26*^{WT/Y}) y mutantes (*Usp26*^{21/Y}). Ninguno de los Abs produjo señal detectable ni en espermatocitos (leptotena, zigotena) ni en espermatogonia o espermátida redonda (Figura 2a), tipos celulares en los que se ha descrito la expresión de USP26 (Wang et al. 2005; Lin et al. 2011). También analizamos la expresión de USP26 por western blot (WB) en extractos totales de testículo procedentes de ratones WT y mutantes. Con el Ab α USP26 W-12 no obtuvimos ninguna banda mientras que con el α USP26-M observamos bandas similares en los extractos *Usp26*^{21/Y} y *Usp26*^{WT/Y} (Figura 2b). Aunque el tamaño esperado para la proteína mutante es de 14 KDa, no se produjeron cambios en la migración de las bandas entre los extractos mutantes y los WT (banda esperada 95 KDa). Estos resultados sugieren firmemente que los Abs utilizados no son capaces de reconocer la proteína endógena y, por lo tanto, las bandas detectadas no corresponden con USP26.

Figura 1. Generación del alelo mutante 21 mediante CRISPR/Cas9. (a-b) Representación esquemática de la estrategia de edición genómica utilizada. Cuadros grises: exones codificantes, blancos: no codificantes. Los cromatogramas muestran las secuencias de la región editada en ratones WT y *Usp26*^{21/21}. ATG, codón de inicio, TAG y *, codones de stop, TGC (Cisteína catalítica, Cys295). Las líneas gruesas (codificantes) y finas (no codificantes) bajo los exones indican la transcripción esperada desde el alelo WT y el alelo nulo. **(b)** Genotipado por PCR de ratones WT, heterocigoto (WT/21) y homocigoto para la mutación (21/21) en *Usp26* con los primers F1 y R1. Se detecta una banda de 411 bp para el alelo WT y una de 307 bp para el alelo 21.

Las mutaciones que generan codones de STOP cerca del inicio de la traducción (5') de una proteína pueden dar lugar a un reinicio de la traducción desde otros codones AUG en pauta que estén aguas abajo del codón de inicio canónico (Makino et al. 2016). En ratón, el locus genómico de *Usp26* codifica para una metionina en pauta en la posición 147 que podría actuar como inicio de traducción alternativo en nuestro modelo mutante (alelo²¹). A partir de esta metionina se podría generar una proteína truncada en el N-terminal (79 KDa, Figura 1a) que presentaría intacto el dominio catalítico. Aunque el análisis por WB de los ratones *Usp26*^{21/Y} no detectó ninguna de las proteínas truncadas predichas (14 KDa-STOP prematuro y 79 KDa-ATG de reinicio, Figura 2b), con los Abs empleados no podemos descartar su expresión. Por ello, dada la imposibilidad de demostrar que el alelo²¹ es un alelo nulo, decidimos generar un segundo mutante con una delección en su dominio proteolítico (mutante inactivo). Este dominio está formado por una tríada catalítica compuesta por un residuo principal, la cisteína catalítica, acompañada de una histidina y una asparagina/aspartato que facilitan el ataque nucleofílico de la cisteína (Nijman et al. 2005; Davis and Simeonov 2015). Para ello, dirigimos el sistema CRISPR/Cas9 a la cisteína catalítica (Cys295) generando una delección de 86 bp que incluía este residuo (nombrado como alelo^{Cat}; Figura 3a-b). Además, esta delección produce un cambio de pauta que genera un codón de STOP en la posición 293 dando lugar, presumiblemente, a una proteína truncada de 34 KDa sin actividad catalítica (Figura 3a).

Figura 3. Generación del alelo mutante Cat mediante CRISPR/Cas9. (a-b) Representación esquemática de la estrategia de edición genómica utilizada. Cuadros grises: exones codificantes, blancos: no codificantes. Los cromatogramas muestran las secuencias de la región editada en ratones WT y *Usp26*^{Cat/Cat}. ATG, codón de inicio, TAG/TAA y *, codones de stop, TGC (Cisteína catalítica, Cys295). Las líneas gruesas (codificantes) y finas (no codificantes) bajo los exones indican la transcripción esperada desde el alelo WT y el alelo nulo. **(b)** Genotipado por PCR de ratones WT, heterocigoto (WT/Cat) y homocigoto para la mutación (Cat/Cat) en *Usp26* con los *primers* F2 y R2. Se detecta una banda de 259 bp para el alelo WT y una banda de 173 bp para el alelo mutante.

El análisis por IF y WB en los mutantes *Usp26*^{Cat/Y} reflejó los mismos resultados que en los mutantes *Usp26*^{21/Y}, es decir, una ausencia de señal detectable en la IF con ambos Abs (Figura 4a) y en WB con el Ab W-12 (Figura 4b). En el caso del α USP26-M el análisis por WB no mostró ningún cambio en la migración de las bandas ni se identificaron las proteínas truncadas putativas de 34 o 58 KDa del alelo^{Cat} (Figura 4b). Por lo tanto, aunque no podemos demostrar la ausencia de USP26 en nuestros mutantes, debido a la falta de Abs capaces de detectar la USP26 endógena, el análisis genético predice que ambos modelos son nulos. En el caso concreto del mutante catalítico, USP26 carece de la información genética que codifica el dominio catalítico activo (Cys295) siendo, por tanto, un mutante inactivo sin capacidad para proteolizar la ubiquitina de sus sustratos.

1.2. Los ratones deficientes en USP26 son fértiles

Los ratones portadores de los alelos 21 y Cat se desarrollaron con normalidad y no presentaron ningún fenotipo evidente. Además, el tamaño del testículo de los ratones mutantes fue similar al de los WT (Figura 5a). Con el fin de determinar el impacto de la ausencia de USP26 en la fertilidad del ratón, los machos *Usp26*^{WT/Y}, *Usp26*^{21/Y}, and *Usp26*^{Cat/Y} se cruzaron con hembras *Usp26*^{WT/WT}.

La presencia de tapón vaginal se monitorizó diariamente descartándose la presencia de defectos de comportamiento en estos animales y se anotó el número de crías obtenido en cada camada. Los resultados obtenidos no mostraron diferencias significativas en el tamaño de las camadas entre los tres grupos analizados (Tabla 1). En línea con este resultado, el análisis histopatológico de los túbulos seminíferos de los ratones *Usp26^{21/Y}* y *Usp26^{Cat/Y}* mostró una progresión normal a lo largo de los 12 estadios del ciclo del epitelio seminífero con una organización y número de los distintos tipos celulares similares a los WT (Figura 5b). Además, la cantidad de espermatozoides que llegaron al epidídimo no se vio afectada en los ratones mutantes (Figura 5b). Por lo tanto, en conjunto estos resultados sugieren que el impacto de USP26 en la fertilidad masculina en ratón, de existir, es muy reducido.

Resultados

		Media (descendientes)	SD	Camadas
Machos	<i>Usp26^{WT/Y}</i>	7,0	1,47	15
	<i>Usp26^{21/Y}</i>	6,9	2,31	18
	<i>Usp26^{Cat/Y}</i>	6,4	2,32	16
Hembras	<i>Usp26^{WT/WT}</i>	7,4	2,33	14
	<i>Usp26^{21/21}</i>	7,2	1,89	16
	<i>Usp26^{Cat/Cat}</i>	7,9	2,28	16

Tabla 1. Datos numéricos (media de crías por camada) obtenidos de los cruces de machos *USP26^{WT/Y}*, *USP26^{21/Y}*, and *USP26^{Cat/Y}* y hembras *USP26^{WT/WT}*, *USP26^{21/21}*, and *USP26^{Cat/Cat}* con hembras *USP26^{WT/WT}* y machos *USP26^{WT/Y}*, respectivamente. Se cruzaron ratones de 8 semanas durante un periodo de 3 meses (n= 6 ratones/genotipo).

En relación con las hembras, *USP26* no se expresa en ovario ni se han descrito hasta el momento mutaciones asociadas con infertilidad femenina. De acuerdo con esto, las hembras *Usp26^{21/21}* y *Usp26^{Cat/Cat}* no mostraron ningún defecto aparente en fertilidad (Tabla 1) ni alteraciones histológicas en los ovarios con respecto a las WT (Figura 5c).

Figura 5. El análisis histopatológico no muestra defectos en los ratones deficientes en *USP26*. (a) El tamaño de los testículos no se ve afectado por la ausencia de *USP26*. (b) Tinción PAS (Periodic Acid-Schiff)-hematoxilina de secciones de testículos *Usp26^{WT/Y}*, *Usp26^{21/Y}* y *Usp26^{Cat/Y}* mostrando en todos ellos una distribución normal de las células germinales en los túbulos seminíferos (TS) y una cantidad normal de espermatozoides en el epidídimo (Ep). Barras de escala 10 μ m (TS) y 20 μ m (Ep). (c) Tinción hematoxilina-eosina (H&E) de secciones de ovarios de hembras *Usp26^{WT/WT}*, *Usp26^{21/21}* y *Usp26^{Cat/Cat}* a los 4 meses de edad. En ausencia de *USP26* se observa la presencia de folículos y cuerpos lúteos y una apariencia normal del estroma. Barras de escala 50 μ m.

1.3. Los espermatoцитos deficientes en USP26 no muestran defectos en sinapsis, dinámica de las DSBs ni recombinación meiótica

Aunque *Usp26* comienza a transcribirse en espermatogonia, este gen alcanza su máximo nivel de expresión durante los primeros estadios de la meiosis (leptotena y zigotena, ver anexo Tablas S1 y S2 (Margolin et al. 2014; da Cruz et al. 2016)). Además, se ha demostrado la implicación del sistema ubiquitin-proteasoma en la regulación de la profase meiótica en ratón (Rao et al. 2017) y más concretamente de USP26 en la reparación de DSBs y la recombinación homóloga en células somáticas (Typas et al. 2016). Con estas evidencias decidimos analizar en los espermatoцитos que carecen de USP26 los principales procesos de la profase I meiótica : la sinapsis entre cromosomas, la generación y resolución de las DSBs y la recombinación meiótica.

Para evaluar la progresión de la sinapsis en nuestros mutantes, analizamos la distribución de SYCP3 y SYCP1 en esparcidos de espermatoцитos de ratones *Usp26*^{21/Y}, *Usp26*^{Cat/Y} y *Usp26*^{WT/Y} (Figura 6). La formación de los AEs fue normal empezando a ensamblarse en leptotena como pequeños fragmentos los cuales se fueron alargando a medida que avanzaba el estadio de zigotena. En paquitena se detectó una correcta sinapsis que abarcaba la longitud completa de los homólogos y no se observaron defectos en la desinapsis desde diplotena a diacinesis. Además, las metafases I y II de los mutantes presentaron una morfología completamente normal (Figura 6). Por lo tanto, estas observaciones sugieren que USP26 no desempeña ninguna función en el establecimiento de la sinapsis y la dinámica cromosómica.

Figura 6. La ablación de USP26 no produce defectos en la sinapsis y la progresión de la meiosis. Doble IF de SYCP1 y SYCP3 en esparcidos de espermatoцитos de ratones *USP26*^{WT/Y}, *USP26*^{21/Y} y *USP26*^{Cat/Y}. El DNA se tiñó con DAPI. No se observan diferencias en los procesos de sinapsis y desinapsis desde leptotena a diplotena, ni defectos en las divisiones meióticas (metafases I y II). Barra de escala 10 μ m.

Resultados

Dada la importancia de la ubiquitinación y sumoilación en los procesos de reparación de las DSBs (Bekker-Jensen and Mailand 2011), decidimos analizar la dinámica de estas roturas en los espermatoцитos deficientes en USP26 mediante el marcaje de γ H2AX, primera modificación detectada tras la formación de las DSBs (Rogakou et al. 1998). La distribución de γ H2AX en leptotena fue muy similar en los espermatoцитos mutantes y los WT observándose en todos los casos una señal intensa en todo el núcleo, indicativo de una correcta formación de las DSBs. Como consecuencia del procesamiento de las DSBs, en los tres genotipos la señal fue disminuyendo a medida que los espermatoцитos alcanzaron el estadio de zigotena. Finalmente, en paquitena la señal se observó restringida al *sex body* indicando una correcta reparación de las DSBs de los autosomas (Figura 7). Estos datos sugieren que la formación y reparación de las DSBs no se ven afectadas por la ausencia de USP26.

Figura 7. La dinámica de las DSBs no se ve afectada por la ausencia de USP26. Doble IF de SYCP3 y γ H2AX en esparcidos de espermatoцитos de ratones *USP26^{WT/Y}*, *USP26^{21/Y}* y *USP26^{Cat/Y}*. El DNA se tiñó con DAPI. No se observan diferencias desde leptotena hasta paquitena indicando una correcta formación y reparación de las DSBs. Los gráficos bajo los paneles muestran la cuantificación de la intensidad de fluorescencia de γ H2AX en los diferentes estadios y genotipos (n= 60 espermatoцитos). Análisis Welch *t*-test de dos colas: ns, no significativo. Barra de escala 10 μ m.

A medida que avanza la profase I, las DSBs van siendo procesadas resolviéndose en la mayor parte de los casos como NCO. Solo una pequeña fracción de ellas se procesarán como CO, dando lugar a la formación de los quiasmas. Teniendo en cuenta la localización del sistema ubiquitin-proteasoma en el SC y su función en la regulación de la tasa de recombinación meiótica y formación de COs (mediada por RNF212 y HEI10) (Qiao et al. 2014; Rao et al. 2017), decidimos evaluar este proceso en los espermatoцитos mutantes para *Usp26*

mediante la tinción de MLH1, proteína *mismatch repair* que marca los futuros COs (Guillon et al. 2005). No se observaron diferencias en el número de foci de MLH1 entre los espermatoцитos $Usp26^{21/Y}$, $Usp26^{Cat/Y}$ y $Usp26^{WT/Y}$ (Figura 8) indicando que la ausencia de USP26 no afecta a la recombinación meiótica.

En conjunto, estos resultados sugieren que en ratón la ausencia de USP26 no afecta a los eventos fundamentales de la profase I meiótica masculina ni tampoco produce una disminución de la fertilidad en ninguno de los sexos, indicando que USP26 no es esencial para la gametogénesis de ratón.

Figura 8. Los ratones deficientes en USP26 no presentan defectos en recombinación meiótica. Doble IF de SYCP3 y MLH1 en esparcidos de espermatoцитos de ratones $Usp26^{WT/Y}$, $Usp26^{21/Y}$ y $Usp26^{Cat/Y}$. El DNA se tiñó con DAPI. El número de foci de MLH1 (futuros quiasmas) es muy similar en los ratones mutantes y los WT. El gráfico a la derecha del panel muestra la cuantificación del número de foci en paquitenas para cada genotipo (n= 48-52 paquitenas). Análisis Welch t-test de dos colas: ns, no significativo. Barra de escala 10 μ m.

2. Una variante en HSF2BP causa POI a través de su nuevo interactor BRME1

2.1. La variante S167L en el gen *HSF2BP* es una firme candidata de POI en humanos

La POI es una de las principales causas de infertilidad femenina y afecta en torno al 1-3% de las mujeres menores de 40 años estimándose que las causas genéticas se encuentran detrás de un 20% de los casos (Nelson 2009; Rossetti et al. 2017; Golezar et al. 2019). Gracias a las estrategias de genética directa e inversa con organismos modelo, fundamentalmente el ratón, se han identificado múltiples genes y rutas moleculares implicados en la regulación de la fertilidad, pero son pocos los casos en los que se ha podido establecer una relación causal directa con la fertilidad en humanos (Schimenti and Handel 2018). Esto se debe, principalmente, a que las variantes que causan infertilidad en humanos tienen poca probabilidad de propagarse en la población, lo cual dificulta su identificación. Sin embargo, en ocasiones estas variantes pueden identificarse mediante el estudio de pedigrís familiares, especialmente cuando existe consanguinidad. Estos casos proporcionan conocimientos fundamentales sobre la función de los genes y los mecanismos moleculares afectados. En la última década, mediante secuenciación del exoma completo de pedigrís afectados por POI, se han identificado variantes patogénicas en varios genes meióticos tales como SYCE1, HFM1, MSH5, MEIOB y SPO11 (Wang et al. 2014; De Vries et al. 2014; Guo et al. 2017; Caburet et al. 2019; Tran and Schimenti 2019). Una de estas variantes, identificada en el gen *STAG3*, fue descrita en un trabajo llevado a cabo por nuestro laboratorio en colaboración con los laboratorios del Dr. Reiner A. Veitia y el Dr. Eric Vilain (Caburet et al. 2014).

Teniendo en cuenta estos antecedentes y la experiencia e interés de nuestro grupo en la determinación de las causas genéticas subyacentes a la infertilidad en humanos abordamos, en colaboración con el grupo del Dr. Reiner A. Veitia (Institut Jacques Monod, Université de Paris, France), el estudio de una familia consanguínea afectada por POI. En esta familia de origen árabe israelí, los padres son primos en primer grado y de las cinco hijas, tres están afectadas por POI. Las tres pacientes tuvieron la menarquia a una edad normal (13-14 años) aunque con menstruaciones irregulares que cesaron alrededor de los 25 años (amenorrea secundaria). De ellas, solo una pudo tener un hijo tras un tratamiento de fertilidad asistida (Figura 9a). Con el fin de identificar las bases genéticas de este caso familiar de POI, se llevó a cabo la secuenciación del exoma completo de dos de las pacientes con POI (III-2 y III-3) y de una hermana fértil (III-10). Las variantes encontradas se filtraron siguiendo los siguientes criterios (Ver anexo, Tablas S3 y S4):

1. Su presencia en homocigosis en las pacientes afectadas.
2. Su ausencia o presencia en heterocigosis en la hermana fértil.
3. Su ausencia en los controles fértiles no relacionados con la familia.
4. Una frecuencia alélica menor a 0,01 en todas las bases de datos analizadas.

Este proceso de filtrado condujo a la identificación de una mutación con cambio de sentido (*missense*) en el gen *HSF2BP*: rs200655253 (21:43630396 G>A, GRCh38). Esta mutación se localiza en el exón 6 del transcrito de referencia ENST00000291560.7 (NM_007031.2:c.500C>T) y cambia el codón TCG a TTG (NP_008962.1:p.Ser167Leu). Esta variante es muy infrecuente (Frecuencia de la Variante Alélica/VAF 0,0001845 en la base de datos GnomAD y 0,0005 en la base de datos GME Variome, específica de poblaciones de Oriente Medio) y está ausente en homocigosis en todas las bases de datos disponibles.

Mediante secuenciación Sanger se verificó la variante y se comprobó su segregación mendeliana con la POI dentro de la familia. Todas las hermanas afectadas (III-1, III-2 y III-3) fueron homocigotas para la mutación mientras que los padres y los hermanos fértiles fueron portadores heterocigotos (Figura 9b). No se identificaron varones homocigotos en la familia impidiendo el análisis del impacto de esta variante en la fertilidad masculina. La serina mutada (Ser167) está altamente conservada y 11 de los 18 predictores de patogenicidad disponibles en dbNSFP 3.5 predicen la variante como patogénica o deletérea (ver anexo, Tabla S5, Figura S1 y Figura S2). Estos análisis fueron llevados a cabo por el grupo del Dr. Reiner A.Veitia y los resultados obtenidos sugieren que la variante S167L en *HSF2BP* podría ser la responsable del desarrollo de POI en esta familia.

Figura 9. Identificación de la variante S167L en *HSF2BP* en una familia consanguínea afectada por POI. (a) Pedigrí de la familia de estudio. Cuando se conocen, se indican el año de nacimiento y la edad de la menarquia. La paciente III-3 tuvo un embarazo normal con ayuda de un tratamiento de fertilidad y un segundo intento fallido. El genotipo de cada individuo para la variante identificada en *HSF2BP* se indica en azul en base a lo determinado por secuenciación de Sanger para los DNAs disponibles. **(b)** Cromatogramas obtenidos por secuenciación Sanger de los familiares a los que no se realizó secuenciación del exoma completo. Se puede observar la segregación mendeliana de la variante.

Resultados

Al comienzo de este trabajo la única información disponible sobre HSF2BP (HSF2 Binding Partner) era su identificación como un interactor del factor de transcripción HSF2 (Heat Shock Factor 2) específico de testículo y su interacción con BNC1 (Basonuclin 1), ambas detectadas mediante cribado de doble híbrido en levaduras (Y2H, *Yeast Two Hybrid*) (Yoshima et al. 1998; Wu et al. 2013). La deficiencia de ambas proteínas se ha relacionado con problemas de fertilidad en ratón (Kallio et al. 2002; Ma et al. 2006). Esto, junto con el patrón de expresión altamente restringido a testículo de *HSF2BP* (proteintlas.org/ENSG00000160207-HSF2BP/summary/rna), nos llevó a considerar este gen como un buen candidato causal de la POI de la familia en estudio. Para analizarlo, en primer lugar, encargamos el desarrollo de Abs policlonales frente a la proteína HSF2BP murina (Proteogenix, ver material y métodos) y analizamos su localización en esparcidos de espermatoцитos y oocitos WT mediante IF (Figura 10a-b). HSF2BP mostró un patrón de localización típico de la maquinaria de recombinación meiótica formando foci sobre los ejes cromosómicos que empezaron a aparecer al inicio de zigotena y fueron aumentando durante este estadio para después disminuir a lo largo de paquitena y finalmente desaparecer totalmente en paquitena tardía (Figura 10a-b). Esta localización sugiere que HSF2BP podría ser una nueva proteína de recombinación meiótica y apoya la idea de que la variante *HSF2BP*-S167L es una firme candidata a producir la POI en la familia de estudio.

Figura 10. Localización de HSF2BP en meiocitos WT. (a-b) Doble IF con SYCP3 y HSF2BP en esparcidos de (a) espermatoцитos y (b) oocitos WT. En ambos casos HSF2BP muestra un patrón de foci localizados sobre los ejes cromosómicos desde zigotena temprana hasta el final de paquitena alcanzando su máximo nivel en zigotena tardía. Barras de escala, 10 μ m.

2.2. Generación de los modelos murinos mutantes para *Hsf2bp*

Con el fin de determinar si la variante *HSF2BP*-S167L es realmente la causante de POI en la familia de estudio y de analizar la función de esta nueva proteína en la meiosis, generamos dos modelos murinos mediante la tecnología de edición del genoma CRISPR/Cas9 (Figura 11a-b y material y métodos). El primero es portador de la variante causante de POI (*Hsf2bp*^{S167L/S167L}) y el segundo es un modelo de pérdida de función

de HSF2BP que permite un análisis comparativo directo entre el comportamiento de la variante S167L y el alelo nulo (KO, knockout o *Hsf2bp*^{-/-}; Figura 11b-c).

Figura 11. Generación y caracterización de los modelos murinos mutantes para *Hsf2bp*. (a-b) Representación esquemática de la estrategia de edición genómica utilizada para la generación del (a) modelo portador de la variante S167L y (b) el modelo nulo. Las guías del sistema CRISPR (sgRNAs) se localizan en el exón 6. Los cuadros grises representan los exones codificantes y los blancos los no codificantes. Los cromatogramas muestran las secuencias de la región editada en ratones WT, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-}. ATG, codón de inicio, TGA/TAA y *, codones de stop. (a) El ssODN lleva la mutación deseada (TCA>TTA, S171L) en el residuo homólogo al de la variante asociada con POI en humanos (S167L) aunque en todo el trabajo se denomina al alelo murino con la nomenclatura del residuo humano (S167L). (b) Las líneas gruesas (secuencias codificantes) y las finas (no codificantes) localizadas bajo los exones indican la transcripción esperada desde el alelo WT y el alelo nulo. (c) Genotipado por PCR de ratones WT (+/+), heterocigoto (+/-) y homocigoto para el alelo nulo (-/-) de *Hsf2bp* con los primers F1 y R1. Se detecta una banda de 383 bp para el alelo WT y una banda de 355 bp para el alelo mutante. (d) WB de extractos de testículo procedentes de ratones de 13 dpp (WT, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-}) con un Ab frente a HSF2BP. Como control de carga se empleó Tubulina. (e) Doble IF frente a SYCP3 y HSF2BP en espermatocitos *Hsf2bp*^{+/+} y *Hsf2bp*^{-/-}. Esta IF, junto con el WB, demuestran la ausencia de HSF2BP en el modelo nulo, así como la especificidad de los Abs generados frente a HSF2BP. (f) Doble IF frente a SYCP3 y EGFP-HSF2BP en espermatocitos tras la electroporación *in vivo* en testículo de EGFP-HSF2BP mostrando un patrón punteado muy similar al obtenido con los Abs frente al HSF2BP endógeno (panel e). Barras de escala, 10 μ m.

La validación del modelo nulo se realizó por WB e IF de extractos y preparaciones de testículo obtenidas de ratones *Hsf2bp*^{+/+} y *Hsf2bp*^{-/-} utilizando los Abs previamente generados frente a HSF2BP. En ambos casos se observó la ausencia de proteína en estos ratones en comparación con los ratones WT (Figura 11d-e) demostrando que se trata realmente de un alelo nulo y validando la especificidad de estos Abs. Esta validación también se llevó a cabo mediante electroporación *in vivo* de un plásmido que codifica para EGFP-HSF2BP en los testículos de ratón. Transcurridas 48 horas tras la electroporación, EGFP-HSF2BP se detectó en forma de foci sobre los ejes cromosómicos desde zigotena hasta paquitena (Figura 11f). Esta localización coincide con la observada con los Abs generados frente a HSF2BP.

2.3. HSF2BP es esencial para la fertilidad en ratón y la variante S167L produce una reducción parcial de la misma

Los ratones *Hsf2bp*^{-/-} y *Hsf2bp*^{S167L/S167L} se desarrollaron con normalidad y no mostraron ningún fenotipo aparente. Sin embargo, los machos *Hsf2bp*^{-/-} fueron estériles y las hembras presentaron una subfertilidad muy penetrante, cercana a la infertilidad (Figuras 12b y 13b). Con el fin de analizar el impacto de la variante S167L en la fertilidad del ratón, establecimos cruces de hembras y machos *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{+/+} con machos y hembras *Hsf2bp*^{+/+}, respectivamente. La presencia de tapón vaginal se monitorizó diariamente y se anotaron las camadas obtenidas y el número de crías en cada una de ellas. Ambos sexos fueron capaces de reproducirse, pero presentaron cierta tendencia hacia una reducción en la fertilidad, siendo esta más evidente en el caso de las hembras en las que se observó una reducción significativa en el número de camadas (Figura 12a). Estos resultados sugieren que la ausencia de HSF2BP produce defectos severos en la fertilidad del ratón y la variante S167L tiene cierto impacto en la misma.

El análisis histológico de los ovarios mostró una drástica reducción de la reserva folicular en las hembras *Hsf2bp*^{-/-}. Sin embargo, en los ovarios *Hsf2bp*^{S167L/S167L} no se observaron diferencias aparentes en la morfología ni en el número de folículos en comparación con los ovarios WT (Figura 12b-c). En machos, los testículos de los ratones *Hsf2bp*^{-/-} mostraron una reducción de tamaño del 70% con respecto a los testículos WT, representado como la relación peso del testículo/peso corporal. Esta reducción también se observó, aunque menos acusada, en los ratones *Hsf2bp*^{S167L/S167L} (reducción del 21%, Figura 13a). De acuerdo con esto, el análisis histológico de los testículos *Hsf2bp*^{-/-} reveló un bloqueo en el estadio IV del ciclo del epitelio seminífero acompañado por una ausencia total de espermatozoides en el epidídimo (Figura 13b). Los túbulos bloqueados en este estadio se caracterizan por la presencia de una apoptosis masiva de espermatoцитos con apariencia similar a zigonemas conjuntamente con espermatogonias intermedias (In) dividiéndose a espermatogonias B (Ahmed and de Rooij 2009). Estos defectos dan lugar a una azoospermia no obstructiva y la consiguiente infertilidad en estos ratones. En el caso de los testículos *Hsf2bp*^{S167L/S167L} se observó un bloqueo parcial con túbulos seminíferos que presentaban un elevado número de divisiones meióticas en apoptosis y una menor cantidad de espermatozoides en el epidídimo (Figura 13b). Conforme con estos resultados, los cortes procedentes de ratones *Hsf2bp*^{S167L/S167L} presentaron un mayor número de células positivas para la tinción TUNEL (marcaje de apoptosis, *TdT mediated dUTP Nick End Labelling*) que correspondían siempre con divisiones meióticas (Figura 13c). Además, la cuantificación de espermatozoides en el epidídimo corroboró la existencia de una reducción del 23% en el número de espermatozoides de los ratones *Hsf2bp*^{S167L/S167L} con respecto a los *Hsf2bp*^{+/+} (Figura 13d). Por último, en ninguno de los mutantes se

vio alterada la presencia de espermatogonias, células de Sertoli y células de Leydig (Figura 13b). En conjunto, estos resultados indican que HSF2BP es esencial para la fertilidad del ratón y su ausencia produce un fenotipo sexualmente dimórfico. Además, el fenotipo observado en los ratones *Hsf2bp*^{S167L/S167L}, con evidentes defectos en espermatogénesis y una reducción significativa de la fertilidad en hembras, sugiere que los ratones portadores de la variante causante del POI fenocopian parcialmente la enfermedad humana.

Figura 12. Análisis de fertilidad y caracterización de los ovarios en los mutantes para *Hsf2bp*. (a) Análisis de fertilidad de hembras y machos *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{+/+} mostrando el número de camadas por mes y el número de crías por camada. n=6 ratones por genotipo (excepto en hembras *Hsf2bp*^{S167L/S167L}; n=7). (b) Tinción de H&E en secciones de ovarios adultos (8 semanas). Mientras los ovarios *Hsf2bp*^{-/-} presentan una fuerte depleción del pool de folículos, los ovarios *Hsf2bp*^{+/+} muestran una apariencia normal. Barras de escala, 100 µm. (c) Cuantificación del número de folículos por ovario en hembras de 5 dpp, 6 semanas y 13 meses. n (5 dpp)= 5 ovarios *+/+* y *S167L/S167L*, 4 *-/-*; n (6 semanas) = 5 ovarios *+/+* y *S167L/S167L*, 3 *-/-*; n (13 meses)= 3 ovarios *+/+* y *S167L/S167L*. Análisis Welch t-test de dos colas: ns, no significativo, *p<0,05; **p<0,01; ***p<0,001.

Figura 13. Análisis y caracterización de los defectos en la espermatogénesis de los mutantes para *Hsf2bp*. (a) Tamaño de los testículos de ratones $Hsf2bp^{S167L/S167L}$ y $Hsf2bp^{-/-}$ en comparación con los correspondientes WT. El gráfico representa la cuantificación del peso del testículo referido al peso corporal de cada animal (%). $n = 14$ testículos $+/+$, 12 $S167L/S167L$ y 6 $-/-$. (b) Tinción de PAS-hematoxilina en secciones de testículo de los tres genotipos. Los asteriscos azules marcan las divisiones meióticas en apoptosis y las flechas indican la apoptosis masiva observada en el KO. Barras de escala, 10 μm (paneles de arriba) y 20 μm (paneles de abajo). TS, Túbulos Seminíferos; Ep, Epidídimo. (c) Detección inmunohistoquímica (TUNEL) de células apoptóticas en secciones de testículo mostrando el incremento de apoptosis en divisiones meióticas observadas en túbulos en estadio XII en los ratones $Hsf2bp^{S167L/S167L}$. El gráfico a la derecha del panel representa la cuantificación. $n = 3$ ratones por genotipo. Barra de escala, 25 μm . (d) Cuantificación del número de espermatozoides presentes en el epidídimo de ratones $Hsf2bp^{S167L/S167L}$ y $Hsf2bp^{+/+}$ adultos. $n = 8$ epidídimos por genotipo. Análisis Welch t-test de dos colas: ns, no significativo, * $p < 0,05$; ** $p < 0,01$; *** $p < 0,001$; **** $p < 0,0001$.

2.4. La ausencia de HSF2BP produce fuertes defectos en sinapsis y la variante S167L provoca una pérdida parcial de función

A la vista de los defectos en gametogénesis y fertilidad de ambos mutantes nos propusimos caracterizar en detalle la progresión de la meiosis en los mismos. Para ello, en primer lugar analizamos el ensamblaje/desensamblaje del SC en los meiocitos de estos ratones mediante el seguimiento de la distribución de las proteínas SYCP1 y SYCP3. De acuerdo con el bloqueo en estadio IV del ciclo epitelial seminífero, los machos $Hsf2bp^{-/-}$ mostraron un bloqueo en un estadio que se asemeja al de zigotena (zigotena-like) (Figura 14a). Muchos de los espermatocitos presentaban un fenotipo denominado *de cambio de pareja* (*partner-switch phenotype*), frecuentemente observado en mutantes con defectos en recombinación (Romanienko and Camerini-Otero 2000; Kumar et al. 2010). En estos espermatocitos los AEs

se alargan correctamente, pero no son capaces de aparearse y sinapsar con su homólogo y, finalmente, establecen una sinapsis aberrante entre regiones de cromosomas no homólogos (Figura 14a). En el caso de los espermátocitos *Hsf2bp*^{S167L/S167L} no se observaron defectos en la sinapsis y desinapsis desde leptotena hasta diplotena (Figura 14b). Sin embargo, observamos un elevado número de células apoptóticas que mediante tinción con SYCP3 y ACA (marcaje de centrómeros) mostraron ser metafases. Estas células presentaban una pérdida total o parcial de la inmunoreactividad frente a los Abs utilizados, característica de las células apoptóticas (Figura 14c). Estos resultados son acordes al bloqueo parcial observado en el análisis histológico (Figura 13b-c).

Figura 14. Análisis de la sinapsis en los machos mutantes de *Hsf2bp*. (a-b) Doble IF con SYCP1 y SYCP3 en espermátocitos (a) *Hsf2bp*^{-/-} y (b) *Hsf2bp*^{S167L/S167L}. La variante S167L no produce defectos en sinapsis mientras que la ausencia de HSF2BP da lugar a un bloqueo en estadio zigoteno-like. (c) Doble IF con SYCP3 y ACA (centrómeros) en aplastados de túbulos seminíferos de ratones WT y S167L. Los ratones S167L presentan un elevado número de metafases apoptóticas con una pérdida parcial o total de inmunoreactividad de las proteínas marcadas. Barras de escala, 10 μ m.

En hembras, los oocitos *Hsf2bp*^{-/-} mostraron un retraso en la progresión de la profase I. A 17,5 días post-coito (dpc) casi todos los oocitos *Hsf2bp*^{+/+} habían alcanzado ya el estadio de paquitena mientras que la mayor parte de los oocitos *Hsf2bp*^{-/-} de esta misma edad se encontraban en zigotena. Además, se observó un aumento en el porcentaje de oocitos con defectos en sinapsis en ausencia de HSF2BP (*Hsf2bp*^{-/-}: 45,5% $\pm$ 1,5 frente a las *Hsf2bp*^{+/+}: 7,5% $\pm$ 1,5; n=2 por genotipo; **p<0,01; Figura 15a). Sin embargo, en las hembras *Hsf2bp*^{S167L/S167L} no observamos ningún defecto en sinapsis/desinapsis (Figura 15b). En conjunto, estos resultados sugieren que la ausencia de HSF2BP produce defectos severos en sinapsis y que la variante S167L presenta una pérdida parcial de función (alelo hipomorfo).

Figura 15. Análisis de la sinapsis en hembras mutantes para *Hsf2bp*. (a-b) Doble IF de SYCP1 y SYCP3 en esparcidos de oocitos de hembras (a) *Hsf2bp*^{-/-} y (b) *Hsf2bp*^{S167L/S167L}. De igual manera que en machos, las hembras *Hsf2bp*^{S167L/S167L} no muestran defectos en sinapsis durante la profase I pero las hembras *Hsf2bp*^{-/-} muestran un retraso en la progresión de la misma y un elevado número de células con defectos en sinapsis. Barras de escala, 10 µm.

2.5. La variante S167L produce una reducción de la expresión/estabilidad de la proteína

Para confirmar si la variante S167L se comporta como un alelo hipomorfo, analizamos la localización de HSF2BP en los meiocitos *Hsf2bp*^{S167L/S167L}. Tanto en espermatocitos como en oocitos, se observó una importante reducción de HSF2BP en los ejes cromosómicos tanto en número de foci como en su intensidad (Figura 16). Además, mediante WB de extractos de testículo de ratones de 13 dpp (día en el que todas las células están en zigotena dada la sincronía de la primera ronda meiótica) observamos que la reducción detectada por IF correlacionaba con una reducción en la expresión de la proteína en los ratones *Hsf2bp*^{S167L/S167L} (Figura 17). Estos resultados sugieren que la variante HSF2BP-S167L produce una reducción de la expresión y/o estabilidad de la proteína.

Figura 16. Localización de HSF2BP en los meiocitos *Hsf2bp*^{S167L/S167L}. Doble IF de SYCP3 y HSF2BP en esparcidos de espermatocitos (arriba) y oocitos (17,5 dpc, abajo) mostrando una fuerte reducción del marcaje de HSF2BP. Gráficos a la derecha de cada panel: cuantificación del número de foci y la intensidad de HSF2BP. n= 30-40 células de cada estadio y genotipo (u.a. unidades arbitrarias). Análisis Welch t-test de dos colas: ****p<0,0001. Barras de escala, 10 µm.

Figura 17. Expresión de HSF2BP en los testículos de los mutantes para *Hsf2bp*. WB frente a HSF2BP de extractos totales de testículo obtenidos de ratones de 13 dpp. Se observa la ausencia total de proteína en los ratones *Hsf2bp*^{-/-} y una reducción del 65% en los ratones *Hsf2bp*^{S167L/S167L}. Como control de carga se empleó Tubulina. Análisis Welch t-test de dos colas: *p<0,05.

2.6. HSF2BP es esencial para la recombinación meiótica y la variante S167L produce defectos severos en este proceso

Dado que HSF2BP presenta una localización similar a la maquinaria de recombinación meiótica decidimos analizar el comportamiento de diferentes proteínas implicadas en este proceso mediante IF en los meiocitos *Hsf2bp*^{-/-} y *Hsf2bp*^{S167L/S167L}.

En primer lugar, analizamos la dinámica de reparación de las DSBs mediante la detección de γ H2AX y RPA por IF. La fosforilación de H2AX es la primera modificación que aparece tras la inducción de las DSBs en leptotena y se observa como una señal difusa en la cromatina. A medida que las células avanzan a través de zigotena esta señal va disminuyendo, quedando restringida al *sex body* en paquitena. Sin embargo, en los espermatocitos *Hsf2bp*^{-/-} la señal de γ H2AX no disminuyó quedándose acumulada en todo el núcleo de las zigonemas-like bloqueadas (Figura 18a). Por su parte, el 80% de las paquinemas *Hsf2bp*^{S167L/S167L} presentaron un intenso marcaje de γ H2AX sobre los autosomas mientras que en los ratones WT solo se detectó este patrón en un 9% de ellas (Figura 18a). En ambos mutantes, esta acumulación de γ H2AX correlacionó perfectamente con la acumulación de la proteína de unión a ssDNA RPA. Así, en los ratones *Hsf2bp*^{-/-} observamos niveles de RPA similares al WT en leptotena pero una acumulación significativa en el bloqueo en zigotena-like (Figura 18b). En los espermatocitos *Hsf2bp*^{S167L/S167L} observamos que, aunque el número de focos de RPA en leptotena y zigotena era similar al de los ratones WT, se producía una gran acumulación de foci intensos en paquitena (Figura 18b). De igual forma, en las hembras observamos una intensa acumulación de γ H2AX en los oocitos en paquitena del mutante nulo que fue menos pronunciada en los oocitos portadores de la variante S167L (Figura 18c). Sin embargo y sorprendentemente, no observamos acumulación de RPA en los oocitos de ninguno de los dos modelos mutantes, sugiriendo la existencia de mecanismos alternativos durante estos procesos de la recombinación meiótica en oocitos (Figura 18d). En conjunto, estos resultados indican que la variante S167L produce importantes defectos en la reparación de las DSBs que son similares a los producidos por la ausencia de HSF2BP, aunque mucho más leves.

Dados los defectos en reparación del DNA de los mutantes de *Hsf2bp*, a continuación, analizamos el comportamiento de las recombinasas RAD51 y DMC1, proteínas encargadas de la invasión del cromosoma homólogo durante la recombinación meiótica. El análisis por IF mostró que la ausencia de HSF2BP produce una drástica reducción, casi total, de la localización de ambas proteínas en los ejes cromosómicos en ambos sexos (Figura 19a-d). Esta reducción también se observó en menor medida en los ratones portadores de la variante causante de POI (Figura 19a-d). Por tanto, HSF2BP es necesaria para la localización de las recombinasas en los RNs tempranos y la variante S167L reduce sustancialmente esta función.

Figura 18. Reparación de las DSBs en los mutantes de *Hsf2bp*. (a-c) Doble IF de SYCP3 y γH2AX en espermátocitos de (a) espermátocitos y (b) oocitos (17,5 dpc) de ratones WT, *Hsf2bp*^{S167L/S167L} y (b) *Hsf2bp*^{-/-}. (b-d) Doble IF de SYCP3 y RPA en espermátocitos (b) espermátocitos y (d) oocitos (16,5 y 17,5 dpc) de ratones WT, *Hsf2bp*^{S167L/S167L} y (b) *Hsf2bp*^{-/-}. (a-c) En machos se observa una acumulación de ambas proteínas en ambos mutantes, siendo esta mayor en los ratones *Hsf2bp*^{-/-}. (b-d) En hembras, se produce una acumulación de γH2AX, aunque no de RPA, que es más acusada los oocitos *Hsf2bp*^{-/-}. Los gráficos a la derecha de los paneles representan las cuantificaciones en cada estadio y genotipo. Cuantificación % de paquitenas con acúmulos: n>360 paquitenas. Cuantificación intensidad γH2AX: (a) n= 53-60 células (c) n= 19-21 células (17,5 dpc). Cuantificación foci RPA: (b) n= 29-46 células (d) n= 23-42 células. Análisis Welch t-test de dos colas: ns, no significativo, *p<0,05; **p<0,01; ****p<0,0001. Barras de escala, 10 μm.

Figura 19. La carga de las recombinasas se encuentra afectada en los mutantes de *Hsf2bp*. (a-c) Doble IF de SYCP3 con (a,c) DMC1 o (b,d) RAD51 en esparcidos de (a,b) espermatozootos y (c,d) oocitos (17,5 dpc) de ratones *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-}. Se observa una fuerte reducción (*Hsf2bp*^{-/-}) y una reducción intermedia (*Hsf2bp*^{S167L/S167L}) del número de foci de ambas recombinasas en comparación con los niveles WT. Los gráficos debajo de cada panel representan la cuantificación de foci en cada estadio y genotipo. n (a y b)= 37-43 células, n (c)= 18-30, n (d)= 35-42. Análisis Welch t-test de dos colas: ns, no significativo; **p<0,01; ***p<0,001; ****p<0,0001. Barras de escala, 10 μ m.

A continuación, decidimos analizar el estado de SPATA22, proteína que forma un heterodímero con MEIOB y que a su vez interacciona con RPA participando en la protección del ssDNA tras la resección, así como en la captura del segundo extremo en los D-loops durante el procesamiento de los RNs intermedios (Luo et al. 2013; Xu et al. 2017). El análisis por IF de SPATA22 en los espermatozootos *Hsf2bp*^{-/-} mostró una

Resultados

intensa acumulación de esta proteína en los ejes cromosómicos que fue menor en presencia de la variante S167L (Figura 20). En los oocitos *Hsf2bp*^{-/-} también se observó una fuerte acumulación de SPATA22 mientras que en los *Hsf2bp*^{S167L/S167L} solo se produjo un ligero aumento no significativo del marcaje (Figura 20). Estos resultados concuerdan con la acumulación de RPA y la deslocalización de las recombinasas observados en nuestros mutantes. Los defectos en la carga/estabilización de las recombinasas provocarían defectos en el desplazamiento del complejo de unión a ssDNA RPA/SPATA22/MEIOB de las regiones de las DSBs provocando su acumulación.

Figura 20. Análisis de SPATA22 en los mutantes *Hsf2bp*. Doble IF de SYCP3 y SPATA22 en espermátocitos (arriba) y oocitos (17,5 dpc, abajo) de ratones *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-}. SPATA22 se acumula en los espermátocitos y oocitos KO y muestra una acumulación intermedia en los espermátocitos *Hsf2bp*^{S167L/S167L}. En los oocitos *Hsf2bp*^{S167L/S167L} solo se observa una ligera acumulación no significativa. Los gráficos bajo los paneles representan la cuantificación de la intensidad de SPATA22 en cada estadio y genotipo. n (machos)= 20 células, n (hembras)= 38-41 células. Análisis Welch *t*-test de dos colas: ns, no significativo; ***p*<0,01; *****p*<0,0001. Barras de escala, 10 μ m.

El objetivo final de la recombinación meiótica es la formación de al menos un CO por bivalente estableciéndose así las uniones físicas (quiasmas) entre los cromosomas homólogos necesarias para la correcta segregación durante la primera división meiótica. En vista de la acumulación de RPA/SPATA22 y los defectos en el reclutamiento de RAD51/DMC1, analizamos si todo ello alteraba la formación de los COs. De acuerdo con el bloqueo en zigotena-like, no se detectaron foci de MLH1 en los espermátocitos *Hsf2bp*^{-/-} (Figura 21a). En el caso de los espermátocitos *Hsf2bp*^{S167L/S167L}, se observó una reducción significativa en el número total de COs en comparación con los ratones WT (Figura 21a). Además, esta reducción estaba acompañada de un mayor porcentaje de bivalentes que no mostraban foci de MLH1, afectando en mayor medida al bivalente sexual (Figura 21b). Acorde con la reducción de los COs, los ratones *Hsf2bp*^{S167L/S167L} mostraron un mayor porcentaje de paquinemas con los cromosomas sexuales desinapsados (Figura 21c) y, consecuentemente, la presencia de metafases I con univalentes libres (Figura 21d). Estos resultados podrían explicar el elevado número de metafases apoptóticas observadas en estos mutantes (Figura 13b-c y 14b).

Figura 21. Formación de COs en los machos *Hsf2bp*^{S167L/S167L}. (a) Doble IF de SYCP3 y MLH1 en paquitenas WT, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-} mostrando una reducción de foci de MLH1 en los mutantes S167L y la ausencia total en los ratones nulos. El gráfico bajo el panel muestra la cuantificación del número total de foci. n = 60-89. (b) Cuantificación del % de espermatoцитos con algún autosoma o el bivalente sexual sin foci de MLH1. n (autosomas) = 90-94, n (XY) = 59-64. (c) Doble IF de SYCP3 y γH2AX en paquitenas WT y S167L permitiendo identificar claramente el *sex body*. El gráfico a la derecha representa el % de paquitenas con el bivalente XY sinapsado/desinapsado. n = 150. (d) IF con SYCP3 (DNA teñido con DAPI) mostrando los univalentes observados en las metafases I de los ratones *Hsf2bp*^{S167L/S167L} (flechas). Análisis Welch *t*-test de dos colas: **p* < 0,05; ***p* < 0,01; ****p* < 0,0001. Barras de escala, 10 μm.

En los oocitos la cuantificación de los COs se llevó a cabo usando CDK2 como marcador. Durante la profase I, CDK2 se localiza en los telómeros de los cromosomas desde leptotena hasta diplotena. Sin embargo, adicionalmente, en paquitenas aparecen señales intersticiales de CDK2 que colocalizan con MLH1 y marcan los sitios de CO participando en la regulación de la formación de los mismos (Ashley et al. 2001; Palmer et al. 2020). Por tanto, como medida del número de COs en oocitos se consideraron únicamente los foci intersticiales de CDK2. Esta cuantificación mostró una importante reducción en el número total de COs en las hembras *Hsf2bp*^{-/-} con respecto a las WT que fue menor en las hembras *Hsf2bp*^{S167L/S167L} (Figura 22a). Además, el porcentaje de bivalentes sin foci intersticiales de CDK2 se vio fuertemente aumentado en los oocitos *Hsf2bp*^{S167L/S167L} y, aún más, en los oocitos *Hsf2bp*^{-/-} (Figura 22b).

Figura 22. Formación de COs en las hembras mutantes para *Hsf2bp*. (a) Doble IF de SYCP3 y CDK2 en oocitos (17,5 dpc) WT, *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-} mostrando una reducción de foci intersticiales de CDK2 en las mutantes S167L más acusada en ausencia de HSF2BP. El gráfico bajo el panel muestra la cuantificación del número total de foci. n= 60-89. (b) Cuantificación del % de oocitos con algún bivalente sin foci intersticiales de CDK2. n = 46-79. Análisis Welch t-test de dos colas: *p<0,05; ***p<0,001; ****p<0,0001. Barras de escala, 10 µm.

En conjunto, todos estos resultados indican que HSF2BP es esencial para la carga de las recombinasas RAD51/DMC1 y, por tanto, para la recombinación meiótica. Además, la comparativa directa con el modelo nulo demuestra que la variante S167L tiene efectos menos severos que la ausencia de proteína, apoyando la hipótesis de que esta variante se comporta como un alelo hipomorfo. Por último, aunque tanto los machos como las hembras *Hsf2bp*^{S167L/S167L} presentan alteraciones moleculares similares en la ruta de recombinación meiótica, en el ámbito reproductivo su efecto es diferente en ambos sexos.

2.7. Los defectos meióticos producidos por la variante S167L no están mediados por BRCA2

Una vez determinada la función de HSF2BP en meiosis y caracterizadas las alteraciones meióticas que produce la presencia de la variante S167L en ratón, profundizamos en el mecanismo por el cual esta mutación produce los defectos descritos. Recientemente, se ha demostrado que HSF2BP interacciona con BRCA2 (Brandsma 2016), una proteína esencial para la recombinación meiótica (Sharan et al. 2004; Martinez et al. 2016). Esta interacción involucra al menos a la Arg200 de HSF2BP y a la región Gly2270-Thr2337 del fragmento C-terminal de BRCA2 (comunicación personal) por lo que nos preguntamos si la variante S167L en HSF2BP afectaría a esta interacción. Dada la imposibilidad de detectar la proteína BRCA2 endógena en nuestros ratones debido a la ausencia de Abs funcionales, llevamos a cabo ensayos *in vitro* de interacción/colocalización entre HSF2BP y el extremo C-terminal de BRCA2 (BRCA2-C) en sistemas heterólogos (células HEK293T y U2OS). En primer lugar, llevamos a cabo un ensayo de co-inmunoprecipitación (co-IP) entre BRCA2-C y las dos variantes de HSF2BP (WT y S167L) mediante la transfección en células HEK293T de estas proteínas marcadas con diferentes epítomos (Flag o GFP). Los resultados mostraron que BRCA2-C co-inmunoprecipita de manera similar con HSF2BP-WT y HSF2BP-S167L

(Figura 23a). A continuación, mediante IF en células U2OS transfectadas con GFP-BRCA2-C y Flag-HSF2BP analizamos el patrón de localización de cada una de ellas en ausencia o en presencia de la otra. En las transfecciones simples ambas variantes de HSF2BP mostraron una localización homogénea en el núcleo y en el citoplasma mientras que BRCA2-C mostró una localización únicamente nuclear. Cuando se co-transfectaron BRCA2-C y HSF2BP, su localización cambió drásticamente hacia un patrón punteado en el núcleo (Figura 23b). Sin embargo, cualitativa y cuantitativamente, este cambio de patrón fue similar en presencia de ambas variantes de HSF2BP (Figura 23b-c). Ambos resultados, sugieren que los efectos de la variante HSF2BP-S167L observados no están mediados directamente por una alteración de la interacción con BRCA2-C o una deslocalización de la misma.

Figura 23. Comparativa de la interacción entre BRCA2-C y las dos variantes de HSF2BP. (a) co-IP de BRCA2-C con ambas variantes mostrando una interacción similar en ambos casos. Las células HEK293T se transfectaron con Flag-HSF2BP (WT y S167L) y GFP-BRCA2-C. Los extractos proteicos se incubaron ON con los Abs indicados en verde y se analizaron mediante WB con los Abs indicados en rojo. (b) Doble IF con Abs frente a EGFP y Flag en células U2OS transfectadas con los plásmidos indicados a la izquierda. Se observa el cambio de patrón de EGFP-BRCA2-C en presencia de HSF2BP, siendo similar con ambas variantes. Barra de escala, 20 μm. (c) Cuantificación de porcentaje de células con focos nucleares detectadas en las transfecciones de U2OS. n>450 células. Análisis Welch *t*-test de dos colas: ns, no significativo; *p<0,05; ***p<0,001.

2.8. BRME1, un nuevo interactador de HSF2BP

Con el fin de profundizar en el mecanismo subyacente a la patogenicidad de la variante HSF2BP-S167L, nos propusimos identificar proteínas que interaccionen con HSF2BP mediante un *screening* de Y2H utilizando una librería de cDNAs (DNA complementario) de testículo de ratón. La secuenciación de los clones obtenidos mostró que 19 de los 98 clones correspondían a un gen no caracterizado denominado 4930432K21Rik en ratón (C19ORF57 en humano) y ahora renombrado como BRME1 (Break Repair Meiotic recombinase recruitment factor 1). Este nuevo interactador putativo de HSF2BP tiene una longitud de 600 aminoácidos con un alto contenido en residuos ácidos, no tiene dominios funcionales reconocibles y presenta una estructura intrínsecamente desordenada. Esta interacción se validó mediante co-IP de ambas proteínas obtenidas por transfección transitoria de plásmidos que codifican para Flag-HSF2BP y GFP-BRME1. Además, con este ensayo comprobamos que ambas variantes (HSF2BP-WT y HSF2BP-S167L) interaccionaban de manera

similar con BRME1 (Figura 24a) sugiriendo que la mutación causante de POI no altera la interacción entre ambas proteínas. A continuación, validamos la interacción *in vivo* mediante co-IP de ambas proteínas a partir de extractos totales de testículo de ratón con Abs específicos frente a cada una de las proteínas (Figura 24b). Los Abs frente a BRME1 fueron generados por encargo por Proteogenix (ver material y métodos).

Figura 24. BRME1 es un nuevo interactor de HSF2BP. (a) co-IP de BRME1 con ambas variantes mostrando una interacción similar en ambos casos. Las células HEK293T se transfectaron con Flag-HSF2BP (WT y S167L) y GFP-BRME1. Los extractos proteicos se incubaron ON con los Abs indicados en verde y se analizaron mediante WB con los Abs indicados en rojo. (b) IP de extractos de testículo con Abs (verde) frente BRME1, HSF2BP e IgGs como control negativo y detección por WB con los Abs indicados en rojo.

2.9. BRME1 es un nuevo componente de los RNs que co-localiza con HSF2BP

Tras demostrar la interacción entre BRME1 y HSF2BP tanto *in vitro* como *in vivo*, nos preguntamos cuál sería su función en meiosis y si, de igual manera que HSF2BP, formaría parte de los RNs meióticos. Para responder a esta pregunta llevamos a cabo un análisis mediante IF en esparcidos de espermatoцитos y oocitos de ratones WT utilizando los Abs policlonales generados frente a BRME1. Esta proteína se detectó asociada a los ejes cromosómicos desde leptotena hasta paquitena mostrando un patrón de foci discretos típico de los RNs (Figura 25a-b). De acuerdo con los resultados obtenidos en los experimentos de Y2H y co-IP, BRME1 co-localizó perfectamente con HSF2BP en los ejes (Figura 25c), co-localización que fue verificada mediante microscopía de super-resolución (Figura 25d). La cuantificación de esta co-localización se realizó mediante el conteo del número de foci positivos para HSF2BP solo, para BRME1 solo o para ambos a la vez. Como control negativo se realizó este mismo conteo girando una de las dos imágenes 90 grados con respecto a la otra, método que permite calcular el porcentaje de co-localización inespecífica o aleatoria. Los resultados mostraron una co-localización de HSF2BP y BRME1 del 90-95% en zigotena tardía y paquitena temprana (Tabla 2). La menor co-localización de BRME1 con HSF2BP observada en zigotena temprana (Tabla 2) podría explicarse por una menor sensibilidad de los Abs frente a HSF2BP en comparación con los de BRME1 o bien por una carga más tardía de HSF2BP a los RNs.

Figura 25. BRME1 forma parte de los RNs donde co-localiza con HSF2BP. (a-b) Doble IF de espermatozoides de (a) espermatozoides y (b) oocitos de ratones WT con un Ab policlonal generado frente BRME1 (proteína completa recombinante) y SYCP3, mostrando que BRME1 se localiza en los RNs desde leptotena hasta paquítena intermedia alcanzando la máxima señal en zigotena tardía. (c) Triple IF de SYCP3, BRME1 y HSF2BP en espermatozoides de ratones WT mostrando la alta co-localización de BRME1 y HSF2BP especialmente en zigotena tardía y paquítena (ver la cuantificación en la Tabla 2). (d) Doble IF de BRME1 y HSF2BP en espermatozoides de ratones WT analizada mediante microscopía de super-resolución (STED, Stimulated emission depletion). Barras de escala, 10µm (a,b y c), 5µm (d).

ESTADIO	Foci HSF2BP positivos para BRME1		Foci BRME1 positivos para HSF2BP	
	Específica (sin rotación)	Inespecífica (con rotación)	Específica (sin rotación)	Inespecífica (con rotación)
Zigotena temprana	91,5% ± 2,0	4,1% ± 1,3	68,7% ± 1,0	3,1% ± 1,0
Zigotena tardía	94,1% ± 1,2	3,1% ± 0,7	88,0% ± 3,0	2,9% ± 0,7
Paquítena temprana	92,6% ± 2,6	2,8% ± 1,7	89,6% ± 2,8	2,7% ± 1,7

Tabla 2. Cuantificación de la co-localización entre BRME1 y HSF2BP. La cuantificación se llevó a cabo en tres células de cada estadio.

Dada la localización de BRME1 y HSF2BP en los RNs desde profase temprana (leptotena-zigotena temprana) hasta paquítena, decidimos analizar su co-localización con dos proteínas clásicas de los RNs que muestran una localización ligeramente diferente a lo largo de la profase, RPA y DMC1. De acuerdo con la fuerte co-localización entre BRME1 y HSF2BP y la función de HSF2BP en reparación de las DSBs y recombinación meiótica, ambas proteínas co-localizaron con los foci de RPA y DMC1 (Figura 26a-b). Sin embargo, HSF2BP y BRME1 mostraron una mejor co-localización espacio-temporal con RPA que con DMC1 (Tabla 3), posiblemente debido a la restringida localización temporal de DMC1 que es eliminada rápidamente

tras la invasión del homólogo (Moens 2002). En conjunto, estos resultados sugieren que BRME1 es una nueva proteína de los RNs con un patrón altamente similar al de HSF2BP y RPA que, probablemente, tiene alguna función en la reparación de las DSBs y la recombinación meiótica.

Figura 26. Co-localización de BRME1 y HSF2BP con RPA y DMC1. (a-b) Triple IF de BRME1 (izda) y HSF2BP (dcha) con SYCP3 y (a) RPA o (b) DMC1 en espermátocitos WT. La localización de BRME1 y HSF2BP es más similar a la de RPA que a la de DMC1 a lo largo de la meiosis. Ver cuantificaciones en Tabla 3. Barras de escala, 10µm.

2.10. BRME1 interacciona con HSF2BP a través de su extremo C-terminal

A continuación, abordamos el análisis de la región de interacción de BRME1 con HSF2BP. Para ello, y dada la ausencia de dominios reconocibles en BRME1, dividimos arbitrariamente la proteína en tres fragmentos (N-terminal, región central y C-terminal) y generamos 3 plásmidos que codificaban para versiones de BRME1 que carecían de cada uno de estos fragmentos (Figura 27). Mediante la sobreexpresión de HSF2BP junto con cada una de estas 3 versiones de BRME1 acoplada a co-IP, observamos que únicamente en ausencia del extremo C-terminal no se producía la interacción, indicando que en esa región reside el dominio de interacción de BRME1 con HSF2BP (residuos 475-600 de la proteína murina, Figura 27).

ESTADIO	Foci BRME1 positivos para RPA2		Foci RPA2 positivos para BRME1	
	Específica (sin rotación)	Inespecífica (con rotación)	Específica (sin rotación)	Inespecífica (con rotación)
Zigotena temprana	90,5% ± 2,3	2,2% ± 0,9	84,6 ± 3,9	2,1% ± 0,8
Zigotena tardía	90,2% ± 1,5	2,0% ± 0,4	95,2% ± 0,6	2,1% ± 0,4
Paquitena temprana	87,7% ± 5,0	1,8% ± 0,2	92,4% ± 4,8	1,9% ± 0,3

ESTADIO	Foci HSF2BP positivos para RPA1		Foci RPA1 positivos para HSF2BP	
	Específica (sin rotación)	Inespecífica (con rotación)	Específica (sin rotación)	Inespecífica (con rotación)
Zigotena temprana	76,3% ± 2,0	5,5% ± 4,4	69,2% ± 0,8	5,0% ± 4,1
Zigotena tardía	81,8% ± 3,1	3,0% ± 0,9	82,4% ± 3,4	3,0% ± 1,0
Paquitena temprana	79,7% ± 3,5	2,8% ± 1,7	88,6% ± 3,5	3,1% ± 2,0

ESTADIO	Foci BRME1 positivos para DMC1		Foci DMC1 positivos para BRME1	
	Específica (sin rotación)	Inespecífica (con rotación)	Específica (sin rotación)	Inespecífica (con rotación)
Zigotena temprana	58,3% ± 4,6	5,1% ± 2,4	55,5% ± 5,0	4,9% ± 2,5
Zigotena tardía	30,8% ± 2,6	2,7% ± 1,0	78,8% ± 13,7	7,0% ± 1,7

ESTADIO	Foci HSF2BP positivos para DMC1		Foci DMC1 positivos para HSF2BP	
	Específica (sin rotación)	Inespecífica (con rotación)	Específica (sin rotación)	Inespecífica (con rotación)
Zigotena temprana	74,0% ± 11,1	1,0% ± 0,5	87,3% ± 8,1	1,2% ± 0,7
Zigotena tardía	55,3% ± 13,4	1,9% ± 1,1	89,1% ± 6,1	3,3% ± 2,5

Figura 27. Determinación de la región de interacción de BRME1 con HSF2BP. Esquema de la proteína BRME1 completa y las construcciones generadas con diferentes deleciones (Δ). En verde se muestran las construcciones que interactúan y en rojo la que no. Los WB bajo el esquema muestran las co-IPs con cada construcción. Las células HEK293T se transfectaron con GFP-HSF2BP y las distintas construcciones de BRME1 unidas a Flag. Los extractos proteicos se incubaron ON con los Abs indicados en verde y se analizaron mediante WB con los Abs indicados en rojo. La deleción $\Delta 475-600$ es la única que elimina la interacción, indicando que la región C-terminal de BRME1 es la esencial para su interacción con HSF2BP.

A la vista de este resultado, analizamos si el resto de BRME1 era esencial para su función mediante el desarrollo de un modelo murino *Brme1* ^{$\Delta 142-472/\Delta 142-472$} que expresa esta proteína sin su parte central (330 aminoácidos, más de la mitad de la proteína). Estos ratones fueron fértiles y no mostraron deslocalización de BRME1 (Figura 28a-d). Además, no presentaron ningún defecto en la sinapsis ni ninguna alteración en la localización de HSF2BP en los ejes cromosómicos (Figura 28e-f). Estos resultados indican que una gran fracción de la proteína codificante de BRME1 no es esencial para la función de BRME1/HSF2BP *in vivo*.

Figura 28. Los ratones *Brme1*^{Δ142-472/Δ142-472} no muestran defectos en meiosis. **(a)** Esquema de la estrategia de edición genómica utilizada para la generación del modelo *Brme1*^{Δ142-472/Δ142-472}. Los cuadros grises representan los exones codificantes y los blancos los no codificantes. Las líneas gruesas (codificantes) y finas (no codificantes) localizadas bajo los exones indican la transcripción esperada desde cada alelo. El cromatograma muestra la secuencia de la región editada. ATG, codón de inicio, TAA y *, codones de stop. **(b)** Genotipado por PCR de ratones WT (+/+), heterocigoto (+/Δ) y homocigoto para el alelo delta (Δ/Δ) de *Brme1*. **(c)** WB de extractos de testículo procedentes de ratones WT y *Brme1*^{Δ142-472/Δ142-472} con un Ab frente a BRME1. BRME1 se detecta a 130 KDa (tamaño esperado 65 KDa), muy probablemente debido a su alto contenido de residuos ácidos. De igual manera, BRME1 Δ142-472 debería detectarse en torno a 30 KDa pero aparece más alta. Tubulina, control de carga. **(d)** Doble IF de SYCP3 y BRME1 en espermatocitos de ratones WT y *Brme1*^{Δ142-472/Δ142-472}. *In vivo* BRME1 Δ142-472 se localiza en los RNs de igual manera que la proteína completa. **(e)** Doble IF de SYCP3 y SYCP1 en espermatocitos de ratones WT y *Brme1*^{Δ142-472/Δ142-472} mostrando la ausencia de defectos en sinapsis en los mutantes. **(f)** Doble IF de SYCP3 y HSF2BP en espermatocitos WT y *Brme1*^{Δ142-472/Δ142-472} mostrando que la ausencia de la región central de BRME1 no altera la localización de HSF2BP. Barras de escala, 10 μm.

2.11. HSF2BP es necesaria para la localización/estabilidad de BRME1

Teniendo en cuenta la interacción entre BRME1 y HSF2BP y la fuerte co-localización de ambas en los RNs, nos propusimos determinar si la localización de BRME1 sería dependiente de HSF2BP. Para ello llevamos a cabo un análisis por IF en los meiocitos *Hsf2bp*^{-/-} y *Hsf2bp*^{S167L/S167L}. Tanto en machos como en hembras, los resultados mostraron una ausencia total de señal de BRME1 en los ratones deficientes en HSF2BP y una reducción significativa del número de foci en los ratones *Hsf2bp*^{S167L/S167L} en comparación con los WT (Figura 29a-b). Además, los foci detectados en los espermatoцитos portadores de la variante S167L presentaban una intensidad significativamente menor (Figura 29a). El análisis de la expresión de BRME1 mediante WB de extractos de testículo de ratones de 13 dpp mostró una fuerte reducción en los testículos *Hsf2bp*^{S167L/S167L} que fue mayor en los testículos *Hsf2bp*^{-/-} (Figura 29c). Estos resultados sugieren que HSF2BP es necesaria no sólo para la localización de BRME1 en los RNs sino también para su expresión y/o estabilidad.

Figura 29. Análisis de BRME1 en los mutantes de *Hsf2bp*. (a-b) Doble IF de SYCP3 y BRME1 en espermatoцитos y (b) oocitos (17,5 dpc) procedentes de ratones *Hsf2bp*^{S167L/S167L} y *Hsf2bp*^{-/-} mostrando una fuerte reducción en los mutantes S167L y la ausencia total en los KO. (a) Los gráficos a la derecha del panel representan la cuantificación del número de foci y la intensidad de estos en cada estadio y genotipo. n= 15-20 células. (b) El gráfico bajo el panel representa la cuantificación del número de foci en paquitena. n= 18 células. Barras de escala, 10 µm. (c) WB de extractos totales de testículo obtenidos de ratones WT, *Hsf2bp*^{S167L/S167L}, *Hsf2bp*^{-/-} y *Brme1*^{-/-} de 13 dpp. Se observa la ausencia total de proteína en los ratones *Brme1*^{-/-} y una reducción importante en los mutantes de *Hsf2bp*, más acusada en el modelo nulo. Como control de carga se empleó Tubulina. Análisis Welch t-test de dos colas: *p<0,05; ****p<0,0001.

2.12. BRME1 y HSF2BP intervienen en los eventos tempranos tras la formación de DSBs

Para determinar de forma más específica el posicionamiento de BRME1 y HSF2BP en la ruta de recombinación meiótica, analizamos su patrón de distribución en distintos modelos murinos que carecen de proteínas relacionadas con la recombinación y la sinapsis. Estos mutantes presentan defectos de recombinación en diferentes puntos de la vía, lo cual permite acotar los procesos que son esenciales para la función de la proteína de interés. En concreto, los mutantes utilizados carecían de las siguientes proteínas: la cohesina REC8 (Bannister et al. 2004), la proteína del elemento central del SC SIX6OS1 (Gomez-H et al. 2016), las ligasas E3 RNF212 (SUMO) y HEI10 (Ubiquitina) implicadas en la estabilización de las proteínas de recombinación en los RNs tardíos, la proteína del espermatoproteasoma PSMA8 (Gomez-H et al. 2019) y la proteína encargada de la generación de las DSBs SPO11. Los mutantes para *Rnf212*, *Hei10* y *Spo11* se describen por primera vez en este trabajo y han sido validados comparando sus fenotipos (arresto meiótico, defectos en sinapsis y alteraciones moleculares) con los de los mutantes ya publicados (ver material y métodos). El análisis se llevó a cabo en zigotena tardía en los mutantes que presentaban un bloqueo posterior a este estadio, o en zigotena-like en aquellos mutantes que estaban bloqueados en ese punto.

Figura 30. La localización de BRME1 y HSF2BP depende de la formación de DSBs pero no de la sinapsis. (a-b) Doble IF de SYCP3 y (a)BRME1 o (b) HSF2BP en esparcidos de espermatocitos de ratones WT, *Rec8*^{-/-}, *Six6os1*^{-/-}, *Rnf212*^{-/-}, *Hei10*^{-/-}, *Psma8*^{-/-} y *Spo11*^{-/-}. Se observa la correcta localización en los mutantes con defectos en recombinación (de más a menos severos: *Rec8*^{-/-}, *Six6os1*^{-/-}, *Rnf212*^{-/-}, *Hei10*^{-/-} y *Psma8*^{-/-}) pero una ausencia total en los ratones *Spo11*^{-/-}, indicando una dependencia total de la formación de las DSBs y un posicionamiento genético en los eventos tempranos tras la formación de estas. Barras de escala, 10 μ m.

Los resultados mostraron que todos los mutantes analizados presentan una localización de BRME1 y HSF2BP similar a los ratones WT a excepción de los mutantes de *Spo11*, en los que no se detectó ninguna de las dos proteínas (Figura 30a-b). Estos resultados indican que la formación de DSBs por SPO11 es esencial

para la carga de BRME1 y HSF2BP a los RNs. Además, el hecho de que mutantes con bloqueos en los primeros estadios de la profase y defectos tempranos en la reparación de DSBs y la recombinación, como es el caso de *Rec8* y *Six6os1*, no tengan alterada la localización de BRME1/HSF2BP indica que su carga es independiente del establecimiento de la sinapsis y que, por tanto, este complejo estaría implicado en los eventos tempranos que ocurren tras la formación de DSBs.

2.13. BRME1 y HSF2BP no tienen capacidad de unión al DNA

Muchas de las proteínas de los RNs implicadas en recombinación meiótica y reparación de las DSBs presentan capacidad para unirse al DNA como es el caso de RPA, BRCA2, las recombinasas RAD51 y DMC1 o el complejo MEIOB/SPATA22 (Yang et al. 2002; Luo et al. 2013; Souquet et al. 2013; Hinch et al. 2020). Por esta razón, y teniendo en cuenta la dependencia de la formación de DSBs observada para la carga en los ejes de BRME1 y HSF2BP, analizamos si BRME1 y HSF2BP (WT o S167L) se unen directamente al DNA. Para ello expresamos ambas proteínas usando un sistema de transcripción/traducción *in vitro* (TnT) con el cual se obtienen proteínas altamente purificadas y libres de proteínas nucleares y cromatina (Loregian et al. 2004; Souquet et al. 2013). Estas proteínas se incubaron con bolas magnéticas cubiertas con biotina (control negativo), biotina-ssDNA (DNA de hebra simple) o biotina-dsDNA (DNA de doble hebra) de manera que si presentan capacidad de unión al DNA serán retenidas y podrán ser detectadas posteriormente por WB. Como control positivo utilizamos la proteína de unión a ssDNA RPA. Nuestros resultados mostraron que ninguna de las dos proteínas presenta capacidad de unión al DNA *in vitro* (Figura 31) y, por tanto, podría hipotetizarse que su carga a los RNs esté mediada por la interacción con proteínas intermedias como, por ejemplo, BRCA2.

Figura 31. BRME1 y HSF2BP no presentan capacidad de unión al DNA. (a-b) Las proteínas BRME1, HSF2BP (WT y S167L) y RPA1 (proteína de unión a ssDNA, control positivo) se produjeron unidas a Flag en un sistema libre de células (TnT) y se incubaron con bolas magnéticas cubiertas con biotina, biotina-ssDNA o biotina-dsDNA. Las proteínas retenidas se eluyeron y se analizaron por WB con un Ab frente a Flag. BRME1 y HSF2BP no unen DNA *in vitro*.

2.14. BRME1 es esencial para la fertilidad en ratón

Con el fin de analizar funcionalmente la implicación de BRME1 en la fertilidad del ratón y comprobar si su función es redundante con la de HSF2BP, generamos un mutante nulo de esta proteína (*Brme1*^{-/-}) mediante edición genómica con CRISPR/Cas9 (Figura 32a-b). La validación del modelo se realizó por WB e IF de extractos y preparaciones obtenidas de los testículos de ratones *Brme1*^{+/+} y *Brme1*^{-/-}. En ambos casos se

Resultados

observó la ausencia de proteína en estos ratones en comparación con los ratones WT (Figura 32c-d) demostrando que es un modelo nulo y validando la especificidad de los Abs generados frente a BRME1. De igual manera que en el caso de HSF2BP esta validación también se llevó a cabo mediante electroporación *in vivo* en testículos de ratón de un plásmido que codifica para EGFP-BRME1. Transcurridas 48 horas tras la electroporación, EGFP-BRME1 se detectó en forma de foci sobre los ejes cromosómicos desde zigotena hasta paquitena (Figura 32e). Esta localización coincide con la observada con los Abs generados frente a BRME1 (Figura 25a).

Figura 32. Generación y caracterización del modelo *Brme1*^{-/-}. (a) Representación esquemática de la estrategia de edición genómica utilizada para la generación del modelo. Cuadros grises: exones codificantes; cuadros blancos: no codificantes. El cromatograma muestra la secuencia de la región editada en los mutantes. ATG, codón de inicio, TAG/TAA y *, codones de stop. Las líneas gruesas (secuencias codificantes) y finas (no codificantes) localizadas bajo los exones indican la transcripción esperada desde el alelo WT y el alelo nulo. (b) Genotipado por PCR de ratones WT (+/+), heterocigoto (+/-) y homocigoto para el alelo nulo (-/-) de *Brme1* con los primers F2 y R3. Se detecta una banda de 646 bp (alelo WT) y una banda de 606 bp (alelo mutante). (c) WB de extractos de testículo procedentes de ratones *Brme1*^{+/+} y *Brme1*^{-/-} con un Ab frente a BRME1. Tubulina, control de carga. (d) Doble IF frente a SYCP3 y BRME1 en espermatocitos (izda) y oocitos (dcha) *Brme1*^{+/+} y *Brme1*^{-/-}. Esta IF, junto con el WB, demuestran la ausencia de BRME1 en el modelo nulo, así como la especificidad de los Abs generados frente a BRME1. (e) Doble IF frente a SYCP3 y EGFP en espermatocitos tras la electroporación *in vivo* en testículo de EGFP-BRME1 mostrando un patrón punteado idéntico al obtenido con los Abs frente al BRME1 endógeno (panel d). Barras de escala, 10 μ m.

Los ratones *Brme1*^{-/-} se desarrollaron con normalidad y no presentaron ningún fenotipo aparente. Sin

embargo, las hembras *Brme1*^{-/-} presentaron una subfertilidad muy severa, debido a una fuerte reducción de la reserva folicular (Figura 33a). Por su parte, los machos eran infértiles y mostraron una fuerte reducción del tamaño de los testículos (reducción del 76% con respecto a los WT, Figura 33b). El análisis histológico mostró un bloqueo meiótico en el estadio IV del ciclo epitelial seminífero con una presencia de un elevado número espermatoцитos apoptóticos y una ausencia de espermatozoides en el epidídimo (Figura 33c) lo que da lugar a una azoospermia no obstructiva.

Figura 33. La ausencia de BRME1 produce defectos de fertilidad en ratón. (a) Tinción de H&E en secciones de ovarios de hembras WT y *Brme1*^{-/-} (3 meses) mostrando la fuerte depleción de folículos en las hembras *Brme1*^{-/-}. El gráfico de la derecha representa la cuantificación del número de folículos en estas hembras, n= 3 ovarios/genotipo. Escala, 50 μ m. **(b)** Tamaño de testículo de los ratones *Brme1*^{+/+} y *Brme1*^{-/-}. El gráfico representa la cuantificación del peso del testículo referido al peso corporal de cada animal (%). n= 14 testículos +/+ y 6 -/-. **(c)** Tinción PAS-hematoxilina en secciones de testículo mostrando un arresto en estadio IV en los ratones *Brme1*^{-/-}. Los asteriscos rojos indican la apoptosis masiva de espermatoцитos. Escala, 10 μ m (arriba) y 20 μ m (abajo). TS, Túbulo Seminífero; Ep, Epidídimo. Análisis Welch t-test de dos colas: ns, no significativo, *p<0,05; ***p<0,001; ****p<0,0001.

A continuación, dados los defectos observados en el análisis histológico, nos propusimos analizar en detalle la progresión de la meiosis en ausencia de BRME1 siguiendo el ensamblaje/desensamblaje del SC. En los espermatoцитos *Brme1*^{-/-} el marcaje doble de SYCP3 y SYCP1 mostró una sinapsis parcial que en muchos casos ocurría entre cromosomas no homólogos (fenotipo de cambio de pareja) produciendo un bloqueo en zigotena-like (Figura 34a) muy similar al observado en los espermatoцитos *Hsf2bp*^{-/-} (Figura 14a). Una pequeña fracción de los espermatoцитos (3,7% $\pm$ 1,9; n= 3) fue capaz de escapar del bloqueo y alcanzar el estadio de paquitenia temprana. En el caso de las hembras *Brme1*^{-/-} el fenotipo también fue altamente similar

Resultados

al observado en ausencia de HSF2BP (Figura 15a) mostrando un retraso en la profase I y un elevado porcentaje de oocitos con defectos en sinapsis con diferente grado de severidad (*Brme1*^{-/-}: 47,9%±2,2 frente a las *Brme1*^{+/+}: 12%±5,7; n=2 por genotipo; *p<0,05; Figura 34b).

Figura 34. La ausencia de BRME1 produce defectos severos en sinapsis. (a-b) Doble IF con SYCP3 y SYCP1 en espermátocitos de (a) espermátocitos y (b) oocitos (18,5 dpc) de ratones *Brme1*^{+/+} y *Brme1*^{-/-}. (a) Los espermátocitos deficientes en BRME1 mostraron un bloqueo en zigotena-like en el que se observa sinapsis entre cromosomas no homólogos. (b) Los oocitos *Brme1*^{-/-} presentaron un elevado porcentaje de células con defectos en sinapsis con diferente grado de severidad. Barras de escala, 10 μ m.

2.15. La ausencia de BRME1 produce defectos en la recombinación meiótica

Una vez determinados los defectos en la progresión de la meiosis de los mutantes de *Brme1* y, teniendo en cuenta la alta similitud del fenotipo observado con el de los ratones *Hsf2bp*^{-/-} y su localización en los RNs llevamos a cabo un análisis detallado de la recombinación meiótica en ausencia de BRME1.

El marcaje de los espermátocitos *Brme1*^{-/-} con γ H2AX y RPA mostró una acumulación de ambas proteínas en las zigonemas-like bloqueadas (Figura 35a-b) indicando la presencia de DSBs no reparadas. Los oocitos *Brme1*^{-/-} presentaron una acumulación de γ H2AX en paquitena (Figura 35c) pero, pese a esta acumulación de daño, no se observó una acumulación de RPA (Figura 35d), de igual manera que lo observado en los oocitos mutantes para *Hsf2bp* (Figura 18d).

Figura 35. Defectos en la reparación de las DSBs en los meiocitos *Brme1*^{-/-}. (a-b) Doble IF de SYCP3 con (a-c) γH2AX o (b-d) RPA1 en espermátocitos de meiocitos *Brme1*^{+/+} y *Brme1*^{-/-}. (a-b) En los espermátocitos *Brme1*^{-/-} se observa una intensa acumulación de ambas proteínas en las células bloqueadas en zigotena-like. Nótese que en los ratones WT la señal de γH2AX desaparece en paquitena de los autosomas. Los gráficos bajo los paneles muestran la cuantificación en cada estadio y genotipo. n (γH2AX)= 20-30 células; n (RPA1)= 26-34 células. Barras de escala, 10 μm. (c-d) En los oocitos (17,5 dpc) *Brme1*^{-/-} se observa acumulación de γH2AX pero no de RPA en comparación con los WT. Los gráficos bajo los paneles muestran la cuantificación en cada estadio y genotipo. n (γH2AX)= 20-21 células; n (RPA1)= 22-42 células. Barras de escala, 10 μm. Análisis Welch t-test de dos colas: ns, no significativo; *p<0,05; **p<0,01; ****p<0,0001.

Resultados

El análisis de RAD51 y DMC1 en los meiocitos deficientes en BRME1 mostró una fuerte reducción del número de foci de ambas recombinasas (Figura 36a-d), sugiriendo que esta proteína es necesaria para su localización en los RNs. En línea con estas observaciones, los meiocitos *Brme1*^{-/-} presentaron una intensa acumulación de SPATA22 (Figura 37a-b), indicando que los defectos en la carga de las recombinasas y la invasión del homólogo previenen la eliminación de las proteínas que protegen las regiones de ssDNA como son RPA y el complejo MEIOB/SPATA22. Por último, el análisis de la formación de COs en espermatocitos mostró una ausencia total de foci de MLH1, de acuerdo con el bloqueo meiótico observado en estos mutantes (Figura 37c). En hembras, el número total de COs, medidos como foci intersticiales de CDK2, se vio reducido de manera significativa con respecto a los WT, observándose un mayor porcentaje de oocitos con bivalentes sin CO (Figura 37d). Estos resultados son muy similares, si no idénticos, a los descritos para los mutantes nulos de *Hsf2bp* (sección 2.6) y, por tanto, ambos análisis funcionales nos permiten concluir que HSF2BP y BRME1 son dos nuevas proteínas esenciales para la recombinación meiótica que actúan conjuntamente como un complejo heterodimérico funcional y cuya ausencia produce fenotipos altamente similares que incluyen un marcado dimorfismo sexual.

Figura 36. La ausencia de BRME1 provoca una fuerte reducción de la localización de las recombinasas RAD51 y DMC1. (a-b) Doble IF de SYCP3 con (a) RAD51 o (b) DMC1 en esparcidos de espermatocitos *Brme1*^{+/+} y *Brme1*^{-/-}. (c-d) Doble IF de SYCP3 con (a) RAD51 o (b) DMC1 en esparcidos de oocitos (17,5 dpc) *Brme1*^{+/+} y *Brme1*^{-/-}. En todos los casos se observa una disminución significativa del número de foci de recombinasas en los ratones *Brme1*^{-/-}. Los gráficos a la derecha de cada panel muestran la cuantificación en cada estadio y genotipo. n (a, RAD51 machos) = 37-45 células; n (b, DMC1 machos) = 41-50 células. n (c, RAD51 hembras) = 31-42 células; n (d, DMC1 hembras) = 18-31 células. Barras de escala, 10 µm. Análisis Welch *t*-test de dos colas: ****p<0,0001.

Figura 37. Defectos en recombinación en los ratones *Brme1*^{-/-}. (a-b) Doble IF de SYCP3 y SPATA22 en (a) espermatocitos y (b) oocitos (17,5 dpc) *Brme1*^{+/+} y *Brme1*^{-/-} mostrando una mayor intensidad SPATA22 en los mutantes. Los gráficos de la derecha muestran la cuantificación de la intensidad. n (machos)= 20 células; n (hembras)= 40 células. (c) Doble IF de SYCP3 y MLH1 en espermatocitos *Brme1*^{+/+} y *Brme1*^{-/-}, mostrando la ausencia de COs en los ratones *Brme1*^{-/-}. (d) Doble IF de SYCP3 y CDK2 en oocitos (17,5 dpc) *Brme1*^{+/+} y *Brme1*^{-/-} mostrando una reducción del número de COs (foci de CDK2 intersticiales) en las hembras mutantes. Los gráficos bajo el panel muestran la cuantificación del número total de foci y del % de oocitos con bivalentes sin CO. n= 49-79 células. Barras de escala, 10 µm. Análisis Welch t-test de dos colas: ns, no significativo, *p<0,05; ***p<0,001; ****p<0,0001.

2.16. BRME1 es necesaria para la localización/estabilidad de HSF2BP

Teniendo en cuenta la similitud de los fenotipos observados en ausencia de BRME1 y de HSF2BP, así como la deslocalización de BRME1 en los mutantes de *Hsf2bp*, decidimos comprobar si la localización de HSF2BP también era dependiente de BRME1. El análisis por IF de los espermatocitos *Brme1*^{-/-} reveló una importante reducción en el número de foci de HSF2BP con respecto a los *Brme1*^{+/+} (Figura 38a) mientras que en el caso de los oocitos *Brme1*^{-/-} se observó una ausencia total de foci de HSF2BP (Figura 38b). El análisis por WB de la expresión de HSF2BP en extractos de testículo de ratones de 13 dpp mostró una fuerte reducción en los ratones *Brme1*^{-/-} en comparación con la detectada en los WT (Figura 38c). Estos resultados indican que BRME1 es necesaria para la localización y la expresión/estabilidad de HSF2BP y junto con la reducción de BRME1 observada en ausencia de HSF2BP (Figura 29a-c), sugieren una fuerte interdependencia entre ambas proteínas.

Figura 38. Análisis de HSF2BP en los mutantes *Brme1*^{-/-}. (a-b) Doble IF de SYCP3 y HSF2BP en esparcidos de (a) espermatozootos y (b) oocitos (17,5 dpc) mostrando la fuerte reducción de la señal de HSF2BP en machos y una ausencia total en hembras. El gráfico de la derecha del panel (a) muestra la cuantificación del número de foci de HSF2BP en los espermatozootos *Brme1*^{-/-}, n = 30 células/genotipo. Barras de escala, 10 μ m. (c) WB de extractos totales de testículo

Dada la fuerte interdependencia de BRME1 y HSF2BP, decidimos analizar *in vitro* el comportamiento de ambas proteínas de forma individual y conjunta, mediante su transfección (o co-transfección) en células U2OS. En las transfecciones simples, las dos variantes de HSF2BP (WT y S167L) se localizaron en toda la célula de forma homogénea, aunque la intensidad de fluorescencia de la variante HSF2BP-S167L fue menor que la de HSF2BP-WT, indicando una menor expresión/estabilidad de la misma (Figura 39a). La localización de EGFP-BRME1 fue claramente nuclear (Figura 39a). Cuando ambas variantes se co-transfectaron con BRME1 la intensidad de ambas aumentó y cambió su patrón drásticamente pasando a marcar de forma intensa el nucleoplasma formando invaginaciones nucleares similares a las que forma el retículo nucleoplásmico (Malhas et al. 2011). Este patrón se observó en un menor número de células cuando BRME1 se co-transfectó con la variante HSF2BP-S167L, probablemente debido a la menor presencia de esta variante (Figura 39a). A continuación, la menor expresión/estabilidad fue validada mediante WB de extractos procedentes de HEK293T transfectadas con ambas variantes (Figura 39b) observándose además que ambas variantes se acumulaban en presencia de BRME1 (Figura 39b). Esta acumulación también se detectó, aunque en menor medida, cuando las células se trataron con el inhibidor del proteasoma MG132 (Figura 39b), indicando una degradación de HSF2BP parcialmente dependiente del proteasoma que es rescatada por la presencia de BRME1. El hecho de que la acumulación de HSF2BP, tanto en presencia de BRME1 como del inhibidor del proteasoma, sea mayor en el caso la variante S167L probablemente es debido a la menor estabilidad basal de esta variante que se traduce en una mayor degradación vía proteasoma cuando BRME1 no está presente para estabilizarla. En conjunto, tanto los resultados *in vivo* como *in vitro* sugieren la existencia de una fuerte interdependencia funcional entre BRME1 y HSF2BP que da lugar a una menor expresión/estabilidad de ambas en los meiocitos de los ratones mutantes (incluyendo los portadores de la variante causante de POI) que, en última instancia, produce fuertes defectos en la recombinación meiótica.

Figura 39. BRME1 estabiliza a HSF2BP (a) Células U2OS transfectadas con plásmidos que codifican para Flag-HSF2BP (WT o S167L) y EGFP-BRME1 individual o conjuntamente y detectados con Abs frente a Flag o EGFP. Se observa un cambio drástico de ambas proteínas cuando son co-transfectadas, mostrando un patrón de invaginaciones nucleoplásmicas. Este fenotipo es menos severo en presencia de la variante S167L. El gráfico bajo el panel representa la cuantificación del número de células con patrón de retículo nucleoplásmico. $n > 400$ células. Barras de escala, 20 μ m. **(b)** Células HEK293T transfectadas con Flag-HSF2BP (WT o S167L) y EGFP-BRME1 individual o conjuntamente. Además, las células transfectadas se trataron con el inhibidor del proteasoma MG132 (10 μ M). Los extractos se analizaron por WB con un Ab frente a Flag y como control de eficiencia de transfección se co-transfectó un plásmido que codificaba para Cherry. HSF2BP-WT se expresa en mayores niveles que HSF2BP-S167L y la detección se vio incrementada en presencia de BRME1 aunque este incremento fue mayor en la variante S167L. La incubación con MG132 produjo un efecto similar al de BRME1 aunque menos acusado. Los gráficos bajo el panel muestran la cuantificación de la intensidad de las bandas del WB. $n = 3$ transfecciones independientes para cada condición. Análisis Welch t -test de dos colas: ** $p < 0,01$; *** $p < 0,001$; **** $p < 0,0001$.

2.17. BRME1 y HSF2BP forman un complejo multimérico con PALB2 y BRCA2

Por último, con el fin de profundizar en el mecanismo molecular subyacente a los defectos en recombinación observados en los ratones *Brme1*^{-/-}, *Hsf2bp*^{-/-} y *Hsf2bp*^{S167L/S167L} abordamos el estudio del interactoma de BRME1 mediante su inmunoprecipitación en extractos de testículo de ratón acoplado a espectrometría de masas. Como cabía esperar identificamos a HSF2BP como su principal interactor pero también se identificaron BRCA, su interactor clásico PALB2 (Xia et al. 2006), RAD51 y RPA, sugiriendo que todos ellos forman un gran complejo multimérico (Tabla 4).

Genes	ID Proteína	Ab1-Nº de péptidos únicos	Ab2- Nº de péptidos únicos	Control- Nº de péptidos únicos	iBAQ Ab1	iBAQ Ab2	iBAQ Control	Grupo
<i>Hsf2bp</i>	Q9D4G2	20	16	0	168840000	37607000	0	Ab1 y Ab2
<i>C19orf57 (Brme1)</i>	Q6DIA7	16	13	1	77473000	22181000	59511	Ab1 y Ab2
<i>Brca2</i>	P97929	78	77	0	5296400	3125000	0	Ab1 y Ab2
<i>Palb2</i>	Q3U0P1	7	11	0	1102300	896460	0	Ab1 y Ab2
<i>Rad51</i>	Q08297	2	4	0	509370	367390	0	Ab1 y Ab2
<i>Rpa2</i>	Q62193	3	0	0	235990	0	0	Ab1 solo

Tabla 4. Proteínas implicadas en recombinación meiótica identificadas como posibles interactores de BRME1. Estos resultados se obtuvieron mediante IP de BRME1 desde extractos de testículo de ratón acoplada a espectrometría de masas. Se llevaron a cabo dos experimentos en paralelo con dos Abs policlonales de conejo generados frente a la proteína BRME1 completa (Ab1 y Ab2). Como control negativo se usó una IP con IgGs de conejo. El iBAQ representa la intensidad detectada para cada proteína.

Para validar estas interacciones, co-transfectamos los plásmidos que codificaban para cada una de estas proteínas marcadas con diferentes epítomos en células HEK293T y realizamos ensayos de co-IP. BRME1 por sí solo no mostró interacción con ninguna de las regiones de BRCA2 (Nterminal, Medio o Cterminal) (Figura 40a). Sin embargo, en presencia de HSF2BP, BRME1 y BRCA2-C co-inmunoprecipitaron (en ambos sentidos, Figura 40b). Teniendo en cuenta la interacción directa de HSF2BP con BRCA2-C (Figura 23a) y BRME1 (Figura 24a-b), estos resultados indican que las tres proteínas forman un complejo en el que HSF2BP actuaría de puente entre BRCA2 y BRME1.

La validación de las interacciones entre BRME1 y HSF2BP con RPA, RAD51 y PALB2 mostró co-inmunoprecipitación positiva, aunque leve, de HSF2BP con todos ellos y de BRME1 con RAD51 y RPA pero no con PALB2 (Figura 40c). Muchas de las proteínas de la maquinaria implicada en la recombinación meiótica participan también en la recombinación homóloga que se produce en células somáticas, como es el caso de BRCA2, RPA o RAD51. Es por ello, que en este contexto celular (células HEK293T) también se encuentran presentes otras proteínas de la maquinaria de reparación que podrían favorecer la detección de interacciones indirectas. Por este motivo, decidimos validar las interacciones detectadas con RPA, RAD51 y PALB2 expresando cada una de ellas en un sistema libre de células (Sistema TnT) evitando así la presencia de proteínas accesorias. Mediante esta aproximación solo detectamos interacción directa entre BRME1 y HSF2BP (Figura 40d), siendo negativas el resto de las interacciones (Figura 40d) y sugiriendo que estas proteínas no interactúan directamente con HSF2BP y BRME1 sino que forman un complejo (o complejos) *in vivo*, probablemente mediado por BRCA2. La formación de estos complejos estaría afectada tanto en ausencia de BRME1 como de HSF2BP lo que desencadenaría los defectos en recombinación observados en los mutantes nulos. Además, la variante causante de POI (HSF2BP-S167L), dada su menor expresión/estabilidad podría estar alterando, aunque de forma más sutil, la formación de estos grandes complejos de recombinación que incluyen a BRCA2, dando lugar al fenotipo menos severo de estos mutantes.

Figura 40. Co-inmunoprecipitación de BRME1 y HSF2BP con los posibles interactores identificados por espectrometría de masas. (a-c) Los plásmidos codificantes para BRME1, HSF2BP, BRCA2 (C, M o N), RPA1, RAD51 o PALB2 marcados con diferentes epítomos se co-transfectaron en células HEK293T. Los complejos proteicos se inmunoprecipitaron con los Abs indicados en verde y se analizaron por WB con los Abs indicados en rojo. Las IgGs se usaron como control negativo. **(a)** BRME1 y BRCA2-C co-inmunoprecipitan en presencia de HSF2BP. La co-inmunoprecipitación de HSF2BP con BRCA2-C se muestran en la Figura 23a. **(b)** BRME1 no co-inmunoprecipita con BRCA2-N, BRCA2-M o BRCA2-C cuando están los dos solos. **(c)** BRME1 (izquierda) co-inmunoprecipita ligeramente con RPA1 y RAD51 pero no con PALB2. HSF2BP (derecha) co-inmunoprecipita ligeramente con RPA1, RAD51 y PALB2. **(d)** BRME1, HSF2BP y sus posibles interactores (RPA1, RAD51 y PALB2) se expresaron marcados con un epítomo (Flag o HA) en un sistema libre de células TnT. Estas proteínas se incubaron juntas 30 min a RT para permitir formar los complejos y se inmunoprecipitaron con los Abs indicados en verde. El análisis por WB se hizo con los Abs indicados en rojo. La única interacción directa detectada fue entre BRME1 y HSF2BP.

DISCUSIÓN

1. USP26 no es esencial para la fertilidad en ratón

1.1. La ausencia de USP26 no afecta a la meiosis ni la fertilidad en ratón

En el presente estudio hemos generado dos modelos de pérdida de función de *Usp26*, uno de ellos portador de dos inserciones consecutivas (alelo²¹) y otro portador de una delección que elimina la cisteína catalítica de USP26 (mutante inactivo, alelo^{Cat}). Aunque el análisis *in silico* predice que ambos alelos generan proteínas truncadas, no hemos podido demostrar la ausencia de USP26 en nuestros mutantes debido a la falta de sensibilidad de los Abs utilizados, incapaces de detectar la proteína USP26 endógena. Pese a esto, la delección generada en el mutante catalítico (Cat) asegura que, de expresarse parte de la proteína, esta no sería funcional como proteasa específica de ubiquitina dando lugar a un mutante de pérdida de función.

USP26 pertenece a la superfamilia de las enzimas desubiquitinantes encargadas del mantenimiento de la homeostasis de la ubiquitina en las células mediante el procesamiento de sus péptidos precursores y la eliminación de la ubiquitina de sus sustratos (Komander et al. 2009). La ubiquitinación desempeña funciones fundamentales a lo largo de la espermatogénesis, desde el establecimiento y la diferenciación de las SSCs hasta la espermiogénesis (Bose et al. 2014). En este sentido, diferentes enzimas desubiquitinantes se han asociado con defectos en diferentes estadios de la espermatogénesis, así como con defectos en fertilidad (entre ellas USP2 (Bedard et al. 2011), USP7 (Luo et al. 2015), USP9X (Kishi et al. 2017) y Uchl-1 (Kwon et al. 2005)). Sin embargo, nuestros datos muestran que los ratones deficientes en USP26 son fértiles, tienen camadas del mismo tamaño que los silvestres y presentan un número y distribución normal de las células del testículo.

Usp26 alcanza su máximo nivel de transcripción durante los primeros estadios de la profase meiótica (ver anexo, Tablas S1 y S2) (Margolin et al. 2014; da Cruz et al. 2016). Este patrón de expresión, junto con la importancia del sistema ubiquitin-proteasoma en la regulación de la profase en ratón (Rao et al. 2017) y la implicación de USP26 en la reparación del DNA y la recombinación homóloga en células somáticas (Typas et al. 2016), sugieren un posible papel de USP26 en estos procesos. Sin embargo, el análisis citológico de los espermatocitos mutantes no mostró defectos aparentes en sinapsis, reparación de las DSBs y recombinación meiótica sugiriendo que la función de USP26 no estaría relacionada con la regulación de la profase I meiótica.

1.2. Asociación entre USP26 e infertilidad humana

La asociación de esta proteína específica de testículo con la infertilidad masculina ha sido foco de muchos estudios durante las últimas dos décadas, aunque con resultados controvertidos. Diversos trabajos han relacionado mutaciones en *USP26* con infertilidad masculina en humanos (Paduch et al. 2005; Stouffs et al. 2005; Zhang et al. 2007; Lee et al. 2008a; Asadpor et al. 2013; Xia et al. 2014; Li et al. 2015; Ma et al. 2016; Li et al. 2020a; Arafat et al. 2020). Sin embargo, nuestros estudios funcionales indican que USP26 es prescindible para la fertilidad del ratón y concuerdan con los trabajos que muestran una ausencia de asociación entre los polimorfismos genéticos en *USP26* y la infertilidad masculina (Ravel et al. 2006; Stouffs et al. 2006; Christensen et al. 2008; Ribarski et al. 2009; Shi et al. 2011; Zhang et al. 2015; Luddi et al. 2016). Además, nuestras observaciones están respaldadas por la identificación de un paciente normo-zoospérmico portador de una mutación en *USP26* (c.882 C>A) que genera un codón de STOP prematuro y da lugar a una

proteína truncada en la posición 293 y que, por tanto, carece de la cisteína catalítica humana (Cys305) (Luddi et al. 2016).

Nuestros resultados aportan nuevas evidencias que podrían arrojar luz sobre la gran controversia existente en relación con *USP26* y la infertilidad masculina en humanos. Sin embargo, aunque el ratón es el modelo más utilizado en la investigación biomédica (Rosenthal and Brown 2007), no podemos descartar que la relevancia funcional de *USP26* difiera entre ambas especies, teniendo un papel crítico en humanos pero no en ratón. Por otra parte, en paralelo con nuestro estudio Tian et al. (2019) describieron la presencia de defectos en la espermatogénesis y subfertilidad en un modelo nulo para *Usp26* tras más de 6 generaciones de retrocruces en un fondo genético C57BL/6J. En contraposición con este estudio, otro trabajo posterior no detectó defectos en la espermatogénesis ni en la fertilidad de mutantes nulos para *Usp26* retrocruzados 5 veces en ese mismo fondo, pero sí cuando se retrocruzaron de 3 a 5 generaciones en un fondo genético DBA/2 (Sakai et al. 2019). Los ratones retrocruzados en fondo DBA/2 mostraron defectos en sinapsis y una reducción en el número de espermatozoides que condujeron a fenotipos de subfertilidad o, incluso, en algunos casos, infertilidad. En conjunto, estos estudios ponen de manifiesto la influencia del fondo genético en los análisis funcionales realizados en mutantes murinos, así como la importancia de considerarlo a la hora de extrapolar las conclusiones a los humanos. Los dos modelos desarrollados en este trabajo se encuentran en un fondo mixto no consanguíneo B6/CBA F2 (híbridos entre las cepas C57BL/6J y CBA/J) que refleja mejor la variabilidad genética presente en las poblaciones humanas.

Por último, dada la expresión restringida a testículo de *Usp26* y la importancia de la regulación del balance ubiquitinación/desubiquitinación en la célula, la ausencia total de fenotipo de nuestros mutantes resulta sorprendente y podría explicarse por una redundancia genética en la función de *Usp26*. El análisis *in silico* del perfil de transcripción de diferentes USPs durante la espermatogénesis muestra que varias de ellas presentan un patrón altamente similar al de *USP26* (*USP9X*, *USP37*, *USP14*, *USP53*, *USP33*, *USP28* y *USP3*; ver anexo Tablas S1 y S2 (Margolin et al. 2014; da Cruz et al. 2016)) postulándose como posibles candidatas a compensar la pérdida de función de *USP26*. Sería interesante analizar si alguna de estas USPs se encuentra sobreexpresada en ausencia de *USP26*, lo que apoyaría la existencia de algún tipo de redundancia funcional. De identificarse alguna, para obtener la demostración definitiva sería necesaria la generación del modelo de pérdida de función de ambas proteínas simultáneamente.

En conclusión, este trabajo supone el primer análisis funcional de *USP26* en ratón, así como la primera evidencia *in vivo* de que *USP26* no es esencial para la gametogénesis y la fertilidad del ratón, aportando nuevos datos a la gran controversia existente en torno a esta ubiquitin-proteasa.

2. Una variante en HSF2BP causa POI a través de su nuevo interactor BRME1

2.1. La variante S167L en *HSF2BP* afecta a la fertilidad en mujeres

Mediante la secuenciación del exoma de tres casos de POI hereditarios procedentes de una familia endogámica de origen árabe-israelí, se ha identificado la variante S167L en *HSF2BP* como la mejor candidata causal de la infertilidad. En esta familia todos los miembros afectados son homocigotos para la mutación mientras que los miembros sanos son portadores heterocigotos. Esto, junto con la identificación de HSF2BP como una nueva proteína esencial para la recombinación meiótica y la fertilidad en ratón, apoyan la causalidad de la variante HSF2BP-S167L en la POI. Dicha causalidad fue corroborada por el fenotipo meiótico y la subfertilidad observada en las hembras *Hsf2bp*^{S167L/S167L}. Además, los defectos observados en los meiocitos *Hsf2bp*^{S167L/S167L}, como son la reducción del número de foci de RAD51/DMC1 en las DSBs y la consiguiente reducción del número de COs, ponen de manifiesto que esta variante provoca alteraciones en la recombinación meiótica. El análisis comparativo del alelo S167L con el alelo nulo para *Hsf2bp* mostró que esta variante de cambio de sentido se comporta como un alelo hipomorfo, lo cual está de acuerdo con la amenorrea secundaria observada en las pacientes y la fertilidad residual (con asistencia médica) presente en una de las hermanas afectadas. La identificación de *HSF2BP* como un nuevo gen causante de POI debido a alteraciones en la recombinación meiótica concuerda con la creciente aunque reducida lista de genes de esta ruta responsables de POI que incluye a *BRCA2*, *MEIOB*, *SYCE1*, *MSH4* o *DMC1* (De Vries et al. 2014; Carlosama et al. 2017; He et al. 2018; Caburet et al. 2019; Caburet et al. 2020).

Aunque no podemos descartar la aparición con el tiempo de un fenotipo de POI en estos ratones, la subfertilidad de las hembras *Hsf2bp*^{S167L/S167L} representa un fenotipo menos severo que el observado en las pacientes. Esto podría deberse a una menor sensibilidad del ratón con respecto al humano ante los alelos hipomorfos, como sucede con la menor sensibilidad del ratón a la dosis génica en aquellos genes que son haploinsuficientes en humanos (Veitia 2003). Por otro lado, los eventos iniciales de la meiosis femenina en humanos parecen ser más propensos a la aparición de errores que en ratón o que en hombres, como se evidencia por la mayor incidencia de defectos en sinapsis presente en los oocitos humanos. Además, el hecho de que los foci de MLH1 se detecten mucho antes durante la profase I femenina humana en comparación con la masculina o la de ratón, sugiere una diferente regulación temporal de la recombinación en mujeres (Hassold et al. 2007). Por último, recientemente se ha descrito que los oocitos humanos presentan cierta ineficiencia en la maduración de los CO de tal forma que, pese a formarse un mayor número de COs en mujeres, la frecuencia de bivalentes sin CO es más elevada en estas que en los hombres (Wang et al. 2017b). Todas estas observaciones podrían explicar el fenotipo más severo de las pacientes con POI en comparación con las hembras del modelo *Hsf2bp*^{S167L/S167L}.

En el caso de los hombres, la ausencia en la familia de hermanos varones portadores de la mutación en homocigosis impide la comparación directa entre el fenotipo observado en los ratones macho *Hsf2bp*^{S167L/S167L} y los humanos. Sin embargo, la extrapolación a los humanos de nuestros resultados en ratón sugiere que la presencia de la variante S167L en homocigosis provocaría defectos de fertilidad en los varones. No es descartable que estudios futuros identifiquen esta u otras variantes en *HSF2BP* asociadas a

infertilidades masculinas. En este sentido, recientemente se ha descrito una variante en *HSF2BP* (G224*) que afecta a la recombinación meiótica en hombres y podría estar asociada con problemas de fertilidad. Los dos hermanos portadores de esta variante (hombre y mujer) no presentaron problemas de salud pero ninguno de ellos tuvo descendencia, lo que podría indicar que ambos eran infértiles (Halldorsson et al. 2019). Esta observación refuerza la hipótesis de que diferentes variantes en *HSF2BP* podrían afectar también la fertilidad masculina en humanos.

2.2. Los mutantes de *Hsf2bp* y *Brme1* presentan dimorfismo sexual

Los mutantes meióticos en ratón a menudo presentan fenotipos sexualmente dimórficos (Morelli and Cohen 2005; Cahoon and Libuda 2019). Esta disimilitud puede deberse a las diferencias estructurales en el SC descritas entre sexos como son la distinta longitud de los ejes cromosómicos (Gruhn et al. 2013) o la diferente función de ciertas proteínas de los ejes entre machos y hembras, como ocurre en el caso de *RAD21L* (Herrán et al. 2011). En los mutantes de recombinación meiótica, las hembras habitualmente son capaces de avanzar más en la meiosis que los machos. De acuerdo con esto, los mutantes nulos de *Hsf2bp* y *Brme1* mostraron un bloqueo en zigotena-like en machos y, sin embargo, las hembras fueron capaces de generar un reducido número de oocitos fertilizables pese a los defectos observados en sinapsis. Este dimorfismo puede deberse, además de a las diferencias estructurales del SC, a la ausencia de silenciamiento del cromosoma Y cuando el bivalente sexual no sinapsa en los machos mutantes y/o a la presencia de *checkpoints* menos severos durante la oogenesis femenina (Hunt and Hassold 2002).

Los ratones *Hsf2bp*^{S167L/S167L} también mostraron cierto dimorfismo sexual tanto molecular como reproductivo. Aunque ambos sexos comparten gran parte de las alteraciones moleculares en la ruta de recombinación meiótica, destacando la reducción del número de foci de *RAD51/DMC1* y la consiguiente reducción en el número de COs, también se observaron ciertas diferencias. En las hembras la reducción de *DMC1* fue más acusada y no presentaron acumulación de RPA pese a la acumulación de daño en el DNA, algo que también se observó en las hembras de los modelos nulos. Estas diferencias apuntan hacia la existencia de diferentes mecanismos subyacentes en la recombinación meiótica de cada sexo, de manera similar a lo que ocurre con *TEX15*, proteína esencial para la carga de *RAD51/DMC1* en machos pero no en hembras (Yang et al. 2008). La existencia de estos mecanismos moleculares diferenciales explicaría que la recombinación meiótica sea el proceso donde más dimorfismo sexual se ha descrito hasta el momento.

Desde el punto de vista reproductivo también encontramos diferencias entre sexos en los mutantes *Hsf2bp*^{S167L/S167L}. Ambos muestran una disminución significativa del número de COs y, consecuentemente, una mayor frecuencia de meiocitos con bivalentes sin COs. Sin embargo, mientras que en machos esto da lugar a una disminución del número de espermatozoides en el epidídimo, en hembras el número de oocitos de la reserva folicular no se ve afectado. Paradójicamente, los machos no presentaron una reducción de la fertilidad mientras que las hembras fueron subfértiles. El hecho de que no hayamos detectado ningún impacto estadísticamente significativo de la variante S167L en la fertilidad masculina podría deberse a la alta variabilidad obtenida tanto en la frecuencia de las camadas como en el tamaño de las mismas, lo que provocaría una disminución de la potencia estadística del análisis. Por otra parte, se sabe que una fuerte reducción en la cantidad de espermatozoides (de hasta el 60%) no afecta a la fertilidad del ratón lo que

explicaría la fertilidad normal de los machos *Hsf2bp*^{S167L/S167L} a pesar de los evidentes defectos meióticos observados (Schürmann et al. 2002). En el caso de las hembras la leve subfertilidad observada (reducción en la frecuencia de las camadas) podría deberse al menor número de gametos disponibles para ser fertilizados con respecto a los machos, pero también a las diferencias moleculares en la recombinación meiótica de ambos sexos que hemos comentado anteriormente (Cahoon and Libuda 2019). Dado que en hembras la metafase I se completa tras la ovulación, proceso posterior al establecimiento de la reserva de oocitos del ovario (Bolcun-Filas and Handel 2018), la reducción del número de COs observado en las hembras *Hsf2bp*^{S167L/S167L} podría producir defectos en la metafase I dando lugar a oocitos que, tras la fertilización, producirían embriones no viables. Esto explicaría la reducción de la fertilidad pese a no existir diferencias en la reserva folicular. En el caso de las hembras *Hsf2bp*^{-/-} y *Brme1*^{-/-}, los defectos en recombinación son tan severos que los oocitos presentan fuertes alteraciones de la sinapsis que conllevan su eliminación tras la profase I y la consiguiente reducción de la reserva folicular.

2.3. HSF2BP y BRME1 forman un complejo funcional esencial para la recombinación

Mediante diferentes aproximaciones complementarias tanto *in vivo* e como *in vitro* hemos demostrado que las proteínas HSF2BP y BRME1 constituyen un complejo funcional en el que ambas subunidades son esenciales para la recombinación meiótica y para su expresión y/o estabilidad mutua. De acuerdo con esto, la eliminación genética de HSF2BP o BRME1 da lugar a fenotipos muy similares, si no idénticos. La oogénesis de ambos mutantes nulos presenta defectos severos en sinapsis que producen la eliminación de la mayoría de los oocitos antes de la formación de los folículos primordiales. Por su parte, la espermatogénesis se encuentra bloqueada en un estadio zigotena-like dando lugar a la ausencia de espermatozoides. Mecánicamente, estos defectos son debidos a la disminución drástica de las recombinasas RAD51/DMC1 que impide la correcta reparación de las DSBs y produce una ausencia total o una reducción significativa de los COs en machos y hembras, respectivamente. Como consecuencia de estos defectos en reparación los espermatoцитos bloqueados en zigotena-like acumulan las proteínas de unión a ssDNA, RPA y SPATA22, mientras que los oocitos únicamente acumulan SPATA22. El fenotipo de estos mutantes, junto con la alta co-localización de HSF2BP/BRME1 con RPA, sugiere que estas proteínas actúan en los eventos tempranos de la recombinación meiótica tras la formación de las DSBs. De acuerdo con esto, la ausencia de BRME1 y HSF2BP en el mutante nulo de *Spo11* indica que su localización es dependiente de la formación de las DSBs. Además, el hecho de que la localización de BRME1 y HSF2BP no se viera alterada en los mutantes de pérdida de función de *Rnf212* y *Hei10*, con defectos en la designación y maduración de los COs (Qiao et al. 2014), ni en los de *Rec8* y *Six6os1*, en los que no se produce la invasión del homólogo debido a la asinapsis (Bannister et al. 2004; Gomez-H et al. 2016), nos permite determinar que la función de HSF2BP/BRME1 es independiente del establecimiento de la sinapsis y anterior a la invasión del homólogo mediada por RAD51/DMC1.

Durante la caracterización de la función de HSF2BP y BRME1 en meiosis y de cómo la variante S167L modifica la fertilidad en ratones, se han descrito dos modelos knockout para *Hsf2bp* (Brandsma et al. 2019; Zhang et al. 2019) y cuatro modelos knockout para *Brme1* (Takemoto et al. 2020; Shang et al. 2020; Li et al. 2020b; Zhang et al. 2020b) lo que pone de manifiesto el interés creciente en el desciframiento de los mecanismos moleculares que gobiernan la meiosis y, más concretamente, la recombinación meiótica. De

forma similar a lo descrito en esta tesis, estos grupos reportaron de manera independiente que HSF2BP y BRME1 son esenciales para la recombinación meiótica y la fertilidad del ratón. En el caso de los mutantes nulos para *Hsf2bp*, el fenotipo de los machos fue igual al observado en nuestros mutantes. Sin embargo, la ausencia de HSF2BP en hembras dio lugar a fenotipos con diferente severidad, siendo en unos casos subfértiles ((Zhang et al. 2019) y esta tesis) y en otros fértiles (Brandsma et al. 2019). En cuanto a los mutantes de pérdida de función de *Brme1*, en todos los casos los machos mostraron problemas de fertilidad y alteraciones moleculares similares aunque con diferente severidad. Nuestro modelo *Brme1*^{-/-} mostró un bloqueo casi total en estadio zigoteno-like y una ausencia de COs. Sin embargo, los mutantes de Zhang et al., Shang et al. y Li et al. presentaron un alto porcentaje de células capaces de alcanzar el estadio de paquitena y formar COs e incluso una pequeña fracción alcanzó diplotena y metafase I. Sorprendentemente, el mutante de Takemoto et al. presentó una cantidad residual de espermatidas y espermatozoides que condujeron a una subfertilidad muy severa pero no a la infertilidad. En cuanto a las hembras, solo tres de los trabajos abordaron superficialmente su análisis mostrando una fertilidad normal, en contraposición con la fuerte reducción de la reserva folicular y los defectos meióticos de nuestras hembras *Brme1*^{-/-} (Takemoto et al. 2020; Shang et al. 2020; Li et al. 2020b). Estas diferencias no parecen deberse a la naturaleza de la mutación de cada modelo dado que el mutante con el fenotipo menos severo es el que tiene la delección más grande (el mutante de Takemoto et al. carece de los últimos 6 exones, delección que abarca la región delecionada en los otros 4 mutantes). Por lo tanto, la mayor severidad del fenotipo de nuestros mutantes *Brme1*^{-/-}, tanto en machos como en hembras, podría explicarse por los distintos fondos genéticos de estos ratones. De igual manera, las diferencias en el fondo genético de los mutantes de *Hsf2bp* descritos podría explicar la severidad variable del fenotipo observado en las hembras.

En relación con el análisis de la función de BRME1 durante la meiosis, además de los defectos en recombinación meiótica ya mencionados, Li et al. (2020b) describen una fuerte alteración de la ubiquitinación tanto en los autosomas como en el *sex body* en ausencia de BRME1. Sus resultados descartan que esta proteína actúe como E3 ligasa o que sea en sí mismo un sustrato de ubiquitinación y proponen que BRME1 podría participar en la progresión de la profase I meiótica regulando la ubiquitinación de proteínas meióticas clave (Li et al. 2020b). Nuestros resultados (*in vivo* e *in vitro*) sugieren una estabilización de HSF2BP dependiente de BRME1 y del proteasoma apoyando la hipótesis de que BRME1 podría participar en la regulación de la profase I y la recombinación meiótica modulando la ubiquitinación de diferentes sustratos. Dado que nuestro análisis *in vivo* sugiere que HSF2BP es necesaria para la estabilidad de BRME1, y teniendo en cuenta la fuerte interdependencia funcional de ambas proteínas, no podemos descartar que HSF2BP también esté implicada en esta regulación dependiente de ubiquitinación.

Finalmente, la inmunoprecipitación de BRME1 de extractos de testículo acoplada a espectrometría de masas detectó principalmente a su pareja HSF2BP, pero también a las proteínas PALB2, RAD51, RPA y BRCA2. Sin embargo, mediante ensayos *in vitro* observamos que la única interacción directa se producía entre BRME1 y HSF2BP. Todo ello sugiere que el resto de interacciones podrían estar mediadas por la proteína multidominio BRCA2 ya que HSF2BP interacciona directamente con ella (este trabajo y Brandsma et al. 2019) y con BRME1 (este trabajo). Además, BRCA2 también interacciona directamente con PALB2 (Xia et al. 2006), con las recombinasas RAD51 y DMC1 y con el DNA, a través de su capacidad de unión a este (Siaud et al.

2011). Estos complejos multiproteicos gobernados por BRCA2 participan en la sucesión ordenada de eventos que ocurren en la reparación de las DSBs como son la unión inicial de RPA a las zonas reseccionadas, el intercambio de RPA por RAD51/DMC1 y la carga de MEIOB/SPATA22 a los complejos de RPA (Zhao et al. 2015; Martínez et al. 2016). Recientemente, se han identificado variantes en pacientes con POI en genes relacionados con BRCA2 e implicados en los procesos tempranos de la reparación de las DSBs meióticas como son *MEIOB*, *DMC1* o el propio *BRCA2* (He et al. 2018; Caburet et al. 2019; Caburet et al. 2020). Todo ello pone de manifiesto la elevada susceptibilidad y la importancia de la regulación de este punto de la ruta meiótica, así como el papel central que BRCA2 desempeña en la misma.

2.4. La variante HSF2BP-S167L se comporta como un alelo hipomorfo en ratón

El análisis de la variante en HSF2BP causante de POI (S167L), tanto *in vivo* como *in vitro*, ha mostrado que se esta comporta como un alelo hipomorfo en ratón dando lugar a una reducción de su propia expresión/estabilidad y la de su interactor BRME1. Como consecuencia de ello, los meiocitos *Hsf2bp*^{S167L/S167L} presentaron cualitativamente las mismas alteraciones moleculares en la ruta de la recombinación meiótica que los meiocitos *Hsf2bp*^{-/-} y *Brme1*^{-/-} pero cuantitativamente menos severas. Estas incluyeron una reducción de RAD51/DMC1 y de los COs, así como una acumulación de RPA (solo en machos) y de SPATA22. Estas alteraciones condujeron a una disminución de la cantidad de espermatozoides sin afectar a la fertilidad de los machos *Hsf2bp*^{S167L/S167L} y a una ligera reducción en la fertilidad en las hembras. De acuerdo con la pérdida parcial de función de la variante S167L, dichos fenotipos fueron menos severos que los observados en los mutantes nulos.

La acumulación de RPA observada en los machos *Hsf2bp*^{S167L/S167L} probablemente se corresponde con la primera oleada de actividad de RPA que tiene lugar durante los primeros estadios de la recombinación. El complejo RPA (RPA1-RPA2-RPA3) se une y estabiliza los intermediarios de ssDNA que se forman durante la reparación del DNA y en meiosis forma un complejo con otras dos proteínas esenciales, MEIOB y SPATA22. (Luo et al. 2013). Se ha postulado que RPA actúa en dos momentos diferentes durante la meiosis. En primer lugar, durante las fases iniciales de la recombinación uniéndose al ssDNA generado tras la resección de las DSBs donde es reemplazado rápidamente por las recombinasas RAD51/DMC1. En segundo lugar, durante la invasión del cromosoma homólogo mediada por RAD51/DMC1 protegiendo las regiones de ssDNA del *D-loop* que se forman tras el desplazamiento de la hebra (Shi et al. 2019). La demostración de que HSF2BP y BRME1 no se unen directamente al DNA sugiere un modelo en el que el complejo HSF2BP/BRME1 se localiza en las DSBs probablemente a través de la interacción con BRCA2 siendo indispensable para la posterior localización de RAD51/DMC1. La ausencia del complejo HSF2BP/BRME1 impide el reemplazamiento de RPA por RAD51/DMC1 en los RNs lo que produciría la acumulación observada de RPA y de su interactor SPATA22. De igual manera, la reducción de la cantidad de HSF2BP/BRME1 como consecuencia de la variante S167L provocaría un reemplazo menos eficiente de RPA por RAD51/DMC1 dando lugar a la acumulación, aunque en menor grado, de RPA/SPATA22 y provocando una disminución del número de COs. En el caso de las hembras no se observó acumulación de RPA en ninguno de los tres modelos mutantes analizados, pero sí de SPATA22. Dado que se desconoce la función *in vivo* de RPA en la ovogénesis y que la carga de MEIOB/SPATA22 a las DSBs es independiente de RPA (Shi et al. 2019), podríamos especular que en la meiosis femenina el

papel de RPA en la unión del ssDNA y su reemplazo por RAD51/DMC1 en los estadios tempranos de la recombinación podría ser llevado a cabo por otro complejo como SPATA22/MEIOB de manera dependiente de HSF2BP/BRME1.

Recientemente, el desarrollo de un mapa de recombinación meiótica de alta resolución ha permitido identificar nuevos loci implicados en el control de este proceso a través del análisis de variantes que modifican la tasa de recombinación y/o la localización de los COs (Halldorsson et al. 2019). Entre ellos están genes relacionados con la formación del SC (*Syce2*, *Rad21l*, *Sycp3* y *Six6os1*) y proteínas de la propia maquinaria de recombinación meiótica. Dentro de esta segunda categoría se identificaron variantes en las E3 ligasas RNF212 (SUMO-ligasa) y HEI10 (Ubiquitin-ligasa), proteínas ampliamente documentadas como determinantes de la tasa de recombinación (Reynolds et al. 2013; Lake and Hawley 2013), pero también en proteínas de los eventos tempranos de recombinación como *Meiob* y, lo que es más importante, en *Hsf2bp* (Halldorsson et al. 2019). Es importante destacar que algunos de estos genes descritos como modificadores de la recombinación meiótica han sido descritos como “genes de fertilidad” en humanos, como es el caso de *SYCP3*, *HFM1* y *HSF2BP* (Wang et al. 2014; Geisinger and Benavente 2017 y este trabajo). Con estos antecedentes, planteamos la hipótesis de que diferentes variantes en el mismo gen pueden alterar la tasa de recombinación en diferente grado pudiendo dar lugar desde fenotipos leves (sin efectos perjudiciales) a fenotipos severos (causantes de infertilidad) cuando esta tasa de recombinación se reduce por debajo del límite de un CO por bivalente. En el caso bajo estudio, los resultados sugieren que la variante S167L provoca una hipoactividad del complejo HSF2BP/BRME1 que da lugar a una reducción de la tasa de recombinación por debajo de ese límite. Hasta donde sabemos, *HSF2BP* es uno de los escasos genes humanos con variantes que afectan tanto a la recombinación meiótica en la población (Halldorsson et al. 2019) como a la fertilidad de manera directa a través de defectos en la segregación cromosómica causados por la reducción de la tasa de recombinación (este trabajo). En línea con esta hipótesis, es factible que la presencia de variantes con efectos aditivos pueda conducir a una mayor reducción de la tasa de recombinación y, por lo tanto, a la aparición de aneuploidías e infertilidad (Schimenti and Handel 2018). Concretamente, variantes aditivas en proteínas implicadas en recombinación meiótica o componentes del SC podrían estar detrás de una parte importante de las infertilidades idiopáticas. En principio, estas variantes deberían estar sometidas a selección negativa y reducirse en gran medida o incluso eliminarse de la población. Sin embargo, esto no ocurriría en aquellos genes con dimorfismo sexual donde los individuos de uno de los sexos son portadores fértiles de dichas variantes, algo que sucede en un gran número de genes meióticos incluidos *Hsf2bp* y *Brme1* (Gershoni and Pietrokovski 2014; Cahoon and Libuda 2019).

En resumen, en este trabajo hemos descrito por primera vez una familia humana donde la POI co-segrega mendelianamente con la variante S167L en *HSF2BP*. Mediante el análisis de los ratones humanizados hemos demostrado que la variante S167L se comporta como un alelo hipomorfo que provoca una menor expresión y/o estabilidad del complejo HSF2BP/BRME1 y fenocopia, aunque de manera menos severa, los defectos meióticos observados en los ratones que carecen de HSF2BP o de su nuevo interacto BRME1.

3. Relevancia del estudio

La infertilidad humana es un problema de salud pública que lejos de atajarse se ve acrecentado por los cambios socioculturales de la población, entre los que destaca el aumento de la edad en la que se decide tener el primer hijo (Balasch and Gratacós 2012). Este hecho hace que cada vez más parejas necesiten la ayuda de técnicas de reproducción asistida para concebir lo que conlleva un gran esfuerzo psicológico y económico. La presencia de alteraciones genéticas que provoquen defectos en la gametogénesis de los pacientes puede afectar de manera importante e incluso impedir el éxito de estas técnicas. En estos casos, el diagnóstico temprano de las alteraciones genéticas subyacentes es fundamental pudiendo evitarse la aplicación de procedimientos costosos e inefectivos. Por lo tanto, descifrar las bases genéticas de la infertilidad humana ampliando la batería de genes relacionados con esta patología resulta fundamental para su diagnóstico y, posiblemente, para su tratamiento futuro. En este sentido, las clínicas de reproducción asistida incluyen en su práctica clínica paneles de genes de fertilidad basados en publicaciones científicas que, en el mejor de los casos, evidencian una relación causal entre las alteraciones genéticas y los defectos de fertilidad en humanos o en ratón. La utilización de estos paneles facilita la identificación y aplicación de los tratamientos y técnicas de reproducción asistida más apropiados para cada pareja (Harper et al. 2018; Cariati et al. 2019). En relación con esto, en la presente tesis hemos abordado el estudio de dos genes candidatos a producir infertilidad en humanos, *USP26* y *HSF2BP*. En el caso de *USP26*, el debate sobre su relación con la infertilidad humana se ha mantenido desde hace décadas y aunque nuestros datos sugieren que no existe una relación directa entre este gen y la infertilidad, al menos en ratón, siguen existiendo resultados controvertidos siendo necesarios más estudios para confirmar o descartar su causalidad. Por el contrario, en el caso de *HSF2BP*, nuestra investigación demuestra una relación directa entre una mutación en este gen y el desarrollo de la infertilidad previamente identificada en humanos.

Por otro lado, los resultados obtenidos en esta tesis doctoral suponen un paso más en el desciframiento de los mecanismos moleculares existentes en la meiosis del ratón. Dado que la meiosis humana no puede ser analizada funcionalmente de forma directa, el uso del ratón como modelo de estudio representa actualmente la mejor alternativa para profundizar en el conocimiento de estos procesos en humanos. La descripción de *USP26* como proteína prescindible para la espermatogénesis de ratón, al menos en ciertos fondos genéticos, abre la posibilidad de que existan mecanismos redundantes en la regulación de la ubiquitinación durante este proceso y que, además, su efectividad dependa del paisaje genético concreto de cada individuo. Esta hipótesis tiene fuertes implicaciones en la extrapolación de los resultados al humano, ya que, de demostrarse, sería necesario un análisis individualizado de cada paciente, así como la identificación de marcadores adicionales que pudieran predecir el efecto de las mutaciones en *USP26* en cada uno de ellos. Además, es factible suponer que puedan existir múltiples genes sometidos a mecanismos compensatorios, aumentando la complejidad del diagnóstico genético en la práctica reproductiva. En un futuro próximo, el desarrollo de células iPS (*induced Pluripotent Stem*) de cada individuo, junto con la diferenciación *in vitro* a línea germinal y la edición genómica, permitirán el análisis de la interrelación de las mutaciones en genes implicados en reproducción con su fondo genético particular con un detalle y precisión sin precedentes (Ishii 2014; Zhou et al. 2016).

Por último, en este trabajo hemos seguido una ruta inversa a la habitual en la que a partir del estudio de una mutación en un gen anónimo candidato a producir POI hereditaria, hemos identificado y caracterizado un nuevo heterodímero (HSF2BP/BRME1) que a su vez forma parte de un supra-complejo proteico ensamblado entorno al supresor tumoral BRCA2. La ausencia de cualquiera de las subunidades del heterodímero, o su disminución como consecuencia de la variante causante de POI, provoca defectos en la recombinación meiótica y la fertilidad del ratón. Estos resultados aportan nuevos e importantes conocimientos sobre los mecanismos moleculares que gobiernan la recombinación meiótica, evidenciando su complejidad y la necesidad de seguir profundizando en estos procesos ya que, a pesar del conocimiento actual de los elementos que participan en esta ruta, todo apunta a que existen elementos por identificar con función aún desconocida.

CONCLUSIONES

1. USP26 no es esencial para la fertilidad del ratón en un fondo genético mixto C57Bl6/CBA.
2. USP26 no participa en la regulación de la profase I, en la reparación de las DSBs ni en la recombinación meiótica del ratón.
3. HSF2BP es esencial para la fertilidad del ratón y su variante S167L produce una reducción parcial de la misma.
4. HSF2BP es esencial para la localización de las recombinasas RAD51/DMC1 y la formación de los COs en el ratón .
5. La variante S167L reduce la expresión y/o estabilidad de HSF2BP y se comporta funcionalmente como un alelo hipomorfo el cual provoca defectos en la recombinación meiótica menos severos que su alelo nulo.
6. C19ORF57/BRME1 es un nuevo componente de los nódulos de recombinación que interacciona fuertemente con HSF2BP.
7. BRME1 es esencial para la fertilidad del ratón y funcionalmente participa en los mismos procesos de la recombinación meiótica que HSF2BP.
8. El complejo HSF2BP/BRME1 muestra una elevada interdependencia funcional en el que ambas proteínas son necesarias para la localización y/o estabilización de la otra.
9. El heterodímero HSF2BP/BRME1 forma parte de un complejo macromolecular superior dependiente de BRCA2.
10. La variante S167L de HSF2BP produce una reducción del complejo HSF2BP/BRME1 *in vivo* que es la responsable de los defectos en recombinación meiótica y la subfertilidad observados en los ratones que la portan.
11. La variante S167L de HSF2BP es la responsable de la POI en el pedigrí familiar produciendo defectos en la recombinación meiótica mediados por la alteración del nuevo complejo HSF2BP/BRME1.

ANEXO

1. Material suplementario

1.1 USP26 no es esencial para la fertilidad en ratón

Tabla S1. Expresión de diferentes USPs (mRNA) descrita por da Cruz et al. (2016). Unidades: RPKM (*reads per Kilobase per million mapped reads*).

Min Max

	Fracción 2C (Espermatogonia y células somáticas)	Fracción LZ (Lepto-Zigotena)	Fracción PS (Paquitena)	Fracción RS (Espermátidas redondas)
Usp54	10,54	10,66	9,72	10,23
Usp33	15,75	18,81	8,31	1,42
Usp10	4,38	4,83	3,54	0,13
Usp14	10,71	14,47	4,85	3,5
Usp9x	30,43	56,25	5,35	10,5
Usp5	8,78	13,9	4,69	1,68
Usp24	20,14	26,96	15,11	7,67
Usp40	10,48	18,98	6,08	1,36
Usp28	6,97	12,99	3,5	1,98
Usp3	16,2	37,89	6,6	4,3
Usp53	7,44	12,29	5,23	3,27
Usp26	5,88	62,51	2,32	3,36
Usp34	41,24	77,15	37,99	16,42
Usp19	10,36	16,62	10,4	6,36
Usp7	33,47	41,83	29,12	32,91
Usp31	3,37	9,08	2,74	1,03
Usp38	4,61	8,48	2,87	5,07
Usp11	9,22	15,43	11,57	9,11
Usp32	13,01	73,83	33,44	14,41
Usp47	20,02	59,6	41,22	28
Usp37	7,38	19,56	15,5	16,56
Usp39	4,39	10,05	5,83	8,19
Usp36	4,73	3,94	1,16	1,11
Usp27x	3,48	2,69	0,29	0
Usp6nl	3,7	4,8	5,47	2,06
Usp22	28,1	15,28	1,43	0,52
Usp48	32,7	14,43	8,06	8,43
Usp12	6,67	6,62	4,94	15,29
Usp11	41	8,72	1,79	6,97
Usp35	8,55	1,39	0,46	0,06
Usp49	4	7,07	7,42	7,52
Usp16	12,92	25,75	32,11	36,11
Usp51	0,14	0,48	0,82	5,24
Usp1	10,56	21,4	100,04	47,7
Usp42	7,47	15,55	42,71	40,61
Usp8	28,16	33,25	103,33	48,87
Usp4	9,74	11,25	13,63	13,89
Usp15	17,33	24,08	50,52	63,35
Usp29	0,17	0,41	1,71	2,63
Usp44	0,58	1,46	8,45	24,44
Usp20	3,11	3,31	5,57	11,15
Usp45	7,58	4,18	5,42	7,41
Usp21	10,02	2,76	7,9	3,27
Usp46	4,66	3,45	3,55	6,59
Usp30	3,6	3,08	6,06	5,93
Usp25	6	1,73	14,9	41,3
Usp2	1,6	2,09	58,64	40,43
Usp50	0,49	0,58	19,72	31,74

Tabla S2. Expresión de diferentes USPs (mRNA) descrita por Margolin et al. (2014).

6dpp, espermatogonia; 10dpp leptotena; 12dpp zigotena; 14dpp paquitena; 16 dpp paquitena-diotena; 18dpp espermatocitos secundarios; 20dpp espermátida redonda; 38 dpp ratón adulto. Unidades: RPKM (*reads per Kilobase per million mapped reads*).

Min									Max
	6dpp	10dpp	12dpp	14dpp	16dpp	18dpp	20dpp	38dpp	
Usp37	19,28	31,96	29,67	26,33	24,01	20,88	19,55	24,25	
Usp26	22,46	35,77	28,20	26,08	20,93	11,53	9,28	1,91	
Usp9x	23,50	38,57	31,72	26,88	21,51	13,56	11,88	6,89	
Usp53	13,20	21,40	16,60	14,08	11,04	8,89	10,38	5,52	
Usp33	17,68	22,67	18,06	16,28	16,79	11,64	13,32	3,16	
Usp14	18,03	25,38	19,78	20,24	18,29	15,30	14,22	7,89	
Usp46	12,56	19,07	15,33	13,55	12,89	13,55	12,57	11,01	
Usp45	3,86	6,51	4,02	4,28	4,82	4,87	5,82	3,85	
Usp28	30,73	31,11	31,38	26,69	25,23	17,54	16,53	6,60	
Usp3	23,55	30,36	32,21	29,69	22,35	15,74	14,58	7,27	
Usp9y	1,73	3,77	3,79	4,10	2,35	1,57	1,96	0,91	
Usp11	16,88	13,72	15,98	11,25	8,87	5,77	4,77	2,44	
Usp35	3,09	2,29	1,84	1,63	0,94	0,67	1,13	0,97	
Usp38	7,71	9,78	9,85	10,90	8,90	8,75	7,48	7,82	
Usp22	59,23	44,31	43,64	33,56	29,21	19,81	16,64	7,98	
Usp48	25,64	21,63	21,75	18,94	18,87	13,17	12,79	8,56	
Usp13	0,75	0,47	0,53	0,34	0,69	0,27	0,23	0,02	
Usp25	7,04	6,00	4,81	5,96	3,97	4,32	4,03	32,76	
Usp12	8,51	10,19	12,36	11,18	11,29	6,93	7,29	14,96	
Usp40	7,66	9,09	9,14	13,84	12,92	11,34	9,79	4,50	
Usp7	40,10	44,71	42,68	49,40	47,25	45,11	41,41	51,80	
Usp24	4,23	5,07	5,19	5,89	6,26	5,60	6,49	3,30	
Usp18	1,03	0,62	0,86	0,42	0,92	0,98	0,62	0,38	
Usp10	20,76	13,51	18,49	16,50	19,11	14,80	12,16	9,48	
Usp32	19,63	40,05	41,44	47,38	50,53	62,04	65,97	37,34	
Usp43	1,41	0,44	0,53	0,83	0,30	0,35	0,52	0,98	
Usp54	8,30	10,37	14,54	16,67	17,18	14,16	11,53	6,65	
Usp52	20,35	15,74	19,49	20,86	17,87	18,17	19,62	11,66	
Usp11	13,07	16,49	18,72	17,57	20,34	28,36	27,89	19,44	
Usp39	23,63	20,49	17,76	23,51	26,71	31,68	26,25	21,59	
Usp47	25,41	36,61	38,63	48,64	48,42	62,73	68,58	57,81	
Usp15	13,11	20,38	20,98	30,34	32,87	46,44	46,80	78,05	
Usp16	9,11	12,95	16,74	19,82	18,62	21,18	22,95	26,52	
Usp1	9,96	12,11	10,39	12,56	19,31	43,84	57,99	50,03	
Usp8	24,17	31,50	38,14	46,93	45,84	81,37	96,31	104,00	
Usp42	14,50	17,39	20,79	23,98	25,90	32,47	37,00	37,01	
Usp5	33,10	22,64	31,03	35,51	38,75	31,92	30,11	20,99	
Usp4	27,31	28,39	32,02	40,50	42,21	50,68	48,29	59,88	
Usp21	18,16	13,33	15,05	16,37	15,34	19,10	20,36	17,18	
Usp44	1,47	0,80	0,98	0,96	2,03	3,98	4,32	38,32	
Usp49	7,89	7,29	9,02	8,73	8,38	8,74	8,52	13,29	
Usp6nl	8,29	6,51	7,65	7,34	6,90	7,29	8,30	7,49	
Usp20	6,34	4,10	6,59	7,28	8,79	15,15	16,00	29,45	
Usp50	0,04	0,00	0,33	0,00	0,20	0,06	0,60	62,96	
Usp30	11,30	7,66	8,69	10,57	9,44	12,89	12,94	12,83	
Usp29	0,17	0,11	0,19	0,44	0,75	0,74	0,51	0,53	
Usp2	5,04	4,48	5,75	15,15	25,94	56,47	66,82	141,24	
Usp19	58,70	40,41	60,20	64,58	67,27	54,04	51,44	43,37	

1.2 Una variante en HSF2BP causa POI a través de su nuevo interactor BRME1

Tabla S3. Secuenciación del exoma completo y datos de mapeo de las pacientes de POI y la hermana no afectada (NA).

	Lecturas pareadas (millones)	Profundidad media	Cobertura > 25 X	Mismatch Lectura 1	Mismatch Lectura 2
POI III-2	56,2	79,05 X	80,1 %	0,13 %	0,23 %
POI III-3	36,5	26,81 X	89,86 %	0,15 %	0,29 %
NA III-10	31,1	38,39X	92,1 %	0,16 %	0,25 %

Tabla S4. Filtrado de las variantes identificadas por secuenciación del exoma completo.

Variantes localizadas en	POI III-2	POI III-3	NA III-10
Total	118545	105899	107840
SNPs	108106	96854	98621
Indels	10439	9045	9219
Variantes en homocigosis	42695	39741	39640
Variantes en heterocigosis	75850	66158	68200
Filtros aplicados a las variantes			# de variantes
Cobertura > 9 lecturas en los 3 exomas			130021
Homocigosis en pacientes o Heterocigosis en la hermana no afectada			8897
En regiones codificantes			4553
Con efecto en la proteína (no sinónimas)			2295
No homocigosis en los controles fértiles no relacionados			66
MAF < 1% en GnomAD			5
Con información funcional coherente			1

Tabla S5: Predicciones de patogenidad de la variante S167L en *HSF2BP*.

Software	Score para la variante	Umbral de patogenidad	Predicción patogenidad
SIFT	0,114	< 0,05	Tolerada
SIFT4G	0,004	< 0,05	Dañina
PolyPhen 2	0,135	> 0,8	Benigna (indicada como probablemente dañina en GnomAD)
M-CAP	0,0232	> 0,025	Tolerada (en el límite)
FATHMM-MKL	0,8975	> 0,5	Dañina
LRT	0,000293	Score is a p-value	Deletérea
MutationTaster	0,994943	0,5	Causante de enfermedad
MutationAssessor	1,87	> 0,65	Baja
FATHMM	-0,06	< -1,5	Tolerada
FATHMM-MKL coding	0,89755	0,5 (default) 0,80 (stringent)	Deletérea
PROVEAN	-2,88	-2,28	Dañina
MetaSVM	-0,8926	0	Tolerada
MetaLR	0,1613	0,5	Tolerada
REVEL	0,068	0,5 (default) 0,75 (stringent)	Benigna
DANN	0,996469	0,96	Dañina
CADD	5,470242	1,75	Dañina
GERP++ RS	4,36	> 4,4	Conservada
phyloP100way_vertebrate	5,335	> 1,6	Altamente conservada

Figura S1. El residuo Ser167 en proteína HSF2BP está altamente conservado en 99 mamíferos. Alineamiento múltiple de los ortólogos de HSF2BP con CLUSTAL W 2.0. La Ser167 humana (verde) está conservada en 97 de las 99 secuencias disponibles de mamíferos. Solo se han identificado dos cambios (resaltados en amarillo) en *Cavia aperea* (Cape, cobaya brasileña) y *Cavia porcellus* (Cpor, cobaya).

ENSP00000291560_Hsap/1-334

ENSTTRP000000000680	Ttru/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVQ---	ELSDSESQFVFALAGIVTNVAA
ENSGGROF000000029257	Ggor/1-339	ILGGGKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPFY000000012806	Pabe/1-326	ILGGDKALKFFSITGQTMESFVKSLDGDVQ---	ELSDSESQFVFALAGIVTNVAA
ENSMFAP000000037975	Mfas/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVQ---	ELSDSDENQFVFALAGIVTNVAA
ENSMAUP000000020636	Maur/1-344	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSAMEP000000011987	Amel/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSSARP000000010935	Sara/1-237	ILAGGAKPKFFRITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSDORP000000027761	Dord/1-337	ILAGEKALKFFRITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSHGLP000000012308	Hqfe/1-336	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSTSY0000000000982	Csyr/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSCWAP000000028492	Cwag/1-334	ILGGDKALKFFFITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPEMP000000025118	Pmba/1-338	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSOGAP000000008822	Ogar/1-334	ILEGDKALKFFFNITGQTMESFVKSLDGDVK---	ELNSDENQFVFALAGIVTNVAA
ENSSSUP000005002765	Ssur/1-367	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSSBOP000000029851	Sbbo/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELNSDENQFVFALAGIVTNVAA
ENSCCAP000000027862	Ccap/1-331	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELNSDENQFVFALAGIVTNVAA
ENSCANP000000035144	Capa/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPVAP0000000009392	Pvam/1-322	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSTSTP000000010567	Itri/1-334	ILGGDKALKFFRITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSBIXP000005017111	Bihy/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSOCUP000000008291	Ocun/1-285	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSETEP000000013339	Etel/1-253	ILGGEKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSDNOP000000003033	Dnov/1-321	ILGGDKALKFFFITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSCAFP000030042050	Cifa/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSBGRP000000033394	Bgru/1-380	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSBIXP000000043346	Bthy/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSVURP000010030416	Vurs/1-338	ILGGDKALKFFSITGQTMESFVKSLDGSIK---	QELSDSDENQFVFALAGIITNVAA
ENSSHAP000000020791	Shar/1-323	ILGGDKALKFFFNITGQTMESFVKSLDGNIK---	QELSDSDENQFVFALAGIITNVAA
ENSCGRP000015029988	Cgpi/1-375	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPITP000000018380	Ptal/1-325	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSHGLP000100007014	Hgma/1-336	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPTRP000000024023	Ptro/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVQ---	ELSDSESQFVFALAGIVTNVAA
ENSPSMP000000001155	Psim/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELNSDENQFVFALAGIVTNVAA
ENSCATP000000038233	Caty/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPPRP000000018732	Ppar/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSRROP000000017466	Rrox/1-305	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSSDAP000000017273	Sdau/1-335	ILGGDKALKFFRITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSJAP0000000007064	Jjac/1-272	ILGGDKALKFFFNITGQTMESFVKSLDEDVK---	ELSDSESQFVFALAGIVTNVAA
ENSRNOP000000001580	Rnor/1-338	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSCDRP000005018641	Cdro/1-334	ILAGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSC LAP000000009399	Clan/1-334	MLGGDKASKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSPAP000000023666	Ppan/1-334	ILGGDKALKFFSITGQTMESFVKSLDGDVQ---	ELSDSESQFVFALAGIVTNVAA
ENSFADP000000004802	Fdam/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSM LUP000000007048	Mluc/1-338	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSMOCP000000006501	Moch/1-339	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSLCNP000005011741	Lcan/1-334	MLGGDKALKFFFNITGQTMESFVKSLDEDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSM SIP000000014116	Mspi/1-338	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSOEP000000015972	Odeg/1-338	MLGGDKASKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSMICP000000004266	Mmur/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSOARP000000012082	Oari/1-334	ILGGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
MGP_PahariEiJ_P0056165_Mpah/1-140		ILGGDKALKFFFNITGQTMESFVKSLDGNVK---	ELSDSDENQFVFALAGIVTNVAA
ENSEASP00005002169	Eaas/1-365	ILGGDKALKFFSITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA
ENSMUGP000000008716	Mung/1-338	ILRGDKALKFFFNITGQTMESFVKSLDGDVK---	ELSDSDENQFVFALAGIVTNVAA

ENSP00000291560_Hsap/1-334 ILGGDKALKFFSITGQTMSFVKSLDGDV--QELDSDESQFVFLAGIVTNVAA
 ENSGAGP00000000454_Gaga/1-336 ILGGSKAVKFFTITGQTMSFVRSLGEDIKHQDLDSDENQFVFLAGIVTNVAA
 ENSCCEP00000018733_Ccae/1-336 ILGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSNMEP00000016416_Nmel/1-382 ILGGSKAAKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSSCAP00000015316_Scan/1-347 ILGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSLCOF00000002986_Lcor/1-336 ILGGSKAGKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSMUNP00000011578_Mund/1-361 ILGGSKAVKFFTITAQTMSFVKSLGEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSCPUP00000017770_Cpug/1-301 LLGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSEGOP00005008603_Egou/1-335 ILGGTKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSSHBP00005011292_Shah/1-305 ILGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSPLCP00000008530_Pcol/1-336 ILGGSKAAKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSACAP00000017574_Acar/1-301 ILGSSKAVKFFFIAGQTMSFVRSLSEDIKQNMDSDESQFVLGLAGIVTNIAA
 ENSPTXP00000015356_Ptex/1-337 ILGSAQAVKFFNIAGQTMSFVCSLNEDSKHQMDSDESQFVLGLAGIVTNVAA
 ENSARWP00000026189_Arow/1-329 ILGGSKAVKFFTITAQTMSFVKSLNEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSCPBP00000017720_Cpbe/1-342 ILGGSKAVKFFTITGQTMSFVRSLGEDTKHQDLDSDENQFVFLAGIVTNVAA
 ENSACOF00000010942_Acol/1-326 ILGGTKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSFALP00000002102_Falb/1-336 ILGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSSMRP00000029796_Smer/1-301 ILGSSKAVKFFRLAGQTMSFVRSLSSENIKQQDMESDESQFVLGLTGIVTNVAA
 ENSNPEP00000012775_Nper/1-329 ILGASKAVKFFTITAQTMSFVKSLNEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSAPLP00000003727_Appl/1-336 ILGGSKAAKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSLSDP00000011344_Lsdo/1-339 ILGGTKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSSPUP00000023120_Spun/1-327 ILGGNKAAFTFSIAGQTMSFVKSLGEDIKQDMDSDENQFVFLAGIVTNVAA
 ENSNSUP00000009799_Nscu/1-335 ILGSAQAVKFFNIAGQTMSFVCSLNEDSKHQMDSDESQFVLGLAGIVTNVAA
 ENSPMJP00000014199_Pmaj/1-336 ILGGSKAVKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSPVIP00000026890_Pvit/1-301 ILGNSKAIIFNNIAGQTMSFVRSLSDDIKQDMDSDDESQFVLGLAGIVTNVAA
 ENSCPIC00010010640_Cpic/1-340 ILGGSKAAKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSCABP00000014577_Cabi/1-327 ILGGSKAVKFFTITGQTMSFVRSLGEDIKHQDLDSDENQFVFLAGIVTNVAA
 ENSAHAP00000003935_Ahaa/1-311 ILGGSKAVKFFTITAQTMSFVKSLNEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSCJPP000000509998_Cjap/1-363 ILGGSKAAKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA
 ENSMVIP00005009255_Mvit/1-336 ILGGSKAGKFFTITAQTMSFVKLSLEDMDKQDLDSDENQFVFLAGIVTNVAA

ENSPSIP00000008390_Psin/1-322
 ENSMGAP00000015454_Mgal/1-322
 ENSABRP00000017191_Abra/1-306
 ENSPMRP00000034335_Pmur/1-345
 ENSDNVP00000009224_Dnov/1-334
 ENSACCP00020000703_Acch/1-340
 ENSJHYP00000017704_Jhye/1-409
 ENSSCUP00000016503_Scau/1-336
 ENSACUP00000019635_Acun/1-326
 ENSAOWP00000021620_Aowe/1-337
 ENSACDP00005009735_Acyg/1-306
 ENSCPGP00000014458_Cpyg/1-336
 ENSTGUP00000006085_Tgut/1-335
 ENSPSTP00000012794_Pcri/1-320
 ENSZALP000000021794_Zalb/1-301
 ENSVKKP00000020715_Vkom/1-321
 ENSANIP00000018222_Anis/1-336
 ENSGALP00000026071_Ggal/1-336
 ENSCPRP00005023694_Cpor/1-308

ILGGSKAVKFFTTITGQTLESFVRS LGEDTKHQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAAKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAAKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGSSKAIKFFSIAGQTMSFVRSLSEDIKQQDMDSDESQFVLGLAGIVTNVAA
 ILGGSKAVKFFTTITAQTMSFVKSLNEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAVKFFTTIAAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAVKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAVKFFTTITAQTMSFVKSLNEDVKKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAVKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAVKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAAKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 LLGGSKAVKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGTKAVKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAAKFFTTITAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGSSKAVKFFNIAGQTMSFVRS LTEDMKQQDLDS DESQFVLGLAGIITNVAA
 ILGGSKAVKFFTTIAAQTMSFVKSLSEDMKQQDLDS DENQFVLALAGIVTNVAA
 ILGGSKAAKFFTTITAQTMSFVKSLSEDTKQQDLDS DENQFVLALAGIVTNVAA
 ILGG-----VTSLSHKMS-----LVII LCSVV-DVAA

ENSPHSA00000291560_Hsap/1-334

ENSOABP00000040411_Oaur/1-343
 ENSGAFP00000024427_Gaff/1-336
 ENSMMOP00000005954_Mmol/1-304
 ENSGACP000000020225_Gacu/1-322
 ENSPKIP00000027491_Pkin/1-345
 ENSSORP00005023028_Sorb/1-309
 ENSIPUP00000020465_Ipun/1-350
 ENSELUP000000033600_Eluc/1-369
 ENSORLP00020013137_Olhn/1-355
 ENSFHPEP00000033032_Phet/1-335
 ENSSMAP00000016094_Smax/1-348
 ENSPNYP000000029528_Nnye/1-356
 ENSLCAP00010015134_Pcal/1-361
 ENSKMAP00000016215_Kmar/1-321
 ENSATEP00000017112_Ates/1-324
 ENSMZEPE00005025586_Mzeb/1-341
 ENSXCOP00000027267_Xcob/1-294
 ENSONIP00000004724_Onil/1-322
 ENSPMPEP00000022553_Pmex/1-346
 ENSLCRP00005033625_Lcro/1-383
 ENSTRUP000000056411_Trub/1-394
 ENSGMOP00000002425_Gmor/1-311
 ENSXMAP00000030843_Xmac/1-338
 ENSPNAP00000012466_Pnat/1-349
 ENSAOCPP00000007621_Aoce/1-327
 ENSHBUP00000021919_Hbur/1-239
 ENSORLP00015015570_Olhs/1-379
 ENSTNIP00000017492_Tnig/1-337
 ENSLOCPP00000001492_Locu/1-342
 ENSPMGP00000019233_Pmag/1-307
 ENSPREP00000026460_Pret/1-339
 ENSMALPP00000017207_Malb/1-332
 ENSMMDP00005003945_Mmur/1-345
 ENSHCOP00000014782_Hcom/1-352
 ENSENLP00000003494_Enau/1-355
 ENSORLP00000025611_Olat/1-355
 ENSGWIP00000005420_Gwil/1-365
 ENSPLAP00000024128_Plat/1-312
 ENSXETP00000025892_Xtro/1-315
 ENSSLDP00000019834_Sldo/1-339
 ENSMAMP00000005262_Marm/1-330
 ENSACLP00000010749_Acal/1-341
 ENSSPAP00000019130_Spar/1-336
 ENSAPOP00000016808_Apol/1-345
 ENSSDUP00000030608_Sdum/1-354
 ENSACIP00000022763_Acit/1-312
 ENSPPOP00000000434_Pfor/1-324
 ENSDARP00000114662_Drer/1-365
 ENSSFAP00005047030_Sfas/1-327
 ENSSAUP00010012427_Saur/1-348
 ENSCVAP00000003098_Cvar/1-257
 ENSSFOP00015035913_Sfor/1-362
 ENSDCDP00000028687_Sclu/1-335
 ENSAMXP00005023676_Ampa/1-356

ILGGD-KALKFFSITGQTMSFV-KSLDGDV----QELDSDESQFVFALAGIVT-NVAA

WLADG-KLQAF LAVAQAQTLESFV-TSLDEEEKTQTEDYNSHEHQFVLAVAGTIT-NVAA
 WLADG-KLP SFLTVAQAQTLESFV-KSMDDEIKTQPGDHNSQEHQFVLALVGTIT-NIAA
 WL--G-KLQSFLEVAQAQTLESFV-ASLDQEVKNLTEDHNSQEHQFVLALAGTIT-NIAA
 WLTDG-KLP SFLTVAQAQTLESFV-GSLDDDANTLAEDHNP-DHQFVLAWLEP LLSDVAA
 ILAGS-YIEPFLALAGAQTLESFV-SSLN-PVKMEKEDSSCQEHQFVLAVAGIIT-NIAA
 ----G-KLRSFVAVAAQTLESFV-RSLDEDAKTQTEDQDSEHQFVLALAGIIT-NIAA
 VLADG-KLEPFLSVAGQTLESFV-KSLDEEEKPQQQYNSHEHQFVLALAGVIT-NIAA
 ILVEG-NLQSF LSMAGQILES YV-KSLDGEARAEQQDSNSHEQQFVLALAGVVT-NVAA
 WLADG-KLLSFVSVSAQAQTLESFV-KSLDEEVKDQSDDYNSHEHQFVLALVGTIT-NVAA
 WLADG-KLP SFLTVAQAQTLESFV-KSMDDEMKT--EPPGSHHQFVLALAGTIT-NVAA
 WLADG-KLQSF LAVAQAQTLESFV-GSLDAEVKNLTEDHNSQEHQFVLGLAGTIT-NIAA
 WL--G-KLQAF LAVAQAQTLESFV-TSLDEEEKSQTEDYNSHEHQFVLAVAGTIT-NVAA
 WLQ-G-KLQSF LTVAGQTLESFV-RSLDSEVKTQTEDHNSQEHQFVLALAGTIT-NIAA
 WL--G-KLQSF LIVATQTLESFV-KSLDDEMKTQTEDPNSTEHQFVLALVGTIT-NIAA
 WL--G-KLKSFLTVAQAQTLESFV-GSLDEEVKTQSEDLNSQEHQFVLALAGTIT-NIAA
 WLADG-KLQAF LAVAQAQTLESFV-TSLDEEEKSQTEDYNSHEHQFVLAVAGTIT-NVAA
 WL--G-KLP SFLTVAQAQTLESFV-KSMDDEIKTQPGDHNSQEHQFVLALVGTIT-NIAA
 WL--G-KLQAF LAVAQAQTLESFV-TSLDEEEKTQTEDYNSHEHQFVLAVAGTIT-NVAA
 WLADG-KLASFLTVAQAQTLESFV-KSMDDEIKTQPGDHNSQEHQFVLALVGTIT-NIAA
 WLADG-KLQSF LVLAQAQTLESFV-RSLDEEDGKAQTEDHNSQEHQFVLALAGTIT-NIAA
 WLADG-MLQSF LTVAGQTLESFV-RSLDDEVKTQTEDHSSQEHQFVLALAGTIT-NIAA
 XXXXG-KLP SFLDLAGQTLQSFV-QSGEDDADL---DQTSTEHQFVLAWLSGQ--DVAA
 WLTDG-KLP SFLTVAQAQTLESFV-KSMDDEIKTQPGDHNSQEHQFVLALVGTIT-NIAA
 ILADG-KLEPFLKVAGQTLESFV-KSLDEEEKPQ-QNYSSHEHQFVLALAGIIT-NIAA
 WL--G-KLQAF LTVAGQTLESFV-RSLDDEMKTQTEDHNSQEHQFVLALAGTIT-NIAA
 WLADG-KLQAF LAVAQAQTLESFV-TSLDEEEKSQTEDYNSHEHQFVLAVAGTIT-NVAA
 WLADG-KLLSFVSVSAQAQTLESFV-KSLDEEVKDQSDDYNSHEHQFVLALVGTIT-NVAA
 WLADG-KLP PFLAVATQTLESFV-TSLDDEVKTQTEDHNSQEHQFVLGLAGTIT-NIAA
 ILAGS-RVQTF LTVAGQTLESFV-KSLDGDGRPGQDLSSDENQFVLALAGVIT-NIAA
 WLSHDHVKLQSF LLAQAQTLESFV-CSLDEEDVKVQTEDPNSQEHQFVLALAGTIT-NIAA
 WLADG-KLP CFLTVAQAQTLESFV-KSMGDEIKTQPGDHNSQEHQFVLALVGTIT-NIAA
 WLADG-KLQAF LAVTAQTLEIFV-KSLDEEVKTQVEDHNCQEHQFVLALARTIT-NIAA
 WVAQG-KLP SFLTVAQAQTLESFV-KSLDEEVNAQSQEDYNTHEHQFVLGLAGVIT-NVAA
 WLADG-SLP AFVAVATQTLGSFV-RTLDEEDVKTRSEDCTQEQYQFVLALAGTIT-NIAA
 WLAD----AFLAIAQAQTLESFI-GSLDVELKPQTEDQNSQEHQFVLALAGTIT-NIAA
 WLADG-KLLSFVSVSAQAQTLESFV-KSLDEEVKDQSDDYNSHEHQFVLALVGTIT-NVAA
 WLTDG-KLP SFLTVAQAQTLESFV-TSLDEETKTQTEDHNSQEHQFVLALAGTIT-NIAA
 WL--G-KLSSFLTVAQAQTLESFV-KSMDDEIKTQPSDHNSQEHQFVLALVGTIT-NIAA
 ILGGS-KAPMFFSLAQAQTLSFV-RSLTSE---QPQDEENESHFVLGLAGVIT-NVAA
 WL--G-KLQAF LAVAQAQTLESFV-RSLDVELKTQTEDQNSQEHQFVLALAGTIT-NIAA
 WLSDG-KLP SFLGLAQAQTLESFV-KSLDDEVKTQIEDHNSQEHQFVLALAGTIT-NIAA
 WLADG-KLQAF LAVAQAQTLESFV-TSLDEEEKSQTEDYNSHEHQFVLAVAGTIT-NVAA
 WL--G-KLQSF LTVAGQTLESFV-SSLDEEMKTQTEDHNSQEHQFVLALAGTIT-NIAA
 WLADG-KLRSFLTVAQAQTLESFV-RSLDEDMKTQTEDHNSQEHQFVLALAGTIT-NIAA
 WLADG-KLP SFLTVAQAQTLESFV-RSLDVELKTQTEDQNSQEHQFVLALAGTIT-NIAA
 WL--G-KLQAF LAVAQAQTLESFV-KSLDEDMTTQTDDHNSHELQFVLAVAGTIT-NIAA
 WLADG-KLSSFLTVAQAQTLESFV-KSMDDEIKTQPSDHNSQEHQFVLALVGTIT-IDIAA
 I LANG-KLEDFLSIAGQTLTFTI-KLLVDEAKPQQSD-NSHEHNFVLALAGVVT-NIAA
 W---G-KLP SFLTVAQAQTLESFV-RSLDEETKTQTEDPSSQEHQFVLALAGSIT-NIAA
 WLADG-KLQSF LSVAAQAQTLESFV-RSLDEEDVKTQTEDHTSQEHQFVLALAGTIT-NIAA
 WLSDG-KLHTFLNIAQAQTLESFV-KSLDEEIKPPAEDHSSNENQFVLALVGTIT-NIAA
 ILADR-KAEAF LAVAQAQTLESFVQSLDREGKVEQQT-SSQEMQFVLALAGIVT-NIAA
 ML--G-KLP SFLSLAGQTLLESFV-KSLGEEVKPPQVEHNSHEHQFVLALAGVVT-NIAA
 ILADG-KLEPFLSVAGQTLESFV-KSLDEEEKPQQQYNSHEHQFVLALAGVIT-NIAA

ENSNMLP00000006028_Nmel/1-345
ENSOMEF00000009520_Omel/1-341
ENSCHAP00000000583_Char/1-368
ENBSLPL000000013271_Bspl/1-344
ENSCGOP000000002370_Cgob/1-345
ENSAPEP000000012225_Aper/1-285
ENSEEEP000000024604_Eele/1-329
ENSCSEP000000009624_Csem/1-343
ENSLACP000000006144_Lcha/1-340
ENSEBUP000000012007_Ebur/1-327
ENSPMAP000000009832_Pmar/1-320

WLADQWKLQSFVVVAAQTLESFV-RSLDEDVKVQTEDPNSQEHKFVLALAGTIT-NIAA
WLVDG-KLPSFLNVAAQTLESFG-KSLDEEVKDQSEDYNSQEHQFVLAVAGTVT-NVAA
ILADG-KLEPFLSVAGQTLESFV-KSLDQDSKLPQQDSNSHEHQFVLALAGVVT-NLAA
WLADG-KLQSFVCVAAQTLESFV-GSLDEEVKA--EDHNSQEHQFVLALAGTIT-NIAA
WLADG-KLQTFLSVAAQTLESFV-RSLDEEAKSQTEDHNSQEHQFVLALAGTIT-NVAA
-----TLESFV-RSLDEDMKTQTEDHNSHEHQFVLALAGTIT-NIAA
I---G-LFKPFLT VAGQTLESFI-KSLDEEETPQQQYNNSHEHQFVLALAGIIT-NVAA
WLADG-KLPSFLGVAAQTLESFV-DSLDAEVKTQTEDQNSLEHQFVLALAGTVT-NIAA
ILGGS-KAVKFFT VAGQTIESFV-KSLDGDVK-QQQDLSDENQFVMGLTGIVT-NIAA
ILNGA-KVEEFFTLTSQTLDSFM-KSMIEEGK--RKNIDEEENQFVLAMAGTIT-NVAA
ILVGC-KVKEFFTLASQTVESYV-KSMPGDFK--TEDAGGDESQFVLAMAGIVT-NIAA

BIBLIOGRAFÍA

- Abir R, Fisch B, Nahum R, Orvieto R, Nitke S, Rafael Z Ben (2001). "Turner's syndrome and fertility: Current status and possible putative prospects." *Hum. Reprod. Update* 7:603–610
- Ahmed EA, de Rooij DG (2009). "Staging of mouse seminiferous tubule cross-sections." *Methods Mol. Biol.* 558:263–277
- Arafat M, Zeadna A, Levitas E, Har Vardi I, Samuelli B, Shaco-Levy R, Dabsan S, Lunenfeld E, Huleihel M, Parvari R (2020). "Novel mutation in USP26 associated with azoospermia in a Sertoli cell-only syndrome patient." *Mol Genet Genomic Med* 8: . <https://doi.org/10.1002/mgg3.1258>
- Aramaki S, Hayashi K, Kurimoto K, Ohta H, Yabuta Y, Iwanari H, Mochizuki Y, Hamakubo T, Kato Y, Shirahige K, Saitou M (2013). "A mesodermal factor, T, specifies mouse germ cell fate by directly activating germline determinants." *Dev Cell* 27:516–529 . <https://doi.org/10.1016/j.devcel.2013.11.001>
- Asadpor U, Totonchi M, Sabbaghian M, Hoseinifar H, Akhound MR, Zari Moradi S, Haratian K, Sadighi Gilani MA, Gourabi H, Mohseni Meybodi A (2013). "Ubiquitin-specific protease (USP26) gene alterations associated with male infertility and recurrent pregnancy loss (RPL) in Iranian infertile patients." *J Assist Reprod Genet* 30:923–931 . <https://doi.org/10.1007/s10815-013-0027-9>
- Ashley T, Walpita D, Rooij DD de (2001). "Localization of two mammalian cyclin dependent kinases during mammalian meiosis." *J Cell Sci* 114:685–693
- Baarends WM, Hoogerbrugge JW, Roest HP, Ooms M, Vreeburg J, Hoeijmakers JHJ, Grootegoed JA (1999). "Histone ubiquitination and chromatin remodeling in mouse spermatogenesis." *Dev Biol* 207:322–333 . <https://doi.org/10.1006/dbio.1998.9155>
- Baarends WM, van der Laan R, Grootegoed JA (2000). "Specific aspects of the ubiquitin system in spermatogenesis." *J Endocrinol Invest* 23:597–604 . <https://doi.org/10.1007/BF03343782>
- Baker SM, Plug AW, Prolla TA, Bronner CE, Harris AC, Yao X, Christie DM, Monell C, Arnheim N, Bradley A, Ashley T, Liskay RM (1996). "Involvement of mouse Mlh1 in DNA mismatch repair and meiotic crossing over." *Nat Genet* 13:336–342 . <https://doi.org/10.1038/ng0796-336>
- Balasch J, Gratacós E (2012). "Delayed childbearing: Effects on fertility and the outcome of pregnancy." *Curr. Opin. Obstet. Gynecol.* 24:187–193
- Baltus AE, Menke DB, Hu YC, Goodheart ML, Carpenter AE, De Rooij DG, Page DC (2006). "In germ cells of mouse embryonic ovaries, the decision to enter meiosis precedes premeiotic DNA replication." *Nat Genet* 38:1430–1434 . <https://doi.org/10.1038/ng1919>
- Bannister LA, Reinholdt LG, Munroe RJ, Schimenti JC (2004). "Positional cloning and characterization of mouse mei8, a disrupted allele of the meiotic cohesin Rec8." *Genesis* 40:184–194 . <https://doi.org/10.1002/gene.20085>
- Barlow AL, Benson FE, West SC, Hultén MA (1997). "Distribution of the Rad51 recombinase in human and mouse spermatocytes." *EMBO J* 16:5207–5215 . <https://doi.org/10.1093/emboj/16.17.5207>
- Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, Coop G, De Massy B (2010). "PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice." *Science (80-)* 327:836–840 . <https://doi.org/10.1126/science.1183439>
- Baudat F, Imai Y, De Massy B (2013). "Meiotic recombination in mammals: Localization and regulation." *Nat Rev Genet* 14:794–806 . <https://doi.org/10.1038/nrg3573>
- Baudat F, Manova K, Yuen JP, Jasin M, Keeney S (2000). "Chromosome synapsis defects and sexually dimorphic meiotic progression in mice lacking Spo11." *Mol Cell* 6:989–998 . [https://doi.org/10.1016/S1097-2765\(00\)00098-8](https://doi.org/10.1016/S1097-2765(00)00098-8)
- Bedard N, Yang Y, Gregory M, Cyr DG, Suzuki J, Yu X, Chian RC, Hermo L, O'Flaherty C, Smith CE, Clarke HJ, Wing SS (2011). "Mice lacking the USP2 deubiquitinating enzyme have severe male subfertility associated with defects in fertilization and sperm motility." *Biol Reprod* 85:594–604 . <https://doi.org/10.1095/biolreprod.110.088542>
- Bekker-Jensen S, Mailand N (2011). "The ubiquitin- and SUMO-dependent signaling response to DNA double-strand breaks." *FEBS Lett* 585:2914–2919 . <https://doi.org/10.1016/j.febslet.2011.05.056>
- Bellani MA, Boateng KA, McLeod D, Camerini-Otero RD (2010). "The Expression Profile of the Major Mouse SPO11 Isoforms Indicates that SPO11 β Introduces Double Strand Breaks and Suggests that SPO11 α Has an Additional Role in Prophase in both Spermatocytes and Oocytes." *Mol Cell Biol* 30:4391–4403

- . <https://doi.org/10.1128/mcb.00002-10>
- Bellani MA, Romanienko PJ, Cairatti DA, Camerini-Otero RD (2005). "SPO11 is required for sex-body formation, and Spo11 heterozygosity rescues the prophase arrest of Atm-/- spermatocytes." *J Cell Sci* 118:3233–3245 . <https://doi.org/10.1242/jcs.02466>
- Boateng KA, Bellani MA, Gregoretti I V., Pratto F, Camerini-Otero RD (2013). "Homologous Pairing Preceding SPO11-Mediated Double-Strand Breaks in Mice." *Dev Cell* 24:196–205 . <https://doi.org/10.1016/j.devcel.2012.12.002>
- Bolcun-Filas E, Handel MA (2018). "Meiosis: the chromosomal foundation of reproduction." *Biol Reprod* 99:112–126 . <https://doi.org/10.1093/biolre/iyoy021>
- Bolcun-Filas E, Schimenti JC (2012). "Genetics of meiosis and recombination in mice." *Int Rev Cell Mol Biol* 298:179–227 . <https://doi.org/10.1016/B978-0-12-394309-5.00005-5>
- Bose R, Manku G, Culty M, Wing SS (2014). "Ubiquitin-proteasome system in spermatogenesis." *Adv Exp Med Biol* 759:181–213 . https://doi.org/10.1007/978-1-4939-0817-2_9
- Bowles J, Knight D, Smith C, Wilhelm D, Richman J, Mamiya S, Yashiro K, Chawengsaksophak K, Wilson MJ, Rossant J, Hamada H, Koopman P (2006). "Retinoid signaling determines germ cell fate in mice." *Science* (80-) 312:596–600 . <https://doi.org/10.1126/science.1125691>
- Brandsma I (2016). "Balancing Pathways in DNA Double Strand Break Repair"
- Brandsma I, Sato K, van Rossum-Fikkert SE, van Vliet N, Sleddens E, Reuter M, Odijk H, van den Tempel N, Dekkers DHW, Bezstarosti K, Demmers JAA, Maas A, Lebbink J, Wyman C, Essers J, van Gent DC, Baarends WM, Knipscheer P, Kanaar R, Zelensky AN (2019). "HSF2BP Interacts with a Conserved Domain of BRCA2 and Is Required for Mouse Spermatogenesis." *Cell Rep* 27:3790-3798.e7 . <https://doi.org/10.1016/j.celrep.2019.05.096>
- Buonomo SBC, Clyne RK, Fuchs J, Loidl J, Uhlmann F, Nasmyth K (2000). "Disjunction of homologous chromosomes in meiosis I depends on proteolytic cleavage of the meiotic cohesin Rec8 by separin." *Cell* 103:387–398 . [https://doi.org/10.1016/S0092-8674\(00\)00131-8](https://doi.org/10.1016/S0092-8674(00)00131-8)
- Caburet S, Arboleda VA, Llano E, Overbeek PA, Barbero JL, Oka K, Harrison W, Vaiman D, Ben-Neriah Z, Garcia-Tunon I, Fellous M, Pendas AM, Veitia RA, Vilain E (2014). "Mutant cohesin in premature ovarian failure." *N Engl J Med* 370:943–949 . <https://doi.org/10.1056/NEJMoa1309635>
- Caburet S, Heddar A, Dardillac E, Creux H, Lambert M, Messiaen S, Tourpin S, Livera G, Lopez BS, Misrahi M (2020). "Homozygous hypomorphic BRCA2 variant in primary ovarian insufficiency without cancer or Fanconi anaemia trait ." *J Med Genet* jmedgenet-2019-106672 . <https://doi.org/10.1136/jmedgenet-2019-106672>
- Caburet S, Todeschini AL, Petrillo C, Martini E, Farran ND, Legois B, Livera G, Younis JS, Shalev S, Veitia RA (2019). "A truncating MEIOB mutation responsible for familial primary ovarian insufficiency abolishes its interaction with its partner SPATA22 and their recruitment to DNA double-strand breaks." *EBioMedicine* 42:524–531 . <https://doi.org/10.1016/j.ebiom.2019.03.075>
- Cahoon CK, Libuda DE (2019). "Leagues of their own: sexually dimorphic features of meiotic prophase I." *Chromosoma* 128:199–214
- Campolo F, Gori M, Favaro R, Nicolis S, Pellegrini M, Botti F, Rossi P, Jannini EA, Dolci S (2013). "Essential role of Sox2 for the establishment and maintenance of the germ cell line." *Stem Cells* 31:1408–1421 . <https://doi.org/10.1002/stem.1392>
- Cariati F, D'Argenio V, Tomaiuolo R (2019). "The evolving role of genetic tests in reproductive medicine." *J. Transl. Med.* 17:267
- Carlosama C, Elzaïat M, Patiño LC, Mateus HE, Veitia RA, Laissue P (2017). "A homozygous donor splice-site mutation in the meiotic gene MSH4 causes primary ovarian insufficiency." *Hum Mol Genet* 26:3161–3166 . <https://doi.org/10.1093/hmg/ddx199>
- Chen H, Lisby M, Symington LS (2013). "RPA Coordinates DNA End Resection and Prevents Formation of DNA Hairpins." *Mol Cell* 50:589–600 . <https://doi.org/10.1016/j.molcel.2013.04.032>
- Christensen GL, Griffin J, Carrell DT (2008). "Sequence analysis of the X-linked USP26 gene in severe male factor infertility patients and fertile controls." *Fertil Steril* 90:851–852 . <https://doi.org/10.1016/j.fertnstert.2007.06.096>

- Clermont Y (1972). "Kinetics of spermatogenesis in mammals: seminiferous epithelium cycle and spermatogonial renewal." *Physiol. Rev.* 52:198–236
- Cloutier JM, Turner JMA (2010). "Meiotic sex chromosome inactivation." *Curr. Biol.* 20
- Cole F, Kauppi L, Lange J, Roig I, Wang R, Keeney S, Jasin M (2012). "Homeostatic control of recombination is implemented progressively in mouse meiosis." *Nat Cell Biol* 14:424–430 .
<https://doi.org/10.1038/ncb2451>
- Cool J, Capel B (2009). "Mixed signals: Development of the testis." *Semin. Reprod. Med.* 27:5–13
- Coop G, Przeworski M (2007). "An evolutionary view of human recombination." *Nat Rev | Genet* 8: .
<https://doi.org/10.1038/nrg1947>
- Costa Y, Speed R, Öllinger R, Aisheimer M, Semple CA, Gautier P, Maratou K, Novak I, Höög C, Benavente R, Cooke HJ (2005). "Two novel proteins recruited by synaptonemal complex protein 1 (SYCP1) are at the centre of meiosis." *J Cell Sci* 118:2755–2762 . <https://doi.org/10.1242/jcs.02402>
- Cox J, Mann M (2008). "MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification." *Nat Biotechnol* 26:1367–1372 .
<https://doi.org/10.1038/nbt.1511>
- da Cruz I, Rodriguez-Casuriaga R, Santinaque FF, Farias J, Curti G, Capoano CA, Folle GA, Benavente R, Sotelo-Silveira JR, Geisinger A (2016). "Transcriptome analysis of highly purified mouse spermatogenic cell populations: gene expression signatures switch from meiotic-to postmeiotic-related processes at pachytene stage." *BMC Genomics* 17:294 . <https://doi.org/10.1186/s12864-016-2618-1>
- Davies AA, Masson JY, McIlwraith MJ, Stasiak AZ, Stasiak A, Venkitaraman AR, West SC (2001). "Role of BRCA2 in control of the RAD51 recombination and DNA repair protein." *Mol Cell* 7:273–282 .
[https://doi.org/10.1016/S1097-2765\(01\)00175-7](https://doi.org/10.1016/S1097-2765(01)00175-7)
- Davis MI, Simeonov A (2015). "Ubiquitin-Specific Proteases as Druggable Targets." *Drug Target Rev* 2:60–64
- De Felici M, Klinger FG, Farini D, Scaldaferrri ML, Iona S, Lobascio M (2005). "Establishment of oocyte population in the fetal ovary: Primordial germ cell proliferation and oocyte programmed cell death." *Reprod Biomed Online* 10:182–191 . [https://doi.org/10.1016/S1472-6483\(10\)60939-X](https://doi.org/10.1016/S1472-6483(10)60939-X)
- de Rooij DG (1973). "SPERMATOGONIAL STEM CELL RENEWAL IN THE MOUSE: I. NORMAL SITUATION." *Cell Prolif* 6:281–287 . <https://doi.org/10.1111/j.1365-2184.1973.tb01617.x>
- de Rooij DG (2001). "Proliferation and differentiation of spermatogonial stem cells." *Reproduction* 121:347–354
- De Rooij DG (1998). "Stem cells in the testis." *Int. J. Exp. Pathol.* 79:67–80
- De Vries L, Behar DM, Smirin-Yosef P, Lagovsky I, Tzur S, Basel-Vanagaite L (2014). "Exome sequencing reveals SYCE1 mutation associated with autosomal recessive primary ovarian insufficiency." *J Clin Endocrinol Metab* 99:E2129–E2132 . <https://doi.org/10.1210/jc.2014-1268>
- Di Pasquale E, Beck-Peccoz P, Persani L (2004). "Hypergonadotropic ovarian failure associated with an inherited mutation of human bone morphogenetic protein-15 (BMP15) gene." *Am J Hum Genet* 75:106–111 . <https://doi.org/10.1086/422103>
- Durcova-Hills G, Tang F, Doody G, Tooze R, Surani MA (2008). "Reprogramming primordial germ cells into pluripotent stem cells." *PLoS One* 3: . <https://doi.org/10.1371/journal.pone.0003531>
- Edelmann W, Cohen PE, Kane M, Lau K, Morrow B, Bennett S, Umar A, Kunkel T, Cattoretti G, Chaganti R, Pollard JW, Kolodner RD, Kucherlapati R (1996). "Meiotic Pachytene Arrest in MLH1-Deficient Mice." *Cell* 85:1125–1134 . [https://doi.org/10.1016/S0092-8674\(00\)81312-4](https://doi.org/10.1016/S0092-8674(00)81312-4)
- Edson MA, Nagaraja AK, Matzuk MM (2009). "The mammalian ovary from genesis to revelation." *Endocr. Rev.* 30:624–712
- Ernst C, Eling N, Martinez-Jimenez CP, Marioni JC, Odom DT (2019). "Staged developmental mapping and X chromosome transcriptional dynamics during mouse spermatogenesis." *Nat Commun* 10:1–20 .
<https://doi.org/10.1038/s41467-019-09182-1>
- Fayomi AP, Orwig KE (2018). "Spermatogonial stem cells and spermatogenesis in mice, monkeys and men."

- Stem Cell Res* 29:207–214 . <https://doi.org/10.1016/j.scr.2018.04.009>
- Ferlin A, Raicu F, Gatta V, Zuccarello D, Palka G, Foresta C (2007). "Male infertility: role of genetic background." *Reprod Biomed Online* 14:734–745
- Fraune J, Alsheimer M, Redolfi J, Brochier-Armanet C, Benavente R (2015). "Protein SYCP2 Is an Ancient Component of the Metazoan Synaptonemal Complex." *Cytogenet Genome Res* 144:299–305 . <https://doi.org/10.1159/000381080>
- Geisinger A, Benavente R (2017). "Mutations in Genes Coding for Synaptonemal Complex Proteins and Their Impact on Human Fertility." *Cytogenet. Genome Res.* 150:77–85
- Gershoni M, Pietrokovski S (2014). "Reduced selection and accumulation of deleterious mutations in genes exclusively expressed in men." *Nat Commun* 5:1–10 . <https://doi.org/10.1038/ncomms5438>
- Gillies CB (1975). "Synaptonemal complex and chromosome structure." *Annu. Rev. Genet.* 9:91–109
- Golezar S, Ramezani Tehrani F, Khazaei S, Ebadi A, Keshavarz Z (2019). "The global prevalence of primary ovarian insufficiency and early menopause: a meta-analysis." *Climacteric* 22:403–411 . <https://doi.org/10.1080/13697137.2019.1574738>
- Gomez-H L, Felipe-Medina N, Sanchez-Martin M, Davies OR, Ramos I, Garcia-Tunon I, de Rooij DG, Dereli I, Toth A, Barbero JL, Benavente R, Llano E, Pendas AM (2016). "C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility." *Nat Commun* 7:13298 . <https://doi.org/10.1038/ncomms13298>
- Gomez-H L, Felipe-Medina1 N, Condezo YB, Garcia-Valiente R, Ramos I, Suja JA, Barbero JL, Roig I, Sanchez-Martin M, De Rooij DG, Llano E, Pendasl AM (2019). "The PSMA8 subunit of the spermatoproteasome is essential for proper meiotic exit and mouse fertility." *PLoS Genet* 15:e1008316 . <https://doi.org/10.1371/journal.pgen.1008316>
- Gómez R, Valdeolmillos A, Parra MT, Viera A, Carreiro C, Roncal F, Rufas JS, Barbero JL, Suja JA (2007). "Mammalian SGO2 appears at the inner centromere domain and redistributes depending on tension across centromeres during meiosis II and mitosis." *EMBO Rep* 8:173–180 . <https://doi.org/10.1038/sj.embor.7400877>
- Grangeia A, Sá R, Carvalho F, Martin J, Girodon E, Silva J, Ferráz L, Barros A, Sousa M (2007). "Molecular characterization of the cystic fibrosis transmembrane conductance regulator gene in congenital absence of the vas deferens." *Genet Med* 9:163–172 . <https://doi.org/10.1097/GIM.0b013e3180318aaf>
- Grey C, Barthès P, Chauveau-Le Friec G, Langa F, Baudat F, de Massy B (2011). "Mouse PRDM9 DNA-Binding Specificity Determines Sites of Histone H3 Lysine 4 Trimethylation for Initiation of Meiotic Recombination." *PLoS Biol* 9:e1001176 . <https://doi.org/10.1371/journal.pbio.1001176>
- Griswold MD (2016). "Spermatogenesis: The Commitment to Meiosis." *Physiol Rev* 96:1–17 . <https://doi.org/10.1152/physrev.00013.2015>
- Griswold MD, Hogarth CA, Bowles J, Koopman P (2012). "Initiating meiosis: The case for retinoic acid." *Biol. Reprod.* 86:35–36
- Gruhn JR, Rubio C, Broman KW, Hunt PA, Hassold T (2013). "Cytological Studies of Human Meiosis: Sex-Specific Differences in Recombination Originate at, or Prior to, Establishment of Double-Strand Breaks." *PLoS One* 8:e85075 . <https://doi.org/10.1371/journal.pone.0085075>
- Guillon H, Baudat F, Grey C, Liskay RM, de Massy B (2005). "Crossover and noncrossover pathways in mouse meiosis." *Mol Cell* 20:563–573 . <https://doi.org/10.1016/j.molcel.2005.09.021>
- Guo K, Li C-H, Wang X-Y, He D-J, Zheng P (2016). "Germ stem cells are active in postnatal mouse ovary under physiological conditions." *Mol Hum Reprod* 22:316–328 . <https://doi.org/10.1093/molehr/gaw015>
- Guo T, Zhao S, Zhao S, Chen M, Li G, Jiao X, Wang Z, Zhao Y, Qin Y, Gao F, Chen ZJ (2017). "Mutations in MSH5 in primary ovarian insufficiency." *Hum Mol Genet* 26:1452–1457 . <https://doi.org/10.1093/hmg/ddx044>
- Gurunath S, Pandian Z, Anderson RA, Bhattacharya S (2011). "Defining infertility-a systematic review of prevalence studies." *Hum. Reprod. Update* 17:575–588
- Hackett JA, Kobayashi T, Dietmann S, Surani MA (2017). "Activation of Lineage Regulators and

- Transposable Elements across a Pluripotent Spectrum." *Stem Cell Reports* 8:1645–1658 .
<https://doi.org/10.1016/j.stemcr.2017.05.014>
- Halldorsson B V., Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP, Gunnarsson B, Oddsson A, Halldorsson GH, Zink F, Gudjonsson SA, Frigge ML, Thorleifsson G, Sigurdsson A, Stacey SN, Sulem P, Masson G, Helgason A, Gudbjartsson DF, Thorsteinsdottir U, Stefansson K (2019). "Human genetics: Characterizing mutagenic effects of recombination through a sequence-level genetic map." *Science* (80-) 363: . <https://doi.org/10.1126/science.aau1043>
- Hamada AJ, Esteves SC, Agarwal A (2013). "A comprehensive review of genetics and genetic testing in azoospermia." *Clin (Sao Paulo)* 68 Suppl 1:39–60
- Hamer G, Gell K, Kouznetsova A, Novak I, Benavente R, Höög C (2006). "Characterization of a novel meiosis-specific protein within the central element of the synaptonemal complex." *J Cell Sci* 119:4025–4032 . <https://doi.org/10.1242/jcs.03182>
- Harper JC, Aittomäki K, Borry P, Cornel MC, De Wert G, Dondorp W, Geraedts J, Gianaroli L, Ketterson K, Liebaers I, Lundin K, Mertes H, Morris M, Pennings G, Sermon K, Spits C, Soini S, Van Montfort APA, Veiga A, Vermeesch JR, Viville S, Macek M (2018). "Recent developments in genetics and medically assisted reproduction: From research to clinical applications." *Eur. J. Hum. Genet.* 26:12–33
- Harris SE, Chand AL, Winship IM, Gersak K, Aittomäki K, Shelling AN (2002). "Identification of novel mutations in FOXL2 associated with premature ovarian failure." *Mol Hum Reprod* 8:729–733 .
<https://doi.org/10.1093/molehr/8.8.729>
- Hassold T, Hall H, Hunt P (2007). "The origin of human aneuploidy: Where we have been, where we are going." *Hum. Mol. Genet.* 16
- Hassold T, Hunt P (2001). "To err (meiotically) is human: The genesis of human aneuploidy." *Nat. Rev. Genet.* 2:280–291
- Hayashi K, Yoshida K, Matsui Y (2005). "A histone H3 methyltransferase controls epigenetic events required for meiotic prophase." *Nature* 438:374–378 . <https://doi.org/10.1038/nature04112>
- He W Bin, Tu CF, Liu Q, Meng LL, Yuan SM, Luo AX, He FS, Shen J, Li W, Du J, Zhong CG, Lu GX, Lin G, Fan LQ, Tan YQ (2018). "DMC1 mutation that causes human non-obstructive azoospermia and premature ovarian insufficiency identified by whole-exome sequencing." *J Med Genet* 55:198–204 .
<https://doi.org/10.1136/jmedgenet-2017-104992>
- He M, Zhou Z, Shah AA, Zou H, Tao J, Chen Q, Wan Y (2016). "The emerging role of deubiquitinating enzymes in genomic integrity, diseases, and therapeutics." *Cell Biosci* 6:62 .
<https://doi.org/10.1186/s13578-016-0127-1>
- Hellmuth S, Pöhlmann C, Brown A, Böttger F, Sprinzl M, Stemmann O (2015). "Positive and negative regulation of vertebrate separase by Cdk1-cyclin B1 may explain why securin is dispensable." *J Biol Chem* 290:8002–8010 . <https://doi.org/10.1074/jbc.M114.615310>
- Herbert M, Levasseur M, Homer H, Yallop K, Murdoch A, McDougall A (2003). "Homologue disjunction in mouse oocytes requires proteolysis of securin and cyclin B1." *Nat Cell Biol* 5:1023–1025 .
<https://doi.org/10.1038/ncb1062>
- Herrán Y, Gutiérrez-Caballero C, Sánchez-Martín M, Hernández T, Viera A, Barbero JL, De Álava E, De Rooij DG, Suja JA, Llano E, Pendás AM (2011). "The cohesin subunit RAD21L functions in meiotic synapsis and exhibits sexual dimorphism in fertility." *EMBO J* 30:3091–3105 .
<https://doi.org/10.1038/emboj.2011.222>
- Heyting C (1996). "Synaptonemal complexes: Structure and function." *Curr Opin Cell Biol* 8:389–396 .
[https://doi.org/10.1016/S0955-0674\(96\)80015-9](https://doi.org/10.1016/S0955-0674(96)80015-9)
- Hinch AG, Becker PW, Li T, Moralli D, Zhang G, Bycroft C, Green C, Keeney S, Shi Q, Davies B, Donnelly P (2020). "The Configuration of RPA, RAD51, and DMC1 Binding in Meiosis Reveals the Nature of Critical Recombination Intermediates." *Mol Cell* 79:689–701.e10 .
<https://doi.org/10.1016/j.molcel.2020.06.015>
- Holloway JK, Booth J, Edelmann W, McGowan CH, Cohen PE (2008). "MUS81 generates a subset of MLH1-MLH3-independent crossovers in mammalian meiosis." *PLoS Genet* 4: .
<https://doi.org/10.1371/journal.pgen.1000186>

- Hu Z, Xia Y, Guo X, Dai J, Li H, Hu H, Jiang Y, Lu F, Wu Y, Yang X, Li H, Yao B, Lu C, Xiong C, Li Z, Gui Y, Liu J, Zhou Z, Shen H, Wang X, Sha J (2011). "A genome-wide association study in Chinese men identifies three risk loci for non-obstructive azoospermia." *Nat Genet* 44:183–186 .
<https://doi.org/10.1038/ng.1040>
- Hunt PA, Hassold TJ (2002). "Sex matters in meiosis." *Science* (80-.). 296:2181–2183
- Irie N, Weinberger L, Tang WWC, Kobayashi T, Viukov S, Manor YS, Dietmann S, Hanna JH, Surani MA (2015). "SOX17 is a critical specifier of human primordial germ cell fate." *Cell* 160:253–268 .
<https://doi.org/10.1016/j.cell.2014.12.013>
- Isaksson R, Tiitinen A (2004). "Present concept of unexplained infertility." *Gynecol. Endocrinol.* 18:278–290
- Ishii T (2014). "Human iPS Cell-Derived Germ Cells: Current Status and Clinical Potential." *J Clin Med* 3:1064–1083 . <https://doi.org/10.3390/jcm3041064>
- Jagarlamudi K, Rajkovic A (2012). "Oogenesis: Transcriptional regulators and mouse models." *Mol. Cell. Endocrinol.* 356:31–39
- Jaroszeski MJ, Heller R, Gilbert R, Muramatsu T (2003). "In Vivo Gene Electroporation in the Mouse Testis." In: *Electrochemotherapy, Electrogenethapy, and Transdermal Drug Delivery*. Humana Press, pp 349–357
- Jauregui EJ, Mitchell D, Topping T, Hogarth CA, Griswold MD (2018). "Retinoic acid receptor signaling is necessary in steroidogenic cells for normal spermatogenesis and epididymal function." *Dev* 145: .
<https://doi.org/10.1242/dev.160465>
- Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012). "A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity." *Science* (80-) 337:816–821 .
<https://doi.org/10.1126/science.1225829>
- Johnson J, Canning J, Kaneko T, Pru JK, Tilly JL (2004). "Germline stem cells and follicular renewal in the postnatal mammalian ovary." *Nature* 428:145–150 . <https://doi.org/10.1038/nature02316>
- Jose-Miller AB, Boyden JW, Frey KA (2007). "Infertility"
- Kallio M, Chang Y, Manuel M, Alastalo TP, Rallu M, Gitton Y, Pirkkala L, Loones MT, Paslaru L, Larney S, Hiard S, Morange M, Sistonen L, Mezger V (2002). "Brain abnormalities, defective meiotic chromosome synapsis and female subfertility in HSF2 null mice." *EMBO J* 21:2591–2601 .
<https://doi.org/10.1093/emboj/21.11.2591>
- Kauppi L, Barchi M, Baudat F, Romanienko PJ, Keeney S, Jasin M (2011). "Distinct properties of the XY pseudoautosomal region crucial for male meiosis." *Science* (80-) 331:916–920 .
<https://doi.org/10.1126/science.1195774>
- Kauppi L, Barchi M, Lange J, Baudat F, Jasin M, Keeney S (2013). "Numerical constraints and feedback control of double-strand breaks in mouse meiosis." *Genes Dev* 27:873–886 .
<https://doi.org/10.1101/gad.213652.113>
- Keeney S, Giroux CN, Kleckner N (1997). "Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family." *Cell* 88:375–384 .
[https://doi.org/10.1016/S0092-8674\(00\)81876-0](https://doi.org/10.1016/S0092-8674(00)81876-0)
- Khalil AM, Boyar FZ, Driscoll DJ (2004). "Dynamic histone modifications mark sex chromosome inactivation and reactivation during mammalian spermatogenesis." *Proc Natl Acad Sci U S A* 101:16583–16587 .
<https://doi.org/10.1073/pnas.0406325101>
- Khil PP, Smagulova F, Brick KM, Camerini-Otero RD, Petukhova G V. (2012). "Sensitive mapping of recombination hotspots using sequencing-based detection of ssDNA." *Genome Res* 22:957–965 .
<https://doi.org/10.1101/gr.130583.111>
- Kishi K, Uchida A, Takase HM, Suzuki H, Kurohmaru M, Tsunekawa N, Kanai-Azuma M, Wood SA, Kanai Y (2017). "Spermatogonial deubiquitinase USP9X is essential for proper spermatogenesis in mice." *Reproduction* 154:135–143 . <https://doi.org/10.1530/REP-17-0184>
- Kitajima TS, Miyazaki Y, Yamamoto M, Watanabe Y (2003). "Rec8 cleavage by separase is required for meiotic nuclear divisions in fission yeast." *EMBO J* 22:5643–5653 .
<https://doi.org/10.1093/emboj/cdg527>
- Kitajima TS, Sakuno T, Ishiguro KI, Iemura SI, Natsume T, Kawashima SA, Watanabe Y (2006). "Shugoshin

- collaborates with protein phosphatase 2A to protect cohesin." *Nature* 441:46–52 .
<https://doi.org/10.1038/nature04663>
- Kobayashi T, Surani MA (2018). "On the origin of the human germline." *Dev* 145: .
<https://doi.org/10.1242/dev.150433>
- Komander D, Clague MJ, Urbe S (2009). "Breaking the chains: structure and function of the deubiquitinases." *Nat Rev Mol Cell Biol* 10:550–563 . <https://doi.org/10.1038/nrm2731>
- Koubova J, Hu YC, Bhattacharyya T, Soh YQS, Gill ME, Goodheart ML, Hogarth CA, Griswold MD, Page DC (2014). "Retinoic Acid Activates Two Pathways Required for Meiosis in Mice." *PLoS Genet* 10: .
<https://doi.org/10.1371/journal.pgen.1004541>
- Koubova J, Menke DB, Zhou Q, Cape B, Griswold MD, Page DC (2006). "Retinoic acid regulates sex-specific timing of meiotic initiation in mice." *Proc Natl Acad Sci U S A* 103:2474–2479 .
<https://doi.org/10.1073/pnas.0510813103>
- Krausz C, Riera-Escamilla A (2018). "Genetics of male infertility." *Nat. Rev. Urol.* 15:369–384
- Kudo NR, Anger M, Peters AHFM, Stemmann O, Theussl HC, Helmhart W, Kudo H, Heyting C, Nasmyth K (2009). "Role of cleavage by separase of the Rec8 kleisin subunit of cohesin during mammalian meiosis I." *J Cell Sci* 122:2686–2698 . <https://doi.org/10.1242/jcs.035287>
- Kumar R, Bourbon HM, De Massy B (2010). "Functional conservation of Mei4 for meiotic DNA double-strand break formation from yeasts to mice." *Genes Dev* 24:1266–1280 .
<https://doi.org/10.1101/gad.571710>
- Kumar R, Ghyselinck N, Ishiguro K ichiro, Watanabe Y, Kouznetsova A, Höög C, Strong E, Schimenti J, Daniel K, Toth A, de Massy B (2015). "MEI4 - a central player in the regulation of meiotic DNA double-strand break formation in the mouse." *J Cell Sci* 128:1800–1811 . <https://doi.org/10.1242/jcs.165464>
- Kumar S, Chatzi C, Brade T, Cunningham TJ, Zhao X, Duester G (2011). "Sex-specific timing of meiotic initiation is regulated by Cyp26b1 independent of retinoic acid signalling." *Nat Commun* 2:1–8 .
<https://doi.org/10.1038/ncomms1136>
- Kwon J, Mochida K, Wang YL, Sekiguchi S, Sankai T, Aoki S, Ogura A, Yoshikawa Y, Wada K (2005). "Ubiquitin C-terminal hydrolase L-1 is essential for the early apoptotic wave of germinal cells and for sperm quality control during spermatogenesis." *Biol Reprod* 73:29–35 .
<https://doi.org/10.1095/biolreprod.104.037077>
- La Salle S, Palmer K, O'Brien M, Schimenti JC, Eppig J, Handel MA (2012). "Spata22, a novel vertebrate-specific gene, is required for meiotic progress in mouse: Germ cells." *Biol Reprod* 86: .
<https://doi.org/10.1095/biolreprod.111.095752>
- Laemmli UK (1970). "Cleavage of structural proteins during the assembly of the head of bacteriophage T4." *Nature* 227:680–685 . <https://doi.org/10.1038/227680a0>
- Lake CM, Hawley RS (2013). "RNF212 marks the spot." *Nat Genet* 45:228–229 .
<https://doi.org/10.1038/ng.2559>
- Lammers JH, Offenberger HH, van Aalderen M, Vink AC, Dietrich AJ, Heyting C (1994). "The gene encoding a major component of the lateral elements of synaptonemal complexes of the rat is related to X-linked lymphocyte-regulated genes." *Mol Cell Biol* 14:1137–1146 . <https://doi.org/10.1128/mcb.14.2.1137>
- Lange J, Yamada S, Tischfield SE, Pan J, Kim S, Zhu X, Socci ND, Jasin M, Keeney S (2016). "The Landscape of Mouse Meiotic Double-Strand Break Formation, Processing, and Repair." *Cell* 167:695–708.e16 .
<https://doi.org/10.1016/j.cell.2016.09.035>
- Lawson KA, Dunn NR, Roelen BAJ, Zeinstra LM, Davis AM, Wright CVE, Korving JPWFM, Hogan BLM (1999). "Bmp4 is required for the generation of primordial germ cells in the mouse embryo." *Genes Dev* 13:424–436 . <https://doi.org/10.1101/gad.13.4.424>
- Lee CY, Horn HF, Stewart CL, Burke B, Bolcun-Filas E, Schimenti JC, Dresser ME, Pezza RJ (2015). "Mechanism and Regulation of Rapid Telomere Prophase Movements in Mouse Meiotic Chromosomes." *Cell Rep* 11:551–563 . <https://doi.org/10.1016/j.celrep.2015.03.045>
- Lee IW, Kuan LC, Lin CH, Pan HA, Hsu CC, Tsai YC, Kuo PL, Teng YN (2008a). "Association of USP26 haplotypes in men in Taiwan, China with severe spermatogenic defect." *Asian J Androl* 10:896–904 .
<https://doi.org/10.1111/j.1745-7262.2008.00439.x>

- Lee J, Kitajima TS, Tanno Y, Yoshida K, Morita T, Miyano T, Miyake M, Watanabe Y (2008b). "Unified mode of centromeric protection by shugoshin in mammalian oocytes and somatic cells." *Nat Cell Biol* 10:42–52 . <https://doi.org/10.1038/ncb1667>
- Lei L, Spradling AC (2013). "Female mice lack adult germ-line stem cells but sustain oogenesis using stable primordial follicles." *Proc Natl Acad Sci U S A* 110:8585–8590 . <https://doi.org/10.1073/pnas.1306189110>
- Li L, Xi Q, Zhang H, Fei J, Jiang Y, Li L, Liu R, Wang R, Zhang G (2020a). "A novel frameshift mutation in ubiquitin-specific protease 26 gene in a patient with severe oligozoospermia." *Biosci Rep* 40: . <https://doi.org/10.1042/BSR20191902>
- Li M, Feng H, Lin Z, Zheng J, Liu D, Guo R, Li J, Li RHW, Ng EHY, Huen MSY, Wang PJ, Yeung WSB, Liu K (2020b). "The Novel Male Meiosis Recombination Regulator (MAMERR) Coordinates the Progression of Meiosis Prophase I." *J Genet Genomics*. <https://doi.org/10.1016/j.jgg.2020.08.001>
- Li Z, Huang Y, Li H, Hu J, Liu X, Jiang T, Sun G, Tang A, Sun X, Qian W, Zeng Y, Xie J, Zhao W, Xu Y, He T, Dong C, Liu Q, Mou L, Lu J, Lin Z, Wu S, Gao S, Guo G, Feng Q, Li Y, Zhang X, Wang J, Yang H, Wang J, Xiong C, Cai Z, Gui Y (2015). "Excess of rare variants in genes that are key epigenetic regulators of spermatogenesis in the patients with non-obstructive azoospermia." *Sci Rep* 5:8785 . <https://doi.org/10.1038/srep08785>
- Lin YW, Hsu TH, Yen PH (2011). "Localization of ubiquitin specific protease 26 at blood-testis barrier and near Sertoli cell-germ cell interface in mouse testes." *Int J Androl* 34:e368-77 . <https://doi.org/10.1111/j.1365-2605.2010.01130.x>
- Lindsay TJ (2015). "Evaluation and Treatment of Infertility"
- Lipkin SM, Moens PB, Wang V, Lenzi M, Shanmugarajah D, Gilgeous A, Thomas J, Cheng J, Touchman JW, Green ED, Schwartzberg P, Collins FS, Cohen PE (2002). "Meiotic arrest and aneuploidy in MLH3-deficient mice." *Nat Genet* 31:385–390 . <https://doi.org/10.1038/ng931>
- Liu YL, Zheng J, Mi YJ, Zhao J, Tian QB (2018). "The impacts of nineteen mutations on the enzymatic activity of USP26." *Gene* 641:292–296 . <https://doi.org/10.1016/j.gene.2017.10.074>
- Llano E, Gómez R, Gutiérrez-Caballero C, Herrán Y, Sánchez-Martín M, Vázquez-Quñones L, Hernández T, De Álava E, Cuadrado A, Barbero JL, Suja JA, Pendás AM (2008). "Shugoshin-2 is essential for the completion of meiosis but not for mitotic cell division in mice." *Genes Dev* 22:2400–2413 . <https://doi.org/10.1101/gad.475308>
- Llano E, Herrán Y, García-Tuñón I, Gutiérrez-Caballero C, de Álava E, Barbero JL, Schimenti J, de Rooij DG, Sánchez-Martín M, Pendás AM (2012). "Meiotic cohesin complexes are essential for the formation of the axial element in mice." *J Cell Biol* 197:877–885 . <https://doi.org/10.1083/jcb.201201100>
- Loregian A, Appleton BA, Hogle JM, Coen DM (2004). "Specific Residues in the Connector Loop of the Human Cytomegalovirus DNA Polymerase Accessory Protein UL44 Are Crucial for Interaction with the UL54 Catalytic Subunit." *J Virol* 78:9084–9092 . <https://doi.org/10.1128/jvi.78.17.9084-9092.2004>
- Luddi A, Crifasi L, Quagliarello A, Governini L, De Leo V, Piomboni P (2016). "Single nucleotide polymorphisms of USP26 in azoospermic men." *Syst Biol Reprod Med* 62:372–378 . <https://doi.org/10.1080/19396368.2016.1238116>
- Luo M, Yang F, Leu NA, Landaiche J, Handel MA, Benavente R, La Salle S, Wang PJ (2013). "MEIOB exhibits single-stranded DNA-binding and exonuclease activities and is essential for meiotic recombination." *Nat Commun* 4:2788 . <https://doi.org/10.1038/ncomms3788>
- Luo M, Zhou J, Leu NA, Abreu CM, Wang J, Anguera MC, de Rooij DG, Jasin M, Wang PJ (2015). "Polycomb protein SCML2 associates with USP7 and counteracts histone H2A ubiquitination in the XY chromatin during male meiosis." *PLoS Genet* 11:e1004954 . <https://doi.org/10.1371/journal.pgen.1004954>
- Ma J, Zeng F, Schultz RM, Tseng H (2006). "Basonuclin: A novel mammalian maternal-effect gene." *Development* 133:2053–2062 . <https://doi.org/10.1242/dev.02371>
- Ma Q, Li Y, Guo H, Li C, Chen J, Luo M, Jiang Z, Li H, Gui Y (2016). "A Novel Missense Mutation in USP26 Gene Is Associated With Nonobstructive Azoospermia." *Reprod Sci* 23:1434–1441 . <https://doi.org/10.1177/1933719116641758>
- Mahadevaiah SK, Turner JMA, Baudat F, Rogakou EP, De Boer P, Blanco-Rodríguez J, Jasin M, Keeney S,

- Bonner WM, Burgoyne PS (2001). "Recombinational DNA double-strand breaks in mice precede synapsis." *Nat Genet* 27:271–276 . <https://doi.org/10.1038/85830>
- Makino S, Fukumura R, Gondo Y (2016). "Illegitimate translation causes unexpected gene expression from on-target out-of-frame alleles created by CRISPR-Cas9." *Sci Rep* 6:39608 . <https://doi.org/10.1038/srep39608>
- Malhas A, Goulbourne C, Vaux DJ (2011). "The nucleoplasmic reticulum: Form and function." *Trends Cell Biol.* 21:362–373
- Mallepally R, Butler PR, Herati AS, Lamb DJ (2017). "Genetic Basis of Male and Female Infertility." *Monogr Hum Genet* 21:1–16 . <https://doi.org/10.1159/000477275>
- Margolin G, Khil PP, Kim J, Bellani MA, Camerini-Otero RD (2014). "Integrated transcriptome analysis of mouse spermatogenesis." *BMC Genomics* 15:39 . <https://doi.org/10.1186/1471-2164-15-39>
- Marston AL (2015). "Shugoshins: Tension-Sensitive Pericentromeric Adaptors Safeguarding Chromosome Segregation." *Mol Cell Biol* 35:634–648 . <https://doi.org/10.1128/mcb.01176-14>
- Martin RH (2008). "Cytogenetic determinants of male fertility." *Hum Reprod Update* 14:379–390 . <https://doi.org/10.1093/humupd/dmn017>
- Martinez JS, Von Nicolai C, Kim T, Ehlén Å, Mazin A V., Kowalczykowski SC, Carreira A (2016). "BRCA2 regulates DMC1-mediated recombination through the BRC repeats." *Proc Natl Acad Sci U S A* 113:3515–3520 . <https://doi.org/10.1073/pnas.1601691113>
- Matzuk MM, Lamb DJ (2008). "The biology of infertility: research advances and clinical challenges." *Nat Med* 14:1197–1213 . <https://doi.org/10.1038/nm.f.1895>
- McGee EA, Hsu SY, Kaipia A, J.w. Hsueh A (1998). "Cell death and survival during ovarian follicle development." In: *Molecular and Cellular Endocrinology. Mol Cell Endocrinol*, pp 15–18
- McGee EA, Hsueh AJW (2000). "Initial and Cyclic Recruitment of Ovarian Follicles 1 ." *Endocr Rev* 21:200–214 . <https://doi.org/10.1210/edrv.21.2.0394>
- McGuire MM, Bowden W, Engel NJ, Ahn HW, Kovanci E, Rajkovic A (2011). "Genomic analysis using high-resolution single-nucleotide polymorphism arrays reveals novel microdeletions associated with premature ovarian failure." *Fertil Steril* 95:1595–1600 . <https://doi.org/10.1016/j.fertnstert.2010.12.052>
- McLaren A (1998). "Gonad development: Assembling the mammalian testis." *Curr Biol* 8:175–177
- McLaren A, Southee D (1997). "Entry of mouse embryonic germ cells into meiosis." *Dev Biol* 187:107–113 . <https://doi.org/10.1006/dbio.1997.8584>
- McMurray CT (2010). "Mechanisms of trinucleotide repeat instability during human development." *Nat. Rev. Genet.* 11:786–799
- Meuwissen RLJ, Offenberger HH, Dietrich AJJ, Riesewijk A, Van Iersel M, Heyting C (1992). "A coiled-coil related protein specific for synapsed regions of meiotic prophase chromosomes." *EMBO J* 11:5091–5100 . <https://doi.org/10.1002/j.1460-2075.1992.tb05616.x>
- Mitsunaga S, Shioda T (2018). "Evolutionarily diverse mechanisms of germline specification among mammals: what about us?" *Stem Cell Investig.* 5
- Miyawaki S, Kuroki S, Maeda R, Okashita N, Koopman P, Tachibana M (2020). "The mouse Sry locus harbors a cryptic exon that is essential for male sex determination." *Science* 370:121–124 . <https://doi.org/10.1126/science.abb6430>
- Miyazaki WY, Orr-Weaver TL (1994). "Sister-chromatid cohesion in mitosis and meiosis." *Annu. Rev. Genet.* 28:167–187
- Moens PB, KNK, TMMECPE. SB (2002). "The time course and chromosomal localization of recombination-related proteins at meiosis in the mouse are compatible with models that can resolve the early DNA-DNA interactions without reciprocal recombination | *Journal of Cell Science.*" *J Cell Sci* 115:1611–1622
- Morelli MA, Cohen PE (2005). "Not all germ cells are created equal: Aspects of sexual dimorphism in mammalian meiosis." *Reproduction* 130:761–781
- Moyes-Oliveira M, Guilherme RDS, Dantas AG, Ueta R, Perez AB, Haidar M, Canonaco R, Meloni VA,

- Kosyakova N, Liehr T, Carnevali GM, Melaragno MI (2015). "Genetic mechanisms leading to primary amenorrhea in balanced X-autosome translocations." *Fertil Steril* 103:1289-1296.e2 . <https://doi.org/10.1016/j.fertnstert.2015.01.030>
- Muciaccia B, Boitani C, Berloco BP, Nudo F, Spadetta G, Stefanini M, de Rooij DG, Vicini E (2013). "Novel Stage Classification of Human Spermatogenesis Based on Acrosome Development1." *Biol Reprod* 89:1–10 . <https://doi.org/10.1095/biolreprod.113.111682>
- Muratori M, Marchiani S, Forti G, Baldi E (2005). "Sperm ubiquitination positively correlates to normal morphology in human semen." *Hum Reprod* 20:1035–1043 . <https://doi.org/10.1093/humrep/deh678>
- Myers S, Bowden R, Tumian A, Bontrop RE, Freeman C, MacFie TS, McVean G, Donnelly P (2010). "Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic recombination." *Science* (80-) 327:876–879 . <https://doi.org/10.1126/science.1182363>
- Neale MJ, Pan J, Keeney S (2005). "Endonucleolytic processing of covalent protein-linked DNA double-strand breaks." *Nature* 436:1053–1057 . <https://doi.org/10.1038/nature03872>
- Nelson LM (2009). "Primary Ovarian Insufficiency." *N Engl J Med* 360:606–614 . <https://doi.org/10.1056/NEJMcp0808697>
- Niikura Y, Niikura T, Tilly JL (2009). "Aged mouse ovaries possess rare premeiotic germ cells that can generate oocytes following transplantation into a young host environment." *Aging (Albany NY)* 1:971–978 . <https://doi.org/10.18632/aging.100105>
- Nijman SM, Luna-Vargas MP, Velds A, Brummelkamp TR, Dirac AM, Sixma TK, Bernards R (2005). "A genomic and functional inventory of deubiquitinating enzymes." *Cell* 123:773–786 . <https://doi.org/10.1016/j.cell.2005.11.007>
- Nikolic A, Volarevic V, Armstrong L, Lako M, Stojkovic M (2016). "Primordial Germ Cells: Current Knowledge and Perspectives." *Stem Cells Int* 2016: . <https://doi.org/10.1155/2016/1741072>
- Nishimune Y, Tanaka H (2006). "Infertility caused by polymorphisms or mutations in spermatogenesis-specific genes." *J Androl* 27:326–334 . <https://doi.org/10.2164/jandrol.05162>
- Novak I, Wang H, Revenkova E, Jessberger R, Scherthan H, Höög C (2008). "Cohesin Smc1 β determines meiotic chromatin axis loop organization." *J Cell Biol* 180:83–90 . <https://doi.org/10.1083/jcb.200706136>
- O'Flynn O'Brien KL, Varghese AC, Agarwal A (2010). "The genetic causes of male factor infertility: A review." *Fertil Steril* 93:1–12 . <https://doi.org/10.1016/j.fertnstert.2009.10.045>
- Oakberg EF (1956). "Duration of spermatogenesis in the mouse and timing of stages of the cycle of the seminiferous epithelium." *Am J Anat* 99:507–516 . <https://doi.org/10.1002/aja.1000990307>
- Oatley JM, Brinster RL (2008). "Regulation of spermatogonial stem cell self-renewal in mammals." *Annu. Rev. Cell Dev. Biol.* 24:263–286
- Offenberg HH, Schalk JAC, Meuwissen RLJ, Van Aalderen M, Kester HA, Dietrich AJJ, Heyting C (1998). "SCP2: a major protein component of the axial elements of synaptonemal complexes of the rat"
- Ohinata Y, Ohta H, Shigeta M, Yamanaka K, Wakayama T, Saitou M (2009). "A Signaling Principle for the Specification of the Germ Cell Lineage in Mice." *Cell* 137:571–584 . <https://doi.org/10.1016/j.cell.2009.03.014>
- Ohinata Y, Payer B, O'Carroll D, Ancelin K, Ono Y, Sano M, Barton SC, Obukhanych T, Nussenzweig M, Tarakhovsky A, Saitou M, Surani MA (2005). "Blimp1 is a critical determinant of the germ cell lineage in mice." *Nature* 436:207–213 . <https://doi.org/10.1038/nature03813>
- Paduch DA, Mielnik A, Schlegel PN (2005). "Novel mutations in testis-specific ubiquitin protease 26 gene may cause male infertility and hypogonadism." *Reprod Biomed Online* 10:747–754
- Palmer N, Talib SZA, Singh P, Goh CMF, Liu K, Schimenti JC, Kaldis P (2020). "A novel function for CDK2 activity at meiotic crossover sites." *PLoS Biol* 18:e3000903 . <https://doi.org/10.1371/journal.pbio.3000903>
- Parra MT, Page J, Yen TJ, He D, Valdeolmillos A, Rufas JS, Suja JA (2002). "Expression and behaviour of CENP-E at kinetochores during mouse spermatogenesis." *Chromosoma* 111:53–61 . <https://doi.org/10.1007/s00412-002-0185-5>

- Parvanov ED, Petkov PM, Paigen K (2010). "Prdm9 controls activation of mammalian recombination hotspots." *Science* (80-.). 327:835
- Pastore LM, Johnson J (2014). "The FMR1 gene, infertility, and reproductive decision-making: A review." *Front. Genet.* 5
- Pedersen T, Peters H (1968). "Proposal for a classification of oocytes and follicles in the mouse ovary." *J Reprod Fertil* 17:555–557 . <https://doi.org/10.1530/jrf.0.0170555>
- Pepling ME, Spradling AC (2001). "Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles." *Dev Biol* 234:339–351 . <https://doi.org/10.1006/dbio.2001.0269>
- Peters AH, Plug AW, van Vugt MJ, de Boer P (1997). "A drying-down technique for the spreading of mammalian meiocytes from the male and female germline." *Chromosom Res* 5:66–68
- Peters H (1969). "The development of the mouse ovary from birth to maturity." *Acta Endocrinol (Copenh)* 62:98–116 . <https://doi.org/10.1530/acta.0.0620098>
- Petronczki M, Siomos MF, Nasmyth K (2003). "Un ménage à quatre: The molecular biology of chromosome segregation in meiosis." *Cell* 112:423–440
- Pittman DL, Cobb J, Schimenti KJ, Wilson LA, Cooper DM, Brignull E, Handel MA, Schimenti JC (1998). "Meiotic prophase arrest with failure of chromosome synapsis in mice deficient for Dmc1, a germline-specific RecA homolog." *Mol Cell* 1:697–705 . [https://doi.org/10.1016/S1097-2765\(00\)80069-6](https://doi.org/10.1016/S1097-2765(00)80069-6)
- Powers NR, Parvanov ED, Baker CL, Walker M, Petkov PM, Paigen K (2016). "The Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at Recombination Hotspots In Vivo." *PLoS Genet* 12: . <https://doi.org/10.1371/journal.pgen.1006146>
- Qiao H, Prasada Rao HBD, Yang Y, Fong JH, Cloutier JM, Deacon DC, Nagel KE, Swartz RK, Strong E, Holloway JK, Cohen PE, Schimenti J, Ward J, Hunter N (2014). "Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination." *Nat Genet* 46:194–199 . <https://doi.org/10.1038/ng.2858>
- Rao HB, Qiao H, Bhatt SK, Bailey LR, Tran HD, Bourne SL, Qiu W, Deshpande A, Sharma AN, Beebout CJ, Pezza RJ, Hunter N (2017). "A SUMO-ubiquitin relay recruits proteasomes to chromosome axes to regulate meiotic recombination." *Science* (80-) 355:403–407 . <https://doi.org/10.1126/science.aaf6407>
- Ravel C, El Houate B, Chantot S, Lourenco D, Dumaine A, Rouba H, Bandyopadhyay A, Radhakrishna U, Das B, Sengupta S, Mandelbaum J, Siffroi JP, McElreavey K (2006). "Haplotypes, mutations and male fertility: the story of the testis-specific ubiquitin protease USP26." *Mol Hum Reprod* 12:643–646 . <https://doi.org/10.1093/molehr/gal063>
- Raverdeau M, Gely-Pernot A, Féret B, Dennefeld C, Benoit G, Davidson I, Chambon P, Marka M, Ghyselinck NB (2012). "Retinoic acid induces Sertoli cell paracrine signals for spermatogonia differentiation but cell autonomously drives spermatocyte meiosis." *Proc Natl Acad Sci U S A* 109:16582–16587 . <https://doi.org/10.1073/pnas.1214936109>
- Reijo R, Alagappan RK, Patrizio P, Page DC (1996). "Severe oligozoospermia resulting from deletions of azoospermia factor gene on Y chromosome." *Lancet* 347:1290–1293
- Reynolds A, Qiao H, Yang Y, Chen JK, Jackson N, Biswas K, Holloway JK, Baudat F, De Massy B, Wang J, Höög C, Cohen PE, Hunter N (2013). "RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis." *Nat Genet* 45:269–278 . <https://doi.org/10.1038/ng.2541>
- Ribarski I, Lehavi O, Yorgev L, Hauser R, Bar-Shira Maymon B, Botchan A, Paz G, Yavetz H, Kleiman SE (2009). "USP26 gene variations in fertile and infertile men." *Hum Reprod* 24:477–484 . <https://doi.org/10.1093/humrep/den374>
- Ribeiro J, Abby E, Livera G, Martini E (2016). "RPA homologs and ssDNA processing during meiotic recombination." *Chromosoma* 125:265–276
- Rimon-Dahari N, Yerushalmi-Heinemann L, Alyagor L, Dekel N (2016). "Ovarian folliculogenesis." *Results Probl Cell Differ* 58:167–190 . https://doi.org/10.1007/978-3-319-31973-5_7
- Robert T, Nore A, Brun C, Maffre C, Crimi B, Bourbon HM, De Massy B (2016). "The Topo VIB-Like protein family is required for meiotic DNA double-strand break formation." *Science* (80-) 351:943–949 .

- <https://doi.org/10.1126/science.aad5309>
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM (1998). "DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139." *J Biol Chem* 273:5858–5868
- Rogers RS, Inselman A, Handel MA, Matunis MJ (2004). "SUMO modified proteins localize to the XY body of pachytene spermatocytes." *Chromosoma* 113:233–243 . <https://doi.org/10.1007/s00412-004-0311-7>
- Romanienko PJ, Camerini-Otero RD (2000). "The mouse Spo11 gene is required for meiotic chromosome synapsis." *Mol Cell* 6:975–987 . [https://doi.org/10.1016/S1097-2765\(00\)00097-6](https://doi.org/10.1016/S1097-2765(00)00097-6)
- Rosenthal N, Brown S (2007). "The mouse ascending: perspectives for human-disease models." *Nat Cell Biol* 9:993–999 . <https://doi.org/10.1038/ncb437>
- Rossetti R, Ferrari I, Bonomi M, Persani L (2017). "Genetics of primary ovarian insufficiency." *Clin. Genet.* 91:183–198
- Saitou M, Barton SC, Surani MA (2002). "A molecular programme for the specification of germ cell fate in mice." *Nature* 418:293–300
- Saitou M, Miyauchi H (2016). "Gametogenesis from Pluripotent Stem Cells." *Cell Stem Cell* 18:721–735
- Saitou M, Yamaji M (2012). "Primordial germ cells in mice." *Cold Spring Harb Perspect Biol* 4: . <https://doi.org/10.1101/cshperspect.a008375>
- Sakai K, Ito C, Wakabayashi M, Kanzaki S, Ito T, Takada S, Toshimori K, Sekita Y, Kimura T (2019). "Usp26 mutation in mice leads to defective spermatogenesis depending on genetic background." *Sci Rep* 9: . <https://doi.org/10.1038/s41598-019-50318-6>
- Sasaki K, Nakamura T, Okamoto I, Yabuta Y, Iwatani C, Tsuchiya H, Seita Y, Nakamura S, Shiraki N, Takakuwa T, Yamamoto T, Saitou M (2016). "The Germ Cell Fate of Cynomolgus Monkeys Is Specified in the Nascent Amnion." *Dev Cell* 39:169–185 . <https://doi.org/10.1016/j.devcel.2016.09.007>
- Sasaki K, Yokobayashi S, Nakamura T, Okamoto I, Yabuta Y, Kurimoto K, Ohta H, Moritoki Y, Iwatani C, Tsuchiya H, Nakamura S, Sekiguchi K, Sakuma T, Yamamoto T, Mori T, Woltjen K, Nakagawa M, Yamamoto T, Takahashi K, Yamanaka S, Saitou M (2015). "Robust In Vitro Induction of Human Germ Cell Fate from Pluripotent Stem Cells." *Cell Stem Cell* 17:178–194 . <https://doi.org/10.1016/j.stem.2015.06.014>
- Scherthan H, Weich S, Schwegler H, Heyting C, Härle M, Cremer T (1996). "Centromere and telomere movements during early meiotic prophase of mouse and man are associated with the onset of chromosome pairing." *J Cell Biol* 134:1109–1125 . <https://doi.org/10.1083/jcb.134.5.1109>
- Schimenti JC, Handel MA (2018). "Unpackaging the genetics of mammalian fertility: strategies to identify the 'reproductive genome'." *Biol Reprod* 99:1119–1128 . <https://doi.org/10.1093/biolre/iox133>
- Schramm S, Fraune J, Naumann R, Hernandez-Hernandez A, Höög C, Cooke HJ, Alsheimer M, Benavente R (2011). "A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility." *PLoS Genet* 7:1002088 . <https://doi.org/10.1371/journal.pgen.1002088>
- Schürmann A, Koling S, Jacobs S, Saftig P, Krauß S, Wennemuth G, Kluge R, Joost H-G (2002). "Reduced Sperm Count and Normal Fertility in Male Mice with Targeted Disruption of the ADP-Ribosylation Factor-Like 4 (Arl4) Gene." *Mol Cell Biol* 22:2761–2768 . <https://doi.org/10.1128/mcb.22.8.2761-2768.2002>
- Schwanhüusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M (2011). "Global quantification of mammalian gene expression control." *Nature* 473:337–342 . <https://doi.org/10.1038/nature10098>
- Scully R, Chen J, Plug A, Xiao Y, Weaver D, Feunteun J, Ashley T, Livingston DM (1997). "Association of BRCA1 with Rad51 in mitotic and meiotic cells." *Cell* 88:265–275 . [https://doi.org/10.1016/S0092-8674\(00\)81847-4](https://doi.org/10.1016/S0092-8674(00)81847-4)
- Shah K, Sivapalan G, Gibbons N, Tempest H, Griffin DK (2003). "The genetic basis of infertility." *Reproduction* 126:13–25
- Shang Y, Huang T, Liu H, Liu Y, Liang H, Yu X, Li M, Zhai B, Yang X, Wei Y, Wang G, Chen Z, Wang S, Zhang L (2020). "MEIOK21: a new component of meiotic recombination bridges required for spermatogenesis." *Nucleic Acids Res* 48:6624–6639 . <https://doi.org/10.1093/nar/gkaa406>

- Sharan SK, Pyle A, Coppola V, Babus J, Swaminathan S, Benedict J, Swing D, Martin BK, Tessarollo L, Evans JP, Flaws JA, Handel MA (2004). "BRCA2 deficiency in mice leads to meiotic impairment and infertility." *Development* 131:131–142
- Shi B, Xue J, Yin H, Guo R, Luo M, Ye L, Shi Q, Huang X, Liu M, Sha J, Wang PJ (2019). "Dual functions for the ssDNA-binding protein RPA in meiotic recombination." *PLOS Genet* 15:e1007952 . <https://doi.org/10.1371/journal.pgen.1007952>
- Shi YC, Wei L, Cui YX, Shang XJ, Wang HY, Xia XY, Zhou YC, Li H, Jiang HT, Zhu WM, Huang YF (2011). "Association between ubiquitin-specific protease USP26 polymorphism and male infertility in Chinese men." *Clin Chim Acta* 412:545–549 . <https://doi.org/10.1016/j.cca.2010.12.005>
- Siaud N, Barbera MA, Egashira A, Lam I, Christ N, Schlacher K, Xia B, Jasin M (2011). "Plasticity of BRCA2 function in homologous recombination: Genetic interactions of the PALB2 and DNA binding domains." *PLoS Genet* 7: . <https://doi.org/10.1371/journal.pgen.1002409>
- Sinclair AH, Berta P, Palmer MS, Hawkins JR, Griffiths BL, Smith MJ, Foster JW, Frischau AM, Lovell-Badge R, Goodfellow PN (1990). "A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif." *Nature* 346:240–244 . <https://doi.org/10.1038/346240a0>
- Singh P, Schimenti JC, Bolcun-Filas E (2015). "A mouse geneticist's practical guide to CRISPR applications." *Genetics* 199:1–15 . <https://doi.org/10.1534/genetics.114.169771>
- Slizynski BM (1957). "Meiotic prophase in female mice." *Nature* 179:638
- Smagulova F, Gregoretti I V., Brick K, Khil P, Camerini-Otero RD, Petukhova G V. (2011). "Genome-wide analysis reveals novel molecular features of mouse recombination hotspots." *Nature* 472:375–378 . <https://doi.org/10.1038/nature09869>
- Smith BE, Braun RE (2012). "Germ cell migration across sertoli cell tight junctions." *Science (80-)* 338:798–802 . <https://doi.org/10.1126/science.1219969>
- Souquet B, Abby E, Hervé R, Finsterbusch F, Tourpin S, Le Bouffant R, Duquenne C, Messiaen S, Martini E, Bernardino-Sgherri J, Toth A, Habert R, Livera G (2013). "MEIOB Targets Single-Strand DNA and Is Necessary for Meiotic Recombination." *PLoS Genet* 9:e1003784 . <https://doi.org/10.1371/journal.pgen.1003784>
- Spradling A, Drummond-Barbosa D, Kai T (2001). "Stem cells find their niche." *Nature* 414:98–104
- Stanzione M, Baumann M, Papanikos F, Dereli I, Lange J, Ramlal A, Tränkner D, Shibuya H, De Massy B, Watanabe Y, Jasin M, Keeney S, Tóth A (2016). "Meiotic DNA break formation requires the unsynapsed chromosome axis-binding protein IHO1 (CCDC36) in mice." *Nat Cell Biol* 18:1208–1220 . <https://doi.org/10.1038/ncb3417>
- Stemmann O, Zou H, Gerber SA, Gygi SP, Kirschner MW (2001). "Dual inhibition of sister chromatid separation at metaphase." *Cell* 107:715–726 . [https://doi.org/10.1016/S0092-8674\(01\)00603-1](https://doi.org/10.1016/S0092-8674(01)00603-1)
- Stouffs K, Lissens W, Tournaye H, Van Steirteghem A, Liebaers I (2005). "Possible role of USP26 in patients with severely impaired spermatogenesis." *Eur J Hum Genet* 13:336–340 . <https://doi.org/10.1038/sj.ejhg.5201335>
- Stouffs K, Lissens W, Tournaye H, Van Steirteghem A, Liebaers I (2006). "Alterations of the USP26 gene in Caucasian men." *Int J Androl* 29:614–617 . <https://doi.org/10.1111/j.1365-2605.2006.00708.x>
- Takemoto K, Tani N, Takada-Horisawa Y, Fujimura S, Tanno N, Yamane M, Okamura K, Sugimoto M, Araki K, Ishiguro K ichiro (2020). "Meiosis-Specific C19orf57/4930432K21Rik/BRME1 Modulates Localization of RAD51 and DMC1 to DSBs in Mouse Meiotic Recombination." *Cell Rep* 31:107686 . <https://doi.org/10.1016/j.celrep.2020.107686>
- Tang WWC, Kobayashi T, Irie N, Dietmann S, Surani MA (2016). "Specification and epigenetic programming of the human germ line." *Nat. Rev. Genet.* 17:585–600
- Thorslund T, Esashi F, West SC (2007). "Interactions between human BRCA2 protein and the meiosis-specific recombinase DMC1." *EMBO J* 26:2915–2922 . <https://doi.org/10.1038/sj.emboj.7601739>
- Tian H, Huo Y, Zhang J, Ding S, Wang Z, Li H, Wang L, Lu M, Liu S, Qiu S, Zhang Q (2019). "Disruption of ubiquitin specific protease 26 gene causes male subfertility associated with spermatogenesis defects in mice." *Biol Reprod* 100:1118–1128 . <https://doi.org/10.1093/biolre/iory258>

- Toby GG, Gherraby W, Coleman TR, Golem EA (2003). "A Novel RING Finger Protein, Human Enhancer of Invasion 10, Alters Mitotic Progression through Regulation of Cyclin B Levels." *Mol Cell Biol* 23:2109–2122 . <https://doi.org/10.1128/mcb.23.6.2109-2122.2003>
- Tran TN, Schimenti JC (2019). "A segregating human allele of SPO11 modeled in mice disrupts timing and amounts of meiotic recombination, causing oligospermia and a decreased ovarian reserve." *Biol Reprod* 101:347–359 . <https://doi.org/10.1093/biolre/iez089>
- Turner JMA, Mahadevaiah SK, Fernandez-Capetillo O, Nussenzweig A, Xu X, Deng CX, Burgoyne PS (2005). "Silencing of unsynapsed meiotic chromosomes in the mouse." *Nat Genet* 37:41–47 . <https://doi.org/10.1038/ng1484>
- Tüttelmann F, Simoni M, Kliesch S, Ledig S, Dworniczak B, Wieacker P, Röpke A (2011). "Copy number variants in patients with severe oligozoospermia and sertoli-cell-only syndrome." *PLoS One* 6: . <https://doi.org/10.1371/journal.pone.0019426>
- Typas D, Luijsterburg MS, Wiegant WW, Diakatou M, Helfricht A, Thijssen PE, van den Broek B, Mullenders LH, van Attikum H (2016). "The de-ubiquitylating enzymes USP26 and USP37 regulate homologous recombination by counteracting RAP80." *Nucleic Acids Res* 44:2976 . <https://doi.org/10.1093/nar/gkv1480>
- Veitia RA (2003). "A sigmoidal transcriptional response: Cooperativity, synergy and dosage effects." *Biol. Rev. Camb. Philos. Soc.* 78:149–170
- Vigodner M, Morris PL (2005). "Testicular expression of small ubiquitin-related modifier-1 (SUMO-1) supports multiple roles in spermatogenesis: Silencing of sex chromosomes in spermatocytes, spermatid microtubule nucleation, and nuclear reshaping." *Dev Biol* 282:480–492 . <https://doi.org/10.1016/j.ydbio.2005.03.034>
- Wang H, La Russa M, Qi LS (2016). "CRISPR/Cas9 in Genome Editing and Beyond." *Annu Rev Biochem* 85:227–264 . <https://doi.org/10.1146/annurev-biochem-060815-014607>
- Wang J, Zhang W, Jiang H, Wu BL (2014). "Mutations in HFM1 in recessive primary ovarian insufficiency." *N. Engl. J. Med.* 370:972–974
- Wang JJ, Ge W, Liu JC, Klinger FG, Dyce PW, De Felici M, Shen W (2017a). "Complete in vitro oogenesis: Retrospects and prospects." *Cell Death Differ.* 24:1845–1852
- Wang PJ, McCarrey JR, Yang F, Page DC (2001). "An abundance of X-linked genes expressed in spermatogonia." *Nat Genet* 27:422–426 . <https://doi.org/10.1038/86927>
- Wang PJ, Page DC, McCarrey JR (2005). "Differential expression of sex-linked and autosomal germ-cell-specific genes during spermatogenesis in the mouse." *Hum Mol Genet* 14:2911–2918 . <https://doi.org/10.1093/hmg/ddi322>
- Wang S, Hassold T, Hunt P, White MA, Zickler D, Kleckner N, Zhang L (2017b). "Inefficient Crossover Maturation Underlies Elevated Aneuploidy in Human Female Meiosis." *Cell* 168:977–989.e17 . <https://doi.org/10.1016/j.cell.2017.02.002>
- Wear HM, McPike MJ, Watanabe KH (2016). "From primordial germ cells to primordial follicles: A review and visual representation of early ovarian development in mice." *J Ovarian Res* 9: . <https://doi.org/10.1186/s13048-016-0246-7>
- Weber JE, Russell LD (1987). "A study of intercellular bridges during spermatogenesis in the rat." *Am J Anat* 180:1–24 . <https://doi.org/10.1002/aja.1001800102>
- Weber S, Eckert D, Nettersheim D, Gillis AJM, Schäfer S, Kuckenberg P, Ehlermann J, Werling U, Biermann K, Looijenga LHJ, Schorle H (2010). "Critical Function of AP-2gamma/TCFAP2C in Mouse Embryonic Germ Cell Maintenance1." *Biol Reprod* 82:214–223 . <https://doi.org/10.1095/biolreprod.109.078717>
- Wei K, Clark AB, Wong E, Kane MF, Mazur DJ, Parris T, Kolas NK, Russell R, Hou H, Kneitz B, Yang G, Kunkel TA, Kolodner RD, Cohen PE, Edelmann W (2003). "Inactivation of exonuclease I in mice results in DNA mismatch repair defects, increased cancer susceptibility, and male and female sterility." *Genes Dev* 17:603–614 . <https://doi.org/10.1101/gad.1060603>
- Wilkins AS, Holliday R (2009). "The evolution of meiosis from mitosis." *Genetics* 181:3–12
- Wosnitzer MS, Mielnik A, Dabaja A, Robinson B, Schlegel PN, Paduch DA (2014). "Ubiquitin Specific Protease 26 (USP26) expression analysis in human testicular and extragonadal tissues indicates

- diverse action of USP26 in cell differentiation and tumorigenesis." *PLoS One* 9:e98638 .
<https://doi.org/10.1371/journal.pone.0098638>
- Wu Y, Liao S, Wang X, Wang S, Wang M, Han C (2013). "HSF2BP represses BNC1 transcriptional activity by sequestering BNC1 to the cytoplasm." *FEBS Lett* 587:2099–2104 .
<https://doi.org/10.1016/j.febslet.2013.04.049>
- Xia B, Sheng Q, Nakanishi K, Ohashi A, Wu J, Christ N, Liu X, Jasin M, Couch FJ, Livingston DM (2006). "Control of BRCA2 Cellular and Clinical Functions by a Nuclear Partner, PALB2." *Mol Cell* 22:719–729 .
<https://doi.org/10.1016/j.molcel.2006.05.022>
- Xia JD, Chen J, Han YF, Chen H, Yu W, Chen Y, Dai YT (2014). "Association of 370-371insACA, 494T>C, and 1423C>T haplotype in ubiquitin-specific protease 26 gene and male infertility: a meta-analysis." *Asian J Androl* 16:720–724 . <https://doi.org/10.4103/1008-682X.129134>
- Xu X, Aprelikova O, Moens P, Deng CX, Furth PA (2003). "Impaired meiotic DNA-damage repair and lack of crossing-over during spermatogenesis in BRCA1 full-length isoform deficient mice." *Development* 130:2001–2012
- Xu Y, Greenberg RA, Schonbrunn E, Jeremy Wang P (2017). "Meiosis-specific proteins MEIOB and SPATA22 cooperatively associate with the single-stranded DNA-binding replication protein A complex and DNA double-strand breaks." *Biol Reprod* 96:1096–1104 . <https://doi.org/10.1093/biolre/iox040>
- Yamada S, Hinch AG, Kamido H, Zhang Y, Edlmann W, Keeney S (2020). "Molecular structures and mechanisms of DNA break processing in mouse meiosis." *Genes Dev* 34:806–818 .
<https://doi.org/10.1101/gad.336032.119>
- Yamaji M, Seki Y, Kurimoto K, Yabuta Y, Yuasa M, Shigeta M, Yamanaka K, Ohinata Y, Saitou M (2008). "Critical function of Prdm14 for the establishment of the germ cell lineage in mice." *Nat Genet* 40:1016–1022 . <https://doi.org/10.1038/ng.186>
- Yang F, Eckardt S, Leu NA, McLaughlin KJ, Wang PJ (2008). "Mouse TEX15 is essential for DNA double-strand break repair and chromosomal synapsis during male meiosis." *J Cell Biol* 180:673–679 .
<https://doi.org/10.1083/jcb.200709057>
- Yang H, Jeffrey PD, Miller J, Kinnucan E, Sun Y, Thomä NH, Zheng N, Chen PL, Lee WH, Pavletich NP (2002). "BRCA2 function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure." *Science* (80-) 297:1837–1848 . <https://doi.org/10.1126/science.297.5588.1837>
- Yang H, Wang H, Jaenisch R (2014). "Generating genetically modified mice using CRISPR/Cas-mediated genome engineering." *Nat Protoc* 9:1956–1968 . <https://doi.org/10.1038/nprot.2014.134>
- Yatsenko AN, Georgiadis AP, Murthy LJ, Lamb DJ, Matzuk MM (2013). "UBE2B mRNA alterations are associated with severe oligozoospermia in infertile men." *Mol Hum Reprod* 19:388–394 .
<https://doi.org/10.1093/molehr/gat008>
- Yi YJ, Manandhar G, Oko RJ, Breed WG, Sutovsky P (2007). "Mechanism of sperm-zona pellucida penetration during mammalian fertilization: 26S proteasome as a candidate egg coat lysin." *Soc Reprod Fertil Suppl* 63:385–408
- Yoshida K, Kondoh G, Matsuda Y, Habu T, Nishimune Y, Morita T (1998). "The mouse RecA-like gene Dmc1 is required for homologous chromosome synapsis during meiosis." *Mol Cell* 1:707–718 .
[https://doi.org/10.1016/S1097-2765\(00\)80070-2](https://doi.org/10.1016/S1097-2765(00)80070-2)
- Yoshima T, Yura T, Yanagi H (1998). "Novel testis-specific protein that interacts with heat shock factor 2." *Gene* 214:139–146 . [https://doi.org/10.1016/S0378-1119\(98\)00208-X](https://doi.org/10.1016/S0378-1119(98)00208-X)
- Youds JL, Boulton SJ (2011). "The choice in meiosis - Defining the factors that influence crossover or non-crossover formation." *J. Cell Sci.* 124:501–513
- Zakharyevich K, Ma Y, Tang S, Hwang PYH, Boiteux S, Hunter N (2010). "Temporally and Biochemically Distinct Activities of Exo1 during Meiosis: Double-Strand Break Resection and Resolution of Double Holliday Junctions." *Mol Cell* 40:1001–1015 . <https://doi.org/10.1016/j.molcel.2010.11.032>
- Zhang B, Tang Z, Li L, Lu LY (2020a). "NBS1 is required for SPO11-linked DNA double-strand break repair in male meiosis." *Cell Death Differ* 27:2176–2190 . <https://doi.org/10.1038/s41418-020-0493-4>
- Zhang H, Liu L, Li X, Busayavalasa K, Shen Y, Hovatta O, Gustafsson JÅ, Liu K (2014). "Life-long in vivo cell-lineage tracing shows that no oogenesis originates from putative germline stem cells in adult mice."

- Proc Natl Acad Sci U S A* 111:17983–17988 . <https://doi.org/10.1073/pnas.1421047111>
- Zhang J, Fujiwara Y, Yamamoto S, Shibuya H (2019). “A meiosis-specific BRCA2 binding protein recruits recombinases to DNA double-strand breaks to ensure homologous recombination.” *Nat Commun* 10:1–14 . <https://doi.org/10.1038/s41467-019-08676-2>
- Zhang J, Gurusaran M, Fujiwara Y, Zhang K, Echbarthi M, Vorontsov E, Guo R, Pendlebury DF, Alam I, Livera G, Emmanuelle M, Wang PJ, Nandakumar J, Davies OR, Shibuya H (2020b). “The BRCA2-MEILB2-BRME1 complex governs meiotic recombination and impairs the mitotic BRCA2-RAD51 function in cancer cells.” *Nat Commun* 11:1–18 . <https://doi.org/10.1038/s41467-020-15954-x>
- Zhang J, Qiu SD, Li SB, Zhou DX, Tian H, Huo YW, Ge L, Zhang QY (2007). “Novel mutations in ubiquitin-specific protease 26 gene might cause spermatogenesis impairment and male infertility.” *Asian J Androl* 9:809–814 . <https://doi.org/10.1111/j.1745-7262.2007.00305.x>
- Zhang W, Liu T, Mi YJ, Yue LD, Wang JM, Liu DW, Yan J, Tian QB (2015). “Evidence from enzymatic and meta-analyses does not support a direct association between USP26 gene variants and male infertility.” *Andrology* 3:271–279 . <https://doi.org/10.1111/andr.295>
- Zhao W, Vaithiyalingam S, San Filippo J, Maranon DG, Jimenez-Sainz J, Fontenay G V., Kwon Y, Leung SG, Lu L, Jensen RB, Chazin WJ, Wiese C, Sung P (2015). “Promotion of BRCA2-Dependent Homologous Recombination by DSS1 via RPA Targeting and DNA Mimicry.” *Mol Cell* 59:176–187 . <https://doi.org/10.1016/j.molcel.2015.05.032>
- Zhou Q, Wang M, Yuan Y, Wang X, Fu R, Wan H, Xie M, Liu M, Guo X, Zheng Y, Feng G, Shi Q, Zhao XY, Sha J, Zhou Q (2016). “Complete Meiosis from Embryonic Stem Cell-Derived Germ Cells in Vitro.” *Cell Stem Cell* 18:330–340 . <https://doi.org/10.1016/j.stem.2016.01.017>
- Zorrilla M, Yatsenko AN (2013). “The Genetics of Infertility: Current Status of the Field.” *Curr Genet Med Rep* 1:247–260 . <https://doi.org/10.1007/s40142-013-0027-1>

PUBLICACIONES

Ubiquitin-specific protease 26 (USP26) is not essential for mouse gametogenesis and fertility

Natalia Felipe-Medina¹ · Laura Gómez-H¹ · Yazmine B. Condezo¹ · Manuel Sanchez-Martín^{2,3} · José Luis Barbero⁴ · Isabel Ramos¹ · Elena Llano^{1,5} · Alberto M. Pendás¹

Received: 24 October 2018 / Revised: 18 February 2019 / Accepted: 22 February 2019 / Published online: 18 March 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Ubiquitin-specific protease 26 (USP26) is a deubiquitylating enzyme belonging to the USPs family with a transcription pattern restricted to the male germline. Since protein ubiquitination is an essential regulatory mechanism during meiosis, many efforts have been focused on elucidating the function of USP26 and its relationship with fertility. During the last decade, several studies have reported the presence of different polymorphisms in *USP26* in patients with non-obstructive azoospermia (NOA) or severe oligozoospermia suggesting that this gene may be associated with human infertility. However, other studies have revealed the presence of these and novel polymorphisms, including nonsense mutations, in men with normal spermatogenesis as well. Thus, the results remain controversial and its function is unknown. In the present study, we describe the *in vivo* functional analysis of mice lacking USP26. The phenotypic analysis of two different *Usp26*-null mutants showed no overt-phenotype with both males and females being fertile. Cytological analysis of spermatocytes showed no defects in synapsis, chromosome dynamics, DNA repair, or recombination. Histopathological analysis revealed a normal distribution and number of the different cell types in both male and female mice. Finally, normal counts were observed in fertility assessments. These results represent the first *in vivo* evidence showing that USP26 is not essential for mouse gametogenesis.

Keywords Fertility · Spermatogenesis · Ubiquitination · Meiosis · Ubiquitin-specific protease

This article is part of a Special Issue on Recent advances in meiosis from DNA replication to chromosome segregation" edited by Valérie Borde and Francesca Cole, co-edited by Paula Cohen and Scott Keeney".

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00412-019-00697-6>) contains supplementary material, which is available to authorized users.

✉ Alberto M. Pendás
amp@usal.es

¹ Molecular Mechanisms Program, Centro de Investigación del Cáncer and Instituto de Biología Molecular y Celular del Cáncer (CSIC-Universidad de Salamanca), 37007 Salamanca, Spain

² Departamento de Medicina, Universidad de Salamanca, Salamanca 37007, Spain

³ Transgenic Facility, Nucleus platform, Universidad de Salamanca, Salamanca 37007, Spain

⁴ Departamento de Biología Celular y Molecular, Centro de Investigaciones Biológicas (CSIC), Madrid 28040, Spain

⁵ Departamento de Fisiología y Farmacología, Universidad de Salamanca, Salamanca 37007, Spain

Introduction

Infertility is a major health issue with an estimated 8–12% of couples affected worldwide (Ombelet et al. 2008; Prieto et al. 2004). Men contribute to 50% and are solely responsible for 20–30% of reported cases (Sharlip et al. 2002). Many factors have been associated with male infertility such as environmental or occupational exposures and genetic abnormalities. Genetic causes are thought to underlie 15–30% of male infertilities being non-obstructive azoospermia (NOA), the complete absence of spermatozoa in the ejaculate due to a testicular failure, the most common outcome associated (Ferlin et al. 2007). Although chromosomal aberrations and microdeletions of the Y chromosome have been largely considered to be the main cause of genetic male infertility (Matzuk and Lamb 2008; Reijo et al. 1996), in the past decades, a large number of fertility-related genes have been identified through the analysis of mouse models, clinical cases, and genome-wide association studies (Caburet et al. 2014; Gomez et al. 2016; Hu et al. 2011). However, much work remains to be done as idiopathic azoospermia is still the most

frequent diagnosis among infertile males, and it is currently believed that a high proportion is related with genetic causes (Hamada et al. 2013).

Spermatogenesis, the process through which spermatogonial stem cells generate mature spermatozoa, represents one of the most complex differentiation programs in developmental biology and requires an accurate spatio-temporal regulation of gene transcription, translation, and protein turnover. In this sense, ubiquitination becomes essential for the correct completion of spermatogenesis as it targets substrates for proteasomal degradation but also can modify their biological activity, stability, or subcellular localization (He et al. 2016). Moreover, the importance of ubiquitination in the regulation of protein turnover during spermatogenesis as well as in mammalian fertilization and sperm quality has been reported (Baarends et al. 2000; Muratori et al. 2005; Yi et al. 2007).

The ubiquitin-specific protease 26 (*Usp26*) is a single-exon gene firstly identified by Wang et al. (2001) from a screen for genes expressed in mouse spermatogonia. USP26 belongs to a family of cysteine-proteases that disassemble polymeric ubiquitin chains from substrates and is responsible for maintaining ubiquitin homeostasis in cells (Komander et al. 2009). *USP26* has attracted increased attention as a candidate infertility gene due to its restricted expression pattern to testis and because of its genetic linkage with X-chromosome (male are hemizygous) which is enriched for testis-specific genes (Nishimune and Tanaka 2006). More specifically, *USP26* expression has been described in spermatogonia (types A and B), preleptotene and leptotene-zygotene spermatocytes, round spermatids, and at the blood-testis barrier from both mouse and human testes (Lin et al. 2011; Wang et al. 2005). Additionally, in humans, it is also expressed in Sertoli and Leydig cells (Wosnitzer et al. 2014).

Genetic polymorphisms identified in the *USP26* locus have been causally associated with male infertility from Sertoli cell only-syndrome to non-obstructive azoospermia or asthenozoospermia (Asadpor et al. 2013; Lee et al. 2008; Li et al. 2015; Ma et al. 2016; Paduch et al. 2005; Stouffs et al. 2005; Xia et al. 2014; Zhang et al. 2007). By contrast, other studies have found the same frequency of *USP26* polymorphisms in fertile and infertile men (Christensen et al. 2008; Luddi et al. 2016; Ravel et al. 2006; Ribarski et al. 2009; Shi et al. 2011; Stouffs et al. 2006; Zhang et al. 2015). In addition, it has been reported that most of the described polymorphisms do not abolish USP26 enzymatic activity (Liu et al. 2018; Zhang et al. 2015).

Given the absence of functional studies, we have addressed the in vivo functional analysis of mice lacking USP26. The analysis of USP26-deficient mice showed no overt-phenotype with both males and females being fertile. The histopathological analysis of testis and ovaries showed a normal number and distribution of cell types, and accordingly, the results obtained in the fertility assessment were similar to the wild-type

ones. Finally, spermatogenesis in the absence of USP26 proceeded without any defect neither in synapsis nor in DNA repair or meiotic recombination. These results represent the first in vivo evidence showing that USP26 is not essential for mouse gametogenesis.

Methods

Production of CRISPR/Cas9-edited mice *Usp26*-sgRNAs were predicted at crispr.mit.edu. *Usp26*-sgRNAs were produced by cloning annealed complementary oligos (Table S1) at the BbsI site of pX330 (#42230, Addgene), generating PCR products containing a T7 promoter sequence, and then performing in vitro transcription using the MEGAshortscript™ T7 Transcription Kit (Life Technologies). The plasmid pST1374-NLS-flag-linker-Cas9 (#44758; Addgene) was used for generating CAS9 mRNA after linearization with AgeI. In vitro transcription and capping were performed using the mMESSAGE mMACHINE T7 Transcription Kit (AM1345; Life Technologies). Products were purified using the RNeasy® Mini Kit (Qiagen). RNA (100 ng/ul Cas9 and 50 ng/ul each guide RNA) was microinjected into zygotes (F1 hybrids between strains C57BL6/J and CBA/J also known as B6CBAF1/J) as described previously (Singh et al. 2015). Edited founders were identified by PCR amplification with primers flanking the targeted regions (Table S2) and subcloned into pBlueScript (Stratagene) followed by standard Sanger sequencing. The lengths of the corresponding wild-type and mutant alleles were 307 bp and 411 bp respectively, in the 21 mutants and 259 bp vs 173 bp in the catalytic mutants. The selected founder was crossed with a wild type (B6CBAF1/J) to eliminate possible unwanted off-targets. *Usp26*^(WT/edited) heterozygous mice were sequenced again by Sanger and crossed to give rise to *Usp26*^(edited/edited) homozygous. Genotyping was performed by agarose gel electrophoresis analysis of PCR products from DNA isolated from tail biopsies.

Fertility assessment *Usp26*^{WT/Y}, *Usp26*^{21/Y}, and *Usp26*^{Cat/Y} males and *Usp26*^{WT/WT}, *Usp26*^{21/21}, and *Usp26*^{Cat/Cat} females (8 weeks old) were mated with wild-type females and males, respectively during 3 months. Six mice per genotype were crossed. The presence of copulatory plug was examined daily, and the number of pups per litter was recorded.

Histology For histological analysis, adult testes/ovaries were fixed in Bouin's fixative, processed into serial paraffin sections, and stained with Periodic acid–Schiff (PAS) and Hematoxylin (testes) or hematoxylin-eosin (ovaries).

Immunocytochemistry and antibodies Testes were detunicated and processed for spreading using the “dry-down” technique and incubated with the indicated antibodies for

Fig. 1 *Usp26* knockout generation by CRISPR-Cas9 genome editing. **a** Schematic representation of the wild-type locus (WT) and the two genome editing strategies at the *Usp26* locus (left: knockout 21, right: catalytic knockout Cat). The sgRNAs, the single-coding exon of *Usp26* (gray) and the corresponding non-coding region (open box) are represented. Thin (non-coding) and thick (coding sequences) lines under the exon represent the expected transcripts derived from WT and *Usp26* edited alleles. ATG, start codon; TAG/TAA, stop codons; TGC, codon encoding the catalytic Cys295; Restart ATG, codons encoding the Met147(21) and Met331(Cat). Nucleotide sequence of the WT and the edited allele derived from PCR amplification of DNA from the *Usp26* edited mice is shown. In the allele^{Cat}, the catalytic cysteine (Cys) is highlighted (red rectangle). The insertion (left panel) and deletion (right panel) generate a frameshift (amino acids in red) and a premature STOP codon. Double-headed arrows represent the predicted size (KDa) of the

peptides that could be expressed in both alleles (from the canonical ATG to the generated STOP codon and from the first ATG after the mutation - restart ATG- to the canonical STOP codon). Arrows represent genotyping primers. **b** PCR analysis of genomic DNA from three pups from *Usp26*^{WT/21} heterozygous crosses. The PCR amplification with primers F and R (arrows) revealed a fragment of 307 bp in the case of the allele^{WT} and 411 bp in the allele²¹ (WT/WT), (WT/21), and (21/21) designate wild-type, heterozygous, and homozygous knockout animals, respectively. **c** PCR analysis of genomic DNA from three pups from *Usp26*^{WT/Cat} heterozygous crosses. The PCR amplification with primers F and R (arrows) revealed a fragment of 259 bp in the case of the allele^{WT} and 173 bp in the allele^{Cat} (WT/WT), (WT/Cat), and (Cat/Cat) designate wild-type, heterozygous, and homozygous knockout females, respectively

immunofluorescence. The primary antibodies used for immunofluorescence were goat α USP26 (W-12) sc-51013 (Santa Cruz, 1:5), rabbit α USP26-M (1:5, provided by Dr. Yi-Wen Lin Institute of Biomedical Sciences, Taipei, Taiwan), mouse α SYCP3 IgG sc-74569 (Santa Cruz, 1:60), rabbit α SYCP3 serum K921 (Prieto et al. 2004), rabbit α SYCP1 IgG ab15090 (Abcam, 1:200), rabbit anti- γ H2AX (ser139) IgG #07-164 (Millipore, 1:150), and mouse α MLH1 51-1327GR (BD Biosciences; 1:5). The secondary antibodies used were

TRITC α -mouse 115-095-146/ α -rabbit 111-025-144 and FITC α -mouse 115-095-146/ α -rabbit 111-095-045 (Jackson ImmunoResearch, all 1:100). Slides were visualized at room temperature using a microscope (Axioplan 2; Carl Zeiss, Inc.) with 63 $\times$ objectives with an aperture of 1.4 (Carl Zeiss, Inc.). Images were taken with a digital camera (ORCA-ER; Hamamatsu) and processed with OPENLAB 4.0.3 and Photoshop (Adobe). Quantification of γ H2AX fluorescence signal was measured by ImageJ software.

Protein extraction and western blotting Whole testis extracts were prepared in RIPA buffer (50 mM Tris HCl (pH 7.5), 150 mM NaCl, 1% NP40, 0.1% SDS, 0.5% sodium deoxycholate). Fifteen micrograms of total protein was loaded onto 10% reducing polyacrylamide SDS gels and transferred to 0.45 μ m Hybond ECL nitrocellulose membranes (RPN303D; GE Healthcare, Life Sciences). The membranes were blocked with 5% nonfat dried milk diluted in TBS-Tween. The proteins were detected by western blotting with the indicated antibodies. Primary antibodies used for western blotting were rabbit goat α USP26 (W-12) sc-51013 (Santa Cruz, 1:2000), rabbit α USP26-M (1:2000, provided by Dr. Yi-Wen Lin, Institute of Biomedical Sciences, Taipei, Taiwan), mouse anti- α -Tubulin (T9026, Sigma, 1:40000), and mouse anti- β -Actin (A5441, Sigma, 1:40000). Secondary horseradish peroxidase-conjugated α -mouse (715-035-150, Jackson

ImmunoResearch), α -rabbit (711-035-152, Jackson ImmunoResearch), or α -goat (705-035-147, Jackson ImmunoResearch) antibodies were used at 1:5000 dilution. Antibodies were detected by using Immobilon™ Western Chemiluminescent HRP Substrate from Millipore.

Bioinformatic analysis of RNAseq datasets Gene expression data were obtained from each article repository and processed to extract the relevant information of the genes of interest (*USP* genes) into a dataframe using regular expressions in R language. Colors have been included to represent a gradient of expression for each individual gene. Red and green represent the lower and upper end-points (stages with lower and higher expression level) and between them an intermediate color gradient has been established.

Fig. 2 Histopathological analysis does not show defects in USP26-deficient mice. **a** Testis size was not affected by the absence of USP26. **b** PAS + hematoxylin-stained testis sections show a normal number and distribution of germ cells in the seminiferous tubules (St) and normal counts of spermatozoa in the epididymides (Ep) of *USP26*^{21/Y} and *USP26*^{Cat/Y} mice in comparison with *USP26*^{WT/Y} mice. Bar in upper panels (St), 10 μ m, and in lower panels (Ep), 20 μ m. **c** Comparative histopathological analysis (hematoxylin + eosin staining) of ovaries from *USP26*^{21/21}, *USP26*^{Cat/Cat}, and *USP26*^{WT/WT} females at 4 months of age showing the presence of follicles and corpora lutea and a normal appearance of the stroma in the absence of USP26. Bars represent 50 μ m

Table 1 USP26 deficiency has no impact on fertility

		Mean (offspring)	SD	n
Males	USP26 ^{WT/Y}	7.0	1.47	15
	USP26 ^{21/Y}	6.9	2.31	18
	USP26 ^{Cat/Y}	6.4	2.32	16
Females	USP26 ^{WT/WT}	7.4	2.33	14
	USP26 ^{21/21}	7.2	1.89	16
	USP26 ^{Cat/Cat}	7.9	2.28	16

Numeric data (mean of pups per litter) obtained from the mating of USP26^{WT/Y}, USP26^{21/Y}, and USP26^{Cat/Y} males and USP26^{WT/W}, USP26^{21/21}, and USP26^{Cat/Cat} females with USP26^{WT/WT} females and USP26^{WT/Y} males, respectively. 8-week-old mice were mated for 3 months (6 mice/genotype)

Results

Usp26 CRISPR-Cas9 knockout lines

To gain further insight into USP26 function, we generated a mouse model by targeting the single-coding exon of *Usp26* through CRISPR-Cas9 genome editing. The selected founder carried two consecutive insertions of 85 and 19 base pairs (bp) separated by 5 bp. The first insertion changes the frame and generates a premature STOP codon at position 123 (14KDa) resulting in a predicted null-allele (herein allele²¹; Fig. 1a, left). This founder was crossed with wild-type (WT) C57BL/6J mice, and the heterozygous offspring was crossed to obtain the homozygous mutant mice, which were identified by PCR (Fig. 1b).

To test whether the generated mutation was a null-allele, we analyzed the expression of USP26 in mouse testis with two different antibodies, the commercial anti-USP26 (W-12) and the anti-USP26-M kindly provided by Dr. Yi-Wen Lin (Lin et al. 2011). We first carried out an immunofluorescence (IF) analysis in spermatocyte spreads from both *Usp26*^{21/Y} and *Usp26*^{WT/Y} mice. None of the antibodies showed detectable signal in spermatocytes (leptonema, zygonema, pachynema, diplonema, or diakinesis), spermatogonia, or round spermatid (Fig. S1a–b and S2a–b), where the expression of USP26 has been previously reported (Lin et al. 2011; Wang et al. 2005). We also performed a western blot (WB) analysis of whole testis extracts from wild-type and mutant mice. We did not observe any signal in the blot using the anti-USP26 (W-12) antibody (Fig. S1c) whereas the anti-USP26-M generated similar bands in *Usp26*^{21/Y} and *Usp26*^{WT/Y} extracts (Fig. S2c). No shift in the migration of any detected band was observed in the mutant extracts in comparison with the wild type (USP26 size: WT, 95 KDa vs 14 KDa, allele²¹). These results strongly suggest that the available antibodies are not able to recognize the endogenous protein and, consequently, the detected bands do not correspond to USP26.

Nonsense mutations close to the 5' of open reading frames (ORF) can lead to a translation restart from downstream AUG codons using the same original frame (Makino et al. 2016). The genomic locus of murine *Usp26* encodes an in-frame methionine at the position 147 which could be used as a downstream AUG initiation codon in our initial mutant (allele²¹) and could potentially lead to the generation of an N-terminal truncated protein (79 KDa, Fig. 1a, left) with an intact catalytic domain. We did not detect the predicted truncated proteins of 14 KDa (premature STOP codon) or 79 KDa (Restart ATG) in the *Usp26*^{21/Y} extracts by western blot (Fig. S1c and S2c). Consequently, the absence of working antibodies against USP26 that allowed us to ensure that the allele²¹ was a null mutant led us to generate a second mutant with a deletion in the proteolytic domain of USP26 (dead mutant). This domain is formed by two short motifs known as Cys and His boxes that contain the catalytic triad, one catalytic cysteine, and an histidine and asparagine/aspartate that facilitate the cysteine nucleophilic attack (Davis and Simeonov 2015; Nijman et al. 2005). To abolish the protease activity of USP26, we targeted the CRISPR-Cas9 to the catalytic cysteine (Cys295) generating an 86 bp deletion (including the cysteine) that produces a frameshift and consequently a premature STOP codon at position 293 (34KDa; Fig. 1a, right). Once the founder transmitted the allele, the heterozygous was crossed to obtain the homozygous mutants (herein, *Usp26*^{Cat/Cat} or *Usp26*^{Cat/Y}), which were identified by PCR (Fig. 1c). Immunofluorescence and western blot analysis were performed for the new catalytic mutant, and the results were identical to those obtained in the *Usp26*^{21/Y} mutant (Fig. S1a–c and S2a–c). The western blot analysis did not reveal any shift on band migration nor identify the truncated proteins of 34 and/or 56 KDa that could be expressed from the allele^{Cat} (See Fig. 1a right and S1c and S2c). Thus, although we cannot demonstrate the absence of USP26 in our mutants due to the lack of working antibodies that recognize the endogenous protein, the genetic strategy predicts that both of them are null. Especially, the resultant catalytic mutant lacks the genomic information to encode an active catalytic domain (Cys) and, consequently, represents a dead mutant without ubiquitin-specific protease activity.

USP26-deficient mice are fertile

Usp26^{21/Y} and *Usp26*^{Cat/Y} mice developed normally and showed no overt-phenotype including testis size and weight (Fig. 2a). To determine the potential effect of USP26 absence on mouse fertility, we performed a fertility test. Adult males from *Usp26*^{WT/Y}, *Usp26*^{21/Y}, and *Usp26*^{Cat/Y} lines were mated with *Usp26*^{WT/WT} females. The presence of copulatory plug was monitored daily to discard behavioral defects, and the number of pups per litter was recorded. USP26 mutants behaved similar to their wild-type counterparts, and no

Fig. 3 Chromosome behavior and synapsis proceed properly in spermatocytes lacking USP26. Double immunolabeling of SYCP3 (red) and SYCP1 (green) on spermatocyte spreads from *USP26*^{21/Y}, *USP26*^{Cat/Y},

and *USP26*^{WT/Y} mice. DNA was stained with DAPI (blue). No differences were observed on synapsis/desynapsis processes between *USP26*-deficient and WT spermatocytes. Bar in panels, 10 μ m

significant differences in litter size between the three groups were observed (Table 1). The histopathological analysis of seminiferous tubules from *Usp26*^{21/Y} and *Usp26*^{Cat/Y} mice showed a normal progression through the 12 stages of epithelial cycle with an organization and number of germ cells similar to the wild-type ones (Fig. 2b). Moreover, a normal amount of spermatozoa were able to reach the epididymides of *USP26* mutants (Fig. 2b). Taken together, our findings demonstrate that *USP26* seems to have a minimal if any role in mouse male fertility.

Usp26 is not expressed in human whole ovary and mutations associated with female infertility have not been reported. Accordingly, the fertility assessment and histological analysis of *Usp26*^{21/21} and *Usp26*^{Cat/Cat} females did not reveal any defect (Table 1 and Fig. 2c).

USP26-deficient spermatocytes showed normal synapsis, DSBs dynamics, and recombination

Although *Usp26* transcription starts in spermatogonia, it is transcribed at maximum levels during the early stages of

meiosis (leptotene and zygotene, Tables S5 and S6 (da Cruz et al. 2016; Margolin et al. 2014)). Moreover, the ubiquitin–proteasome system (UPS) has been involved in the regulation of meiotic prophase in mouse (Rao et al. 2017) and *USP26* has been associated with double-strand breaks (DSBs) repair and homologous recombination in somatic cells (Typas et al. 2016). These data prompted us to analyze in more detail chromosome synapsis, double-strand breaks (DSBs) generation/resolution, and meiotic recombination during the meiotic prophase I of *USP26*-deficient spermatocytes.

Chromosome synapsis is mediated by a proteinaceous scaffold known as synaptonemal complex (SC). The SC is composed of two axial elements (AEs) connected by transverse filaments (TFs) through the assembly of the central element (CE), generating a tripartite zipper-like structure. In mammals, the components of AEs are SYCP2 and SYCP3, TFs consist of SYCP1, and five CE components have been identified as follow: SYCE1-3, TEX12, and the recently described SIX6OS1 (Bolcun-Filas and Schimenti 2012; Gomez et al. 2016). To evaluate the synaptic process, we analyzed the distribution of SYCP3 and SYCP1 in

Fig. 4 DSB dynamics is not affected in the absence of USP26. **a** Double immunofluorescence of γ -H2AX (green) with SYCP3 (red) in wild-type and *USP26* mutant spermatocytes showing no differences in the formation and repair of DSBs. DNA was stained with DAPI (blue). DSB formation occurs at leptotene, and they are processed during zygotene and early pachytene. Consequently, γ -H2AX labels intensely the whole nucleus at leptotene and declines until mid-pachytene where the staining is

restricted to the unsynapsed sex bivalent. **b** Plots under each panel represent the quantification of fluorescence intensity at the different stages and shows no significant differences ($n = 60$ cells from 2 different mice for each genotype). See Table S3 for numeric data; bars represent standard deviation (SD). Welch's t test analysis: * $p < 0.01$; ** $p < 0.001$; *** $p < 0.0001$. Bar in panels, 10 μ m

spermatocyte spreads from *Usp26*^{21/Y}, *Usp26*^{Cat/Y}, and *Usp26*^{WT/Y} mice (Fig. 3). We did not observe defects on AEs formation, starting as short patches in leptotene that got longer as zygotene proceeded. Full synapsis was achieved at pachytene, and no defects were observed as desynapsis progressed from diplotene to diakinesis. Additionally, USP26-deficient metaphase I and II plates showed normal morphology (Fig. 3). Altogether, these observations suggest that USP26 has no function on chromosome behavior and synapsis.

Meiotic recombination establishes the physical tethering between homologous chromosomes essential to ensure the correct chromosome segregation during the first meiotic division. Meiotic recombination is initiated at leptotene with the formation of programmed DSBs by SPO11 which are

repaired during the prophase I by the homologous recombination (HR) machinery (Bolcun-Filas and Schimenti 2012). Given the involvement of protein ubiquitination and SUMOylation in DSBs repair (Bekker-Jensen and Mailand 2011), we analyzed USP26-deficient spermatocytes for the labeling of phosphorylated H2AX (γ -H2AX), the first appearing modification after DSBs induction (Rogakou et al. 1998). The γ -H2AX distribution in USP26-deficient spermatocytes resembled that of wild-type cells from leptotene, showing whole nucleus labeling, to pachytene where the γ -H2AX signal was restricted to the sex body. These data indicate that DSBs formation and repair are not compromised by the absence of USP26 (Fig. 4).

During early prophase I, DSBs are frequently resolved as non-crossovers (NCO) and only a reduced fraction of the

Fig. 5 USP26-deficient mice do not show meiotic recombination defects. **a** Immunostaining of MLH1 (green) and SYCP3 (red). DNA was stained with DAPI (blue). The number of MLH1 foci (corresponding to future chiasmata) in mutant spermatocytes resembles that observed in the wild type. Bar in panels, 10 μ m. **b** Quantification of MLH1 foci in *USP26*^{WT/Y}, *USP26*^{21/Y}, and *USP26*^{Cat/Y} pachytene spermatocytes showing no significant differences ($n = 52$, 52, and 48 cells respectively from 2 different mice for each genotype). See Table S4 for numeric data; bars represent standard deviation (SD). Welch's t -test analysis: * $p < 0.01$; ** $p < 0.001$; *** $p < 0.0001$

DSBs are processed as crossovers (CO), which will give rise to chiasmata. Owing to the association of the UPS to the AEs and its role in regulating meiotic recombination (Rao et al. 2017), we evaluated USP26-deficient spermatocytes for CO formation by MLH1 staining, a mismatch repair protein that marks CO sites (Guillon et al. 2005). No differences were observed in the number of MLH1 foci between *Usp26*^{21/Y}, *Usp26*^{Cat/Y}, and *Usp26*^{WT/Y} spermatocytes (Fig. 5).

Discussion

In the present study, we have generated two *Usp26* knockout models, one carrying two insertions (allele²¹) and a second one lacking the catalytic Cys of the protein (dead mutant, allele^{Cat}). Although the in silico analysis predicts that both alleles generate truncated proteins, we have not been able to demonstrate the absence of USP26 in the mutants, likely due

to the lack of sensitivity of the antibodies which failed to detect the endogenous protein. Even though we could not demonstrate the absence of protein, the generated deletion in the catalytic domain of USP26 (allele^{Cat}) ensures that the protein, if present, will not be functional as ubiquitin-specific protease leading to a reliable loss-of-function mutant.

The testis-specific USP26 belongs to the superfamily of deubiquitinating enzymes (DUBs) that remove ubiquitin (Ub) peptides from substrates, process Ub precursors, and disassemble polymeric Ub chains hence being responsible for maintaining ubiquitin homeostasis in cells (Komander et al. 2009). Ubiquitination plays essential roles during spermatogenesis from the establishment and differentiation of spermatogonial stem cells to spermiogenesis (Bose et al. 2014) and to date several deubiquitinating enzymes (i.e., USP2 (Bedard et al. 2011), USP7 (Luo et al. 2015), USP9X (Kishi et al. 2017), Uchl-1 (Kwon et al. 2005)) have been associated with defects at different stages of spermatogenesis and fertility impairment. However, here we show that mice lacking the deubiquitinating enzyme USP26 are fertile showing litters of the same size than those from wild-type mice and a normal number and distribution of all the testis cell types.

The maximum transcription levels of *Usp26* are reached during the first stages of meiotic prophase (Tables S5 and S6 (da Cruz et al. 2016; Margolin et al. 2014)). This expression pattern along with the involvement of the UPS in the regulation of meiotic prophase in mouse (Rao et al. 2017) and the association of USP26 with DNA repair and homologous recombination in somatic cells (Typas et al. 2016) strongly suggest a potential role of USP26 on these processes. However, our cytological analysis of mutant spermatocytes did not reveal any obvious defect in synapsis, DSBs repair, and meiotic recombination. These results suggest a very specific function of USP26 that seems unrelated to the regulation of the meiotic prophase I.

The association of this testis-specific protein with male infertility has focused many efforts during the last decades though with controversial conclusions. Several studies have linked *USP26* mutations with male infertility (Asadpor et al. 2013; Lee et al. 2008; Li et al. 2015; Ma et al. 2016; Paduch et al. 2005; Stouffs et al. 2005; Xia et al. 2014; Zhang et al. 2007). However, our functional studies showing that USP26 is dispensable for mouse fertility are in agreement with previous works reporting a lack of association between *USP26* genetic polymorphisms and male infertility (Christensen et al. 2008; Luddi et al. 2016; Ravel et al. 2006; Ribarski et al. 2009; Shi et al. 2011; Stouffs et al. 2006; Zhang et al. 2015). Supporting our observations, it has been described a normozoospermic man carrying a nonsense mutation in *USP26* (c.882 C>A) that generates a premature STOP codon (Luddi et al. 2016). This mutation produces a truncated USP26 protein at the position 293 that consequently lacks

the catalytic Cys (Cys305). Our results provide new evidences that could shed light on the existing controversy about the relationship between *USP26* mutations and human male infertility. However, although the mouse is the most commonly used organism model in human disease research (Rosenthal and Brown 2007), we cannot exclude the idea that the functional relevance of this protein could differ between both species, showing a critical role in human that does not recapitulate in the mouse where the genetic background can also drastically modulate the observed phenotypes. Nevertheless, our working models have been developed in an outbred background (B6CBAF1/J) that more closely resembles the greater genetic variability of humans.

Given the *Usp26* restricted expression pattern to testis and the importance of the maintenance of ubiquitin homeostasis in cells, the absence of phenotype in the USP26-deficient mice is counterintuitive. Genetic redundancy or biological robustness can be responsible for this lack of deviation in the reproductive phenotype. The in silico analysis of mRNA expression of different USPs during mouse spermatogenesis showed several USPs with an expression pattern highly similar to that of USP26 (i.e., USP9X, USP37, USP14, USP53, USP33, USP28, and USP3. See Tables S5 and S6 (da Cruz et al. 2016; Margolin et al. 2014)). It would be interesting to analyze which of these putative candidates (if any) are upregulated in the absence of USP26 that would provide an indication for compensation. To prove it, a double loss-of-function analysis should be carried out in future studies.

In conclusion, this work reports for the first time a loss-of-function analysis of USP26 in mice and represents in vivo evidence that USP26 is not essential for mouse gametogenesis and fertility.

Acknowledgements We wish to express our sincere thanks to Dr. Yi-Wen Lin (Institute of Biomedical Sciences, Taipei, Taiwan) for providing the anti-USP26-M antibody. We appreciate to Rodrigo García-Valiente for his help with bioinformatic analysis of RNAseq datasets.

Author contributions NFM with the help of LGH and YBC performed the characterization of the mutant mice including the cytological and histological analysis. MSM carried out the Cas9 injections. JLB provided the α SYCP3 serum K921 and reviewed the manuscript. IR carried out infertility phenotyping of mutant mice. AMP and ELC designed the experiments and helped NFM to write the manuscript with the input of the remaining authors.

Funding information This work was supported by BFU2017-89408-R. NFM and LGH are supported by European Social Fund/JCyLe grants (EDU/310/2015 and EDU/1083/2013) and YBC by a FPI grant from the MINECO. IBMCC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia cofunded by the Castilla–León autonomous government and the European Regional Development Fund (CLC–2017–01). The IBMCC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia cofunded by the Castilla–León autonomous government and the European Regional Development Fund (CLC–2017–01).

References

- Asadpor U, Totonchi M, Sabbaghian M, Hoseinifar H, Akhound MR, Zari Moradi S, Haratian K, Sadighi Gilani MA, Gourabi H, Mohseni Meybodi A (2013) Ubiquitin-specific protease (USP26) gene alterations associated with male infertility and recurrent pregnancy loss (RPL) in Iranian infertile patients. *J Assist Reprod Genet* 30:923–931. <https://doi.org/10.1007/s10815-013-0027-9>
- Baarends WM, van der Laan R, Grootegoed JA (2000) Specific aspects of the ubiquitin system in spermatogenesis. *J Endocrinol Investig* 23:597–604. <https://doi.org/10.1007/BF03343782>
- Bedard N, Yang Y, Gregory M, Cyr DG, Suzuki J, Yu X, Chian RC, Hermo L, O'Flaherty C, Smith CE, Clarke HJ, Wing SS (2011) Mice lacking the USP2 deubiquitinating enzyme have severe male subfertility associated with defects in fertilization and sperm motility. *Biol Reprod* 85:594–604. <https://doi.org/10.1095/biolreprod.110.088542>
- Bekker-Jensen S, Mailand N (2011) The ubiquitin- and SUMO-dependent signaling response to DNA double-strand breaks. *FEBS Lett* 585:2914–2919. <https://doi.org/10.1016/j.febslet.2011.05.056>
- Bolcun-Filas E, Schimenti JC (2012) Genetics of meiosis and recombination in mice. *Int Rev Cell Mol Biol* 298:179–227. <https://doi.org/10.1016/B978-0-12-394309-5.00005-5>
- Bose R, Manku G, Culty M, Wing SS (2014) Ubiquitin-proteasome system in spermatogenesis. *Adv Exp Med Biol* 759:181–213. https://doi.org/10.1007/978-1-4939-0817-2_9
- Caburet S, Arboleda VA, Llano E, Overbeek PA, Barbero JL, Oka K, Harrison W, Vaiman D, Ben-Neriah Z, García-Tuñón I, Fellous M, Pendás AM, Velitia RA, Vilain E (2014) Mutant cohesin in premature ovarian failure. *N Engl J Med* 370:943–949. <https://doi.org/10.1056/NEJMoal309635>
- Christensen GL, Griffin J, Carrell DT (2008) Sequence analysis of the X-linked USP26 gene in severe male factor infertility patients and fertile controls. *Fertil Steril* 90:851–852. <https://doi.org/10.1016/j.fertnstert.2007.06.096>
- da Cruz I, Rodríguez-Casuriaga R, Santiñaque FF, Farias J, Curti G, Caporano CA, Folle GA, Benavente R, Sotelo-Silveira JR, Geisinger A (2016) Transcriptome analysis of highly purified mouse spermatogenic cell populations: gene expression signatures switch from meiotic to postmeiotic-related processes at pachytene stage. *BMC Genomics* 17:294. <https://doi.org/10.1186/s12864-016-2618-1>
- Davis MI, Simeonov A (2015) Ubiquitin-specific proteases as druggable targets. *Drug Target Rev* 2:60–64
- Ferlin A, Raicu F, Gatta V, Zuccarello D, Palka G, Foresta C (2007) Male infertility: role of genetic background. *Reprod BioMed Online* 14:734–745
- Gomez HL et al (2016) C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nat Commun* 7:13298. <https://doi.org/10.1038/ncomms13298>
- Guillon H, Baudat F, Grey C, Liskay RM, de Massy B (2005) Crossover and noncrossover pathways in mouse meiosis. *Mol Cell* 20:563–573. <https://doi.org/10.1016/j.molcel.2005.09.021>
- Hamada AJ, Esteves SC, Agarwal A (2013) A comprehensive review of genetics and genetic testing in azoospermia. *Clinics (Sao Paulo)* 68(Suppl 1):39–60
- He M, Zhou Z, Shah AA, Zou H, Tao J, Chen Q, Wan Y (2016) The emerging role of deubiquitinating enzymes in genomic integrity, diseases, and therapeutics. *Cell Biosci* 6:62. <https://doi.org/10.1186/s13578-016-0127-1>
- Hu Z, Xia Y, Guo X, Dai J, Li HG, Hu H, Jiang Y, Lu F, Wu Y, Yang X, Li H, Yao B, Lu C, Xiong C, Li Z, Gui Y, Liu J, Zhou Z, Shen H, Wang X, Sha J (2011) A genome-wide association study in Chinese men identifies three risk loci for non-obstructive azoospermia. *Nat Genet* 44:183–186. <https://doi.org/10.1038/ng.1040>
- Kishi K, Uchida A, Takase HM, Suzuki H, Kurohmaru M, Tsunekawa N, Kanai-Azuma M, Wood SA, Kanai Y (2017) Spermatogonial deubiquitinase USP9X is essential for proper spermatogenesis in mice. *Reproduction* 154:135–143. <https://doi.org/10.1530/REP-17-0184>
- Komander D, Clague MJ, Urbe S (2009) Breaking the chains: structure and function of the deubiquitinases. *Nat Rev Mol Cell Biol* 10:550–563. <https://doi.org/10.1038/nrm2731>
- Kwon J, Mochida K, Wang YL, Sekiguchi S, Sankai T, Aoki S, Ogura A, Yoshikawa Y, Wada K (2005) Ubiquitin C-terminal hydrolase L-1 is essential for the early apoptotic wave of germinal cells and for sperm quality control during spermatogenesis. *Biol Reprod* 73:29–35. <https://doi.org/10.1095/biolreprod.104.037077>
- Lee IW, Kuan LC, Lin CH, Pan HA, Hsu CC, Tsai YC, Kuo PL, Teng YN (2008) Association of USP26 haplotypes in men in Taiwan, China with severe spermatogenic defect. *Asian J Androl* 10:896–904. <https://doi.org/10.1111/j.1745-7262.2008.00439.x>
- Li Z, Huang Y, Li H, Hu J, Liu X, Jiang T, Sun G, Tang A, Sun X, Qian W, Zeng Y, Xie J, Zhao W, Xu Y, He T, Dong C, Liu Q, Mou L, Lu J, Lin Z, Wu S, Gao S, Guo G, Feng Q, Li Y, Zhang X, Wang J, Yang H, Wang J, Xiong C, Cai Z, Gui Y (2015) Excess of rare variants in genes that are key epigenetic regulators of spermatogenesis in the patients with non-obstructive azoospermia. *Sci Rep* 5:8785. <https://doi.org/10.1038/srep08785>
- Lin YW, Hsu TH, Yen PH (2011) Localization of ubiquitin specific protease 26 at blood-testis barrier and near Sertoli cell-germ cell interface in mouse testes. *Int J Androl* 34:e368–e377. <https://doi.org/10.1111/j.1365-2605.2010.01130.x>
- Liu YL, Zheng J, Mi YJ, Zhao J, Tian QB (2018) The impacts of nineteen mutations on the enzymatic activity of USP26. *Gene* 641:292–296. <https://doi.org/10.1016/j.gene.2017.10.074>
- Luddi A, Crifasi L, Quagliarello A, Governini L, De Leo V, Piomboni P (2016) Single nucleotide polymorphisms of USP26 in azoospermic men. *Syst Biol Reprod Med* 62:372–378. <https://doi.org/10.1080/19396368.2016.1238116>
- Luo M, Zhou J, Leu NA, Abreu CM, Wang J, Anguera MC, de Rooij DG, Jasin M, Wang PJ (2015) Polycomb protein SCML2 associates with USP7 and counteracts histone H2A ubiquitination in the XY chromatin during male meiosis. *PLoS Genet* 11:e1004954. <https://doi.org/10.1371/journal.pgen.1004954>
- Ma Q, Li Y, Guo H, Li C, Chen J, Luo M, Jiang Z, Li H, Gui Y (2016) A novel missense mutation in USP26 gene is associated with nonobstructive azoospermia. *Reprod Sci* 23:1434–1441. <https://doi.org/10.1177/1933719116641758>
- Makino S, Fukumura R, Gondo Y (2016) Illegitimate translation causes unexpected gene expression from on-target out-of-frame alleles created by CRISPR-Cas9. *Sci Rep* 6:39608. <https://doi.org/10.1038/srep39608>
- Margolin G, Khil PP, Kim J, Bellani MA, Camerini-Otero RD (2014) Integrated transcriptome analysis of mouse spermatogenesis. *BMC Genomics* 15:39. <https://doi.org/10.1186/1471-2164-15-39>
- Matzuk MM, Lamb DJ (2008) The biology of infertility: research advances and clinical challenges. *Nat Med* 14:1197–1213. <https://doi.org/10.1038/nm.f1895>
- Muratori M, Marchiani S, Forti G, Baldi E (2005) Sperm ubiquitination positively correlates to normal morphology in human semen. *Hum Reprod* 20:1035–1043. <https://doi.org/10.1093/humrep/deh678>
- Nijman SM, Luna-Vargas MP, Velds A, Brummelkamp TR, Dirac AM, Sixma TK, Bernards R (2005) A genomic and functional inventory of deubiquitinating enzymes. *Cell* 123:773–786. <https://doi.org/10.1016/j.cell.2005.11.007>
- Nishimune Y, Tanaka H (2006) Infertility caused by polymorphisms or mutations in spermatogenesis-specific genes. *J Androl* 27:326–334. <https://doi.org/10.2164/jandrol.05162>
- Ombelet W, Cooke I, Dyer S, Serour G, Devroey P (2008) Infertility and the provision of infertility medical services in developing countries.

- Hum Reprod Update 14:605–621. <https://doi.org/10.1093/humupd/dmn042>
- Paduch DA, Mielnik A, Schlegel PN (2005) Novel mutations in testis-specific ubiquitin protease 26 gene may cause male infertility and hypogonadism. *Reprod BioMed Online* 10:747–754
- Prieto I, Tease C, Pezzi N, Buesa JM, Ortega S, Kremer L, Martínez A, Martínez-A C, Hultén MA, Barbero JL (2004) Cohesin component dynamics during meiotic prophase I in mammalian oocytes. *Chromosom Res* 12:197–213
- Rao HB et al (2017) A SUMO-ubiquitin relay recruits proteasomes to chromosome axes to regulate meiotic recombination. *Science* 355:403–407. <https://doi.org/10.1126/science.aaf6407>
- Ravel C, el Houate B, Chantot S, Lourenço D, Dumaine A, Rouba H, Bandyopadhyay A, Radhakrishna U, Das B, Sengupta S, Mandelbaum J, Siffroi JP, McElreavey K (2006) Haplotypes, mutations and male fertility: the story of the testis-specific ubiquitin protease USP26. *Mol Hum Reprod* 12:643–646. <https://doi.org/10.1093/molehr/gal063>
- Reijo R, Alagappan RK, Patrizio P, Page DC (1996) Severe oligozoospermia resulting from deletions of azoospermia factor gene on Y chromosome. *Lancet* 347:1290–1293
- Ribarski I, Lehar O, Yogev L, Hauser R, Bar-Shira Maymon B, Botchan A, Paz G, Yavetz H, Kleiman SE (2009) USP26 gene variations in fertile and infertile men. *Hum Reprod* 24:477–484. <https://doi.org/10.1093/humrep/den374>
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM (1998) DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* 273:5858–5868
- Rosenthal N, Brown S (2007) The mouse ascending: perspectives for human-disease models. *Nat Cell Biol* 9:993–999. <https://doi.org/10.1038/ncb437>
- Sharlip ID, Jarow JP, Belker AM, Lipshultz LI, Sigman M, Thomas AJ, Schlegel PN, Howards SS, Nehra A, Damewood MD, Overstreet JW, Sadovsky R (2002) Best practice policies for male infertility. *Fertil Steril* 77:873–882
- Shi YC, Wei L, Cui YX, Shang XJ, Wang HY, Xia XY, Zhou YC, Li H, Jiang HT, Zhu WM, Huang YF (2011) Association between ubiquitin-specific protease USP26 polymorphism and male infertility in Chinese men. *Clin Chim Acta* 412:545–549. <https://doi.org/10.1016/j.cca.2010.12.005>
- Singh P, Schimenti JC, Bolcun-Filas E (2015) A mouse geneticist's practical guide to CRISPR applications. *Genetics* 199:1–15. <https://doi.org/10.1534/genetics.114.169771>
- Stouffs K, Lissens W, Tournaye H, Van Steirteghem A, Liebaers I (2005) Possible role of USP26 in patients with severely impaired spermatogenesis. *Eur J Hum Genet* 13:336–340. <https://doi.org/10.1038/sj.ejhg.5201335>
- Stouffs K, Lissens W, Tournaye H, Van Steirteghem A, Liebaers I (2006) Alterations of the USP26 gene in Caucasian men. *Int J Androl* 29:614–617. <https://doi.org/10.1111/j.1365-2605.2006.00708.x>
- Typas D, Luijsterburg MS, Wiegant WW, Diakatou M, Helfricht A, Thijssen PE, van den Broek B, Mullenders LH, van Attikum H (2016) The de-ubiquitylating enzymes USP26 and USP37 regulate homologous recombination by counteracting RAP80. *Nucleic Acids Res* 44:2976. <https://doi.org/10.1093/nar/gkv1480>
- Wang PJ, McCarrey JR, Yang F, Page DC (2001) An abundance of X-linked genes expressed in spermatogonia. *Nat Genet* 27:422–426. <https://doi.org/10.1038/86927>
- Wang PJ, Page DC, McCarrey JR (2005) Differential expression of sex-linked and autosomal germ-cell-specific genes during spermatogenesis in the mouse. *Hum Mol Genet* 14:2911–2918. <https://doi.org/10.1093/hmg/ddi322>
- Wosnitzer MS, Mielnik A, Dabaja A, Robinson B, Schlegel PN, Paduch DA (2014) Ubiquitin specific protease 26 (USP26) expression analysis in human testicular and extragonadal tissues indicates diverse action of USP26 in cell differentiation and tumorigenesis. *PLoS One* 9:e98638. <https://doi.org/10.1371/journal.pone.0098638>
- Xia JD, Chen J, Han YF, Chen H, Yu W, Chen Y, Dai YT (2014) Association of 370-371insACA, 494T>C, and 1423C>T haplotype in ubiquitin-specific protease 26 gene and male infertility: a meta-analysis. *Asian J Androl* 16:720–724. <https://doi.org/10.4103/1008-682X.129134>
- Yi YJ, Manandhar G, Oko RJ, Breed WG, Sutovsky P (2007) Mechanism of sperm-zona pellucida penetration during mammalian fertilization: 26S proteasome as a candidate egg coat lysin. *Soc Reprod Fertil Suppl* 63:385–408
- Zhang J, Qiu SD, Li SB, Zhou DX, Tian H, Huo YW, Ge L, Zhang QY (2007) Novel mutations in ubiquitin-specific protease 26 gene might cause spermatogenesis impairment and male infertility. *Asian J Androl* 9:809–814. <https://doi.org/10.1111/j.1745-7262.2007.00305.x>
- Zhang W, Liu T, Mi YJ, Yue LD, Wang JM, Liu DW, Yan J, Tian QB (2015) Evidence from enzymatic and meta-analyses does not support a direct association between USP26 gene variants and male infertility. *Andrology* 3:271–279. <https://doi.org/10.1111/andr.295>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

A missense in HSF2BP causing primary ovarian insufficiency affects meiotic recombination by its novel interactor C19ORF57/BRME1

Natalia Felipe-Medina^{1†}, Sandrine Caburet^{2,3†}, Fernando Sánchez-Sáez¹, Yazmine B Condezo¹, Dirk G de Rooij⁴, Laura Gómez-H¹, Rodrigo García-Valiente¹, Anne Laure Todeschini^{2,3}, Paloma Duque¹, Manuel Adolfo Sánchez-Martin^{5,6}, Stavit A Shalev^{7,8}, Elena Llano^{1,9}, Reiner A Veitia^{2,3,10*}, Alberto M Pendás^{1*}

¹Molecular Mechanisms Program, Centro de Investigación del Cáncer and Instituto de Biología Molecular y Celular del Cáncer (CSIC-Universidad de Salamanca), Salamanca, Spain; ²Université de Paris, Paris Cedex, France; ³Institut Jacques Monod, Université de Paris, Paris, France; ⁴Reproductive Biology Group, Division of Developmental Biology, Department of Biology, Faculty of Science, Utrecht University, Utrecht, Netherlands; ⁵Transgenic Facility, Nucleus platform, Universidad de Salamanca, Salamanca, Spain; ⁶Departamento de Medicina, Universidad de Salamanca, Salamanca, Spain; ⁷The Genetic Institute, "Emek" Medical Center, Afula, Israel; ⁸Bruce and Ruth Rappaport Faculty of Medicine, Technion, Haifa, Israel; ⁹Departamento de Fisiología y Farmacología, Universidad de Salamanca, Salamanca, Spain; ¹⁰Université Paris-Saclay, Institut de Biologie F. Jacob, Commissariat à l'Energie Atomique, Fontenay aux Roses, France

*For correspondence:

reiner.veitia@ijm.fr (RAV);
amp@usal.es (AMP)

†These authors contributed equally to this work

Competing interests: The authors declare that no competing interests exist.

Funding: See page 26

Received: 17 March 2020

Accepted: 26 August 2020

Published: 26 August 2020

Reviewing editor: Bernard de Massy, CNRS UM, France

© Copyright Felipe-Medina et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Abstract Primary Ovarian Insufficiency (POI) is a major cause of infertility, but its etiology remains poorly understood. Using whole-exome sequencing in a family with three cases of POI, we identified the candidate missense variant S167L in *HSF2BP*, an essential meiotic gene. Functional analysis of the HSF2BP-S167L variant in mouse showed that it behaves as a hypomorphic allele compared to a new loss-of-function (knock-out) mouse model. *Hsf2bp*^{S167L/S167L} females show reduced fertility with smaller litter sizes. To obtain mechanistic insights, we identified C19ORF57/BRME1 as a strong interactor and stabilizer of HSF2BP and showed that the BRME1/HSF2BP protein complex co-immunoprecipitates with BRCA2, RAD51, RPA and PALB2. Meiocytes bearing the HSF2BP-S167L variant showed a strongly decreased staining of both HSF2BP and BRME1 at the recombination nodules and a reduced number of the foci formed by the recombinases RAD51/DMC1, thus leading to a lower frequency of crossovers. Our results provide insights into the molecular mechanism of HSF2BP-S167L in human ovarian insufficiency and sub(in)fertility.

Introduction

The process of gametogenesis is one of the most complex and highly regulated differentiation programs. It involves a unique reductional cell division, known as meiosis, to generate highly specialized cells: the gametes. Indeed, the outcome of meiosis is the production of oocytes and spermatozoa, which are the most distinctive cells of an adult organism and are essential for the faithful transmission of the genome across generations.

The meiotic division is an orderly process that results in the pairing and synapsis of homologous chromosomes and crossover (CO) formation, which ultimately enable homologous chromosomes segregation (Hunter, 2015; Loidl, 2016; Zickler and Kleckner, 2015). In mammals, pairing of homologs is dependent on the repair of self-induced double-strand breaks (DSBs) during prophase I by homologous recombination (Handel and Schimenti, 2010) and it leads to the intimate alignment of homologous chromosomes (synapsis) through the zipper-like synaptonemal complex (SC) (Cahoon and Hawley, 2016). The SC is a proteinaceous tripartite structure that provides the structural framework for DSBs repair (Baudat et al., 2013), as epitomized by the tight association of the recombination nodules (RNs, multicomponent recombinogenic factories) and the axial elements of the SC (Zickler and Kleckner, 2015).

Meiotic DSBs repair is an evolutionarily conserved pathway that is highly regulated to promote the formation of at least one CO per bivalent. This chromosome connection between bivalents through chiasmata is required for a correct reductional division. As other DNA repair processes, proper meiotic recombination is essential for genome stability and alterations can result in infertility, miscarriage and birth defects (Geisinger and Benavente, 2017; Handel and Schimenti, 2010; Webster and Schuh, 2017).

Infertility refers to failure of a couple to reproduce and affects 10–15% of couples (Isaksson and Tiitinen, 2004). Infertility can be due to female factors, male factors, a combination of both or to unknown causes, each category representing approximately 25% of cases (Isaksson and Tiitinen, 2004; Matzuk and Lamb, 2008). There are several underlying causes and physiological, genetic and even environmental and social factors can play a role. Forward and reverse genetic analyses in model organisms have identified multiple molecular pathways that regulate fertility and have allowed to infer reasonable estimates of the number of protein-coding genes essential for fertility (de Rooij and de Boer, 2003; Schimenti and Handel, 2018).

Primary ovarian insufficiency (POI) is a major cause of female infertility and affects about 1–3% of women under 40 years of age. It is characterized by cessation of ovarian function before the age of 40 years. POI results from a depletion of the ovarian follicle pool and can be isolated or syndromic. Genetic causes of POI account for approximately 20% of cases (Rossetti et al., 2017). Although infertility-causing pathogenic variants are inherently unlikely to spread in a population, they can be observed within families, especially when there is consanguinity. Such cases provide crucial insights into the function of the genes and molecular mechanisms that they disrupt. Over the last decade, causative variants in several genes have been found using whole exome sequencing in ‘POI pedigrees’. In particular, pathogenic variants in genes involved in DNA replication, recombination or repair, such as *STAG3*, *SYCE1*, *HFM1*, *MSH5* and *MEIOB* have been formally implicated in this condition by ourselves and others (Caburet et al., 2014; Caburet et al., 2019a; de Vries et al., 2014; Guo et al., 2017; Primary Ovarian Insufficiency Collaboration et al., 2014).

In this study, we have identified in a consanguineous family with POI the candidate S167L missense variant in *HSF2BP*, an essential yet poorly studied meiotic gene. *HSF2BP* encodes an interactor of the heat-shock response transcription factor HSF2 (Yoshima et al., 1998). During the course of this work and, in agreement with our results, two independent groups showed that HSF2BP is essential for meiotic recombination through its ability to interact with BRCA2 (Brandtsma et al., 2019; Zhang et al., 2019). Here, we report that the introduction of the missense variant HSF2BP-S167L in mouse leads to subfertility and DNA repair defects during prophase I. In addition, we identified a protein complex composed of BRCA2, HSF2BP, and the as yet unexplored C19ORF57/BRME1 (meiotic double-stranded break BRCA2/HSF2BP complex associated protein) as a key component of the meiotic recombination machinery. Our studies show that a single substitution (S167L) in HSF2BP leads to a reduced loading of both BRME1 and HSF2BP at the RNs. Furthermore, our results suggest that meiotic progression requires a critical threshold level of HSF2BP/BRME1 for the ulterior loading of the recombinases to the RNs.

Results

Clinical cases

The parents are first-degree cousins of Israeli Arab origin. Of the five daughters, three are affected with POI and presented with early secondary amenorrhea. They had menarche at normal age (at 13–

14) but with irregular menses that stopped around 25. Only one of the patients affected by POI could have a child with the help of a fertility treatment (see pedigree in **Figure 1**). In order to identify the genetic basis of this familial POI case, we performed whole exome sequencing on genomic DNA from two POI patients, III-2 and III-3, and their fertile sister III-10 (**Supplementary file 1a**). Variants were filtered on the basis of (i) their homozygosity in the patients, (ii) their heterozygosity or absence in the fertile sister, (iii) their absence in unrelated fertile in-house controls and (iv) a minor allele frequency (MAF) below 0.01 in all available databases (**Supplementary file 1b**). This filtering process led to the identification of a missense substitution located in the *HSF2BP* gene: rs200655253 (21:43630396 G > A, GRCh38). The variant lies within the sixth exon of the reference transcript ENST00000291560.7 (NM_007031.2:c.500C > T) and changes a TCG codon into a TTG (NP_008962.1:p.Ser167Leu). It is very rare (Variant Allele Frequency/VAF 0.0001845 in the GnomAD database and 0.0005 in the GME Variome dedicated to Middle-East populations) and absent in a

Figure 1. Pedigree of the consanguineous family with the variant *HSF2BP*-S167L. III-1 and III-2 are monozygotic twins, who appear phenotypically dizygotic. Clinical investigation confirmed POI, with normal 46, XX karyotype (500 bands and SKY spectral karyotyping). Year of birth and age of menarche are indicated when known. III-1 became amenorrheic at age 24 and III-2 at age 25, both after irregular menstruations since menarche. III-1 presents with a short stature (152 cm, within the 3–5 percentile), a normal neck, cubitus valgus and metacarpal shortening of 4–5. Ultrasound investigation showed normal uterus and ovaries. Her g-banding karyotyping was normal 46, XX (500 bands) and variants in *FMR1* gene were ruled out. III-2 displays a normal secondary sexual development with no dysmorphic sign. Clinical investigation confirmed POI, with normal 46, XX karyotype (500 bands and SKY spectral karyotyping). The elder sister III-3 was also diagnosed with POI, with no further clinical information. She is 160.5 cm. She had one normal pregnancy with the help of ‘fertility treatment’, and a second unsuccessful attempt. The two fertile sisters, III-6 and III-10 had their menarche at 14–15 and 13–14 respectively, with regular menstruations ever since. They are respectively 150 cm and 151 cm, with no clinical sign, and each one had several children without difficulties. The fertile brother III-7 is 171 cm and shows no health or fertility problem. He developed frontal baldness since the age of 30. The genotype of each individual at the variant genomic position in *HSF2BP* is shown in red, as determined by Sanger sequencing for available DNAs (See **Figure 1—figure supplement 1**).

The online version of this article includes the following figure supplement(s) for figure 1:

Figure supplement 1. Segregation of the S167L Variant in *HSF2BP* in the consanguineous family shows the chromatograms obtained by Sanger sequencing of the *HSF2BP*-S167L variant in the family.

Figure supplement 2. Strong conservation of the Ser167 residue in *HSF2BP* protein in 99 mammals.

Figure supplement 3. Strong conservation of the Ser167 residue in *HSF2BP* in 48 birds and reptiles and 64 fish species.

homozygous state from all available databases. The variant was verified by Sanger sequencing and was found to segregate in a Mendelian fashion within the family: the affected twin III-1 was homozygous for the variant and both parents and fertile siblings were heterozygous carriers (**Figure 1—figure supplement 1**). Therefore, there was no homozygous males identified in this family, preventing the analysis of the impact of this variant on male fertility. Serine 167 is a highly conserved position and the S167L variant is predicted to be pathogenic or deleterious by 11 out of the 18 pathogenicity predictors available in dbNSFP 3.5. (**Supplementary file 1c**, **Figure 1—figure supplement 2** and **Figure 1—figure supplement 3**).

Mice with the HSF2BP S167L variant show a partial reduction of fertility

During the course of this work, two independent groups showed that HSF2BP is essential for meiotic recombination through its ability to interact with the armadillo repeats of BRCA2 (**Zhang et al., 2019**). Both groups showed that genetic disruption of *Hsf2bp* in mouse leads to the accumulation in the chromosomes axes of DNA repair proteins such as γ H2AX (ATR-dependent phosphorylation of H2AX marks DSBs) and the single stranded-DNA binding protein RPA, a strong reduction of the recombinases DMC1 and RAD51 at the RNs and a lack of COs as labelled by MLH1 (**Baker et al., 1996**). The end result is male sterility (**Brandsma, 2006; Brandsma et al., 2019; Zhang et al., 2019**). However, loss of HSF2BP in female mice showed a milder meiotic phenotype (**Zhang et al., 2019** and our own data, see below) and a weak albeit non-statistical significant reduction of fertility (**Brandsma et al., 2019**) despite all of the mutants are nulls though in different genetic backgrounds.

In order to confirm the causality of the S167L variant in this POI family, we generated a knock-in mouse *Hsf2bp*^{S167L/S167L} by genome editing (**Figure 2—figure supplement 1a**). We also generated a loss-of-function model (*Hsf2bp*^{-/-}) for direct comparison (**Figure 2—figure supplement 1b–d**). *Hsf2bp*^{S167L/S167L} male and female mice were able to reproduce but females showed a significant reduction in the number of litters (**Figure 2a**), whilst males only showed a slight non-significant reduction in fertility (**Figure 2a**), suggesting that the S167L variant impacts murine fertility.

Histological analysis of *Hsf2bp*^{S167L/S167L} ovaries revealed no apparent differences in the number of follicles in comparison to wild-type (WT) animals (**Figure 2b–c** and **Figure 2—figure supplement 2a**), in contrast with the drastic reduction of the follicle pool in *Hsf2bp*^{-/-} ovaries (**Figure 2b–c**). Testes from *Hsf2bp*^{S167L/S167L} mice displayed a reduced size (21% reduction compared to WT mice; testis/body weight ratio: S167L 0.26% $\pm$ 0.07 (n = 12) vs 0.33% $\pm$ 0.05 for WT controls (n = 14), **p<0.01, **Figure 2d** and **Figure 2—figure supplement 2b**) and this reduction was stronger in *Hsf2bp*^{-/-} testes (70% reduction compared to WT, testis/body weight ratio: *Hsf2bp*^{-/-} 0.10% $\pm$ 0.005 (n = 6) vs 0.33% $\pm$ 0.05 for WT controls (n = 14) ****p<0.001, **Figure 2d** and **Figure 2—figure supplement 2b**). Histological analysis of adult *Hsf2bp*^{S167L/S167L} testes revealed seminiferous tubules with a partial arrest with apoptotic spermatocytes (meiotic divisions) and their epididymis exhibited scarcer spermatozoa (**Figure 2e**). Consistent with these results, *Hsf2bp*^{S167L/S167L} males showed increased numbers of meiotic divisions positive for TUNEL staining (**Figure 2f**) and a reduction in the number of spermatozoa in the epididymis (3.3×10^6 in the *Hsf2bp*^{S167L/S167L} mutant vs 4.3×10^6 in the WT; **Figure 2g**). During mouse spermatogenesis, the 12 stages of the epithelial cycle can be distinguished in seminiferous tubule sections by identifying groups of associated germ cell types (**Ahmed and de Rooij, 2009**). Following these criteria, the seminiferous epithelium of *Hsf2bp*^{-/-} mice showed a stage IV arrest, characterized by a massive apoptosis of zygotene-like spermatocytes occurring at the same time that In spermatogonia divide into B spermatogonia (**Figure 2e**). The presence of spermatogonia, spermatocytes, Sertoli and Leydig cells was not altered in any of the mutants (**Figure 2e**). These results suggest that mice bearing the POI-causing variant only partially phenocopy the human disease.

Hsf2bp^{S167L/S167L} meocytes show an altered meiotic homologous recombination

To further characterize meiotic defects, *Hsf2bp*^{S167L/S167L} meocytes were first analyzed for the assembly/disassembly of the SC by monitoring the distribution of SYCP1 and SYCP3. We did not observe any difference in synapsis and desynapsis from leptotene to diplotene in both oocytes and

Figure 2. Mice carrying the HSF2BP S167L variant show a partial reduction of fertility. (a) Fertility assessment of males and female *Hsf2bp*^{S167L/S167L} and WT mice showing the number of litters per month and the number of pups per litter (see Materials and methods). Mice: *Hsf2bp*^{+/+} n = 6 females/6 males, *Hsf2bp*^{S167L/S167L} n = 7 females/6 males. Two-tailed Welch's t-test analysis: *p < 0.05. (b) Hematoxylin and eosin stained sections of ovaries from adult (8 weeks) *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} females. *Hsf2bp*^{-/-} ovaries but not *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L} showed a strong depletion

Figure 2 continued on next page

Figure 2 continued

of follicles. Bar in panels 100 μm . (c) Quantification of the number of follicles (primordial, primary, secondary and antral follicles) per ovary in *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} females at 5 dpp and 6 weeks of age showing no differences between *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L} but a strong reduction in the oocyte pool in *Hsf2bp*^{-/-} females. Ovaries: five dpp/6 weeks = 5/5 ovaries from *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L} and 4/3 from *Hsf2bp*^{-/-}. Two-tailed Welch's t-test analysis: ns, no significant differences, **p<0.01, ***p<0.001. (d) Testis size of *Hsf2bp*^{S167L/S167L} (left, 21% reduction) and *Hsf2bp*^{-/-} mice (right, 70% reduction) in comparison with their WT counterparts. See **Figure 2—figure supplement 2b** for the quantification. (e) PAS and Hematoxylin stained testis sections. The S167L variant leads to a partial spermatogenic arrest with an elevated number of apoptotic meiotic divisions (blue asterisks) and a reduction of the number of spermatozoa in the epididymides in comparison with the WT control (*Hsf2bp*^{+/+}). The null allele (*Hsf2bp*^{-/-}) showed a complete spermatogenic arrest at epithelial stage IV and absence of spermatozoa. Massive apoptosis of spermatocytes is indicated (blue arrowheads). Bar: upper panels 10 μm , lower panels 20 μm . (St) Seminiferous tubules, (Ep) Epididymides. (f) Immunohistochemical detection of apoptotic cells by TUNEL staining showing an increase of apoptotic meiotic divisions in stage XII tubules from *Hsf2bp*^{S167L/S167L} males (magnified panel). Plot under the panel represents the quantification. Mice: n = 3 adult mice for each genotype. Two-tailed Welch's t-test analysis: ***p<0.001. Bar in panel, 25 μm . (g) Quantification of epididymal sperm in *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L} adult mice. Epididymides: n = 8 for each genotype. Two-tailed Welch's t-test analysis: *p<0.05.

The online version of this article includes the following figure supplement(s) for figure 2:

Figure supplement 1. Generation and genetic characterization of *Hsf2bp* S167L and *Hsf2bp*-deficient mice.

Figure supplement 2. Fertility defects in *Hsf2bp*^{S167L/S167L} mice.

spermatocytes (**Figure 3—figure supplement 1a–b**). However, we observed an elevated number of apoptotic meiotic divisions in *Hsf2bp*^{S167L/S167L} males (**Figure 2—figure supplement 2c**). These results are consistent with the partial arrest observed in the histological analysis (**Figure 2e**). As expected, this phenotype was exacerbated in *Hsf2bp*^{-/-} spermatocytes that were arrested at a zygotene-like stage (**Figure 3—figure supplement 1c**). *Hsf2bp*^{-/-} oocytes showed a delay in prophase I progression with the majority of cells at zygotene stage in 17.5 days post-coitum (dpc) females, whilst the WT oocytes were mainly at pachytene stage. Additionally, we observed increased numbers of oocytes showing synapsis defects in the *Hsf2bp*^{-/-} oocytes (*Hsf2bp*^{-/-}: 45.5% $\pm$ 1.5 vs WT: 7.5% $\pm$ 1.5; n = 2 (both genotypes), **p<0.01, **Figure 3—figure supplement 1d**). These results strongly suggest that the POI variant S167L is a hypomorphic allele.

Next, we analyzed whether the POI-inducing variant affects the loading/stability of HSF2BP by immunolabeling meiocytes from *Hsf2bp*^{S167L/S167L} mice. We observed a striking reduction of HSF2BP staining at the axes during prophase I in both spermatocytes and oocytes (**Figure 3a–b**). Western blot analysis of WT, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} in whole testis extracts from 13 days post-partum (dpp) animals (**Figure 3c**) revealed that the reduced labeling observed by immunofluorescence correlated with a reduced protein expression level, suggesting that the mutation leads to a reduced expression and/or stability.

Given that HSF2BP is essential for DNA repair, we carried out a comparative staining analysis of γ H2AX, the ssDNA-binding protein RPA, the recombinases RAD51 and DMC1, the ssDNA-binding protein SPATA22 (complexed to RPA during resection) and CO formation in meiocytes from *Hsf2bp*^{S167L/S167L}, *Hsf2bp*^{-/-} and WT animals (**Figures 4, 5 and 6, Figure 4—figure supplement 1, Figure 5—figure supplement 1 and Figure 6—figure supplement 1**). Our results revealed that *Hsf2bp*^{S167L/S167L} spermatocytes showed an increased labeling of γ H2AX at pachytene (**Figure 4a**), an accumulation of RPA at the chromosome axis (**Figure 4b and Figure 4—figure supplement 1a**), a reduction of the recombinases DMC1 and RAD51 staining (**Figure 5a–b and Figure 5—figure supplement 1a–b**), an accumulation of SPATA22 (**Figure 6a and Figure 6—figure supplement 1a**), and a decreased number of COs (measured as MLH1, **Figure 6b and Figure 6—figure supplement 1b**). In accordance with the reduction of COs, we observed the presence of univalents in the XY pair at pachynema as well as univalents in metaphase I spermatocytes from *Hsf2bp*^{S167L/S167L} mice (**Figure 6d and Figure 6—figure supplement 1c**). These results would explain the elevated number of apoptotic metaphases observed (**Figure 2e–f and Figure 2—figure supplement 2c**).

Our analysis in females showed accumulation of γ H2AX staining (**Figure 4c**) but no accumulation in RPA labeling in *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} oocytes (**Figure 4d and Figure 4—figure supplement 1b**). Similar to the spermatocytes, DMC1 and RAD51 staining showed a reduction in both *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} oocytes (**Figure 5c–d and Figure 5—figure supplement 1c–d**). SPATA22 labeling in females showed a clear accumulation in *Hsf2bp*^{-/-} but only a trend towards accumulation in *Hsf2bp*^{S167L/S167L} oocytes (**Figure 6a and Figure 6—figure supplement 1a**). In

Figure 3. Meocytes from *Hsf2bp*^{S167L/S167L} mice show a decrease in the expression of HSF2BP. (a–b) Double immunofluorescence of HSF2BP (green) and SYCP3 (red) in *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L} (a) spermatocyte and (b) oocyte spreads showing a strong reduction in the labeling of HSF2BP at the chromosome axis. Plots under the panels show the quantification. Nuclei analyzed: 30 zygonemas and 40 pachynemas from two adult male mice of each genotype. In females 38/39 zygonemas and 37/35 pachynemas from two 17.5 dpc embryos of *Hsf2bp*^{+/+} and *Hsf2bp*^{S167L/S167L}, respectively. Two-tailed Welch's t-test analysis: ****p<0.0001. (c) Western blot analysis of protein extracts from 13 dpp WT, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} testes using polyclonal antibodies against HSF2BP. Tubulin was used as loading control. Graph on the right represents the relative quantification of the immunoblotting. Mice: n = 2 *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-}. Two-tailed Welch's t-test analysis: *p<0.05. Bar in panels a–c, 10 μ m. The online version of this article includes the following figure supplement(s) for figure 3:

Figure supplement 1. *Hsf2bp*^{S167L/S167L} mice do not show synapsis defects.

agreement with the lower presence of recombinases, the number of COs (measured as interstitial CDK2 foci) was also reduced in *Hsf2bp*^{S167L/S167L} oocytes, and a stronger reduction was observed in *Hsf2bp*^{-/-} oocytes (Figure 6c and Figure 6—figure supplement 1d). Overall, male and female *Hsf2bp*^{S167L/S167L} mice share alterations in the meiotic recombination pathway although with different reproductive outcome.

We next sought to understand how the HSF2BP pathogenic variant was mediating the observed meiotic alteration. HSF2BP has been shown to bind BRCA2, an essential protein for meiotic homologous recombination (Martinez et al., 2016; Sharan et al., 2004), by a direct interaction that involves Arg200 in HSF2BP and the Gly2270-Thr2337 region within the C-terminal fragment of BRCA2

Figure 4. DNA repair in *Hsf2bp*^{S167L/S167L} mice. (a, c) Double labeling of γH2AX (green) and SYCP3 (red) in (a) spermatocyte and (c) oocyte spreads from WT, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} mice. (a) Males display an accumulation of γH2AX patches in *Hsf2bp*^{S167L/S167L} pachynemas and a strong accumulation in the whole nucleus in *Hsf2bp*^{-/-} zygotene-like arrested cells. Plots on the right of the panel represent the percentage of pachynemas with γH2AX labeling (Nuclei: 364 *Hsf2bp*^{+/+} and 376 *Hsf2bp*^{S167L/S167L} from three adult mice) and the quantification of γH2AX intensity on autosomes at Figure 4 continued on next page

Figure 4 continued

early and mid-pachytene stages (Nuclei: 53 early and 60 mid pachynemas from three adult mice of each genotype). Two-tailed Welch's t-test analysis: * $p < 0.05$, **** $p < 0.0001$. (c) In females there is an accumulation of γ H2AX in $Hsf2bp^{S167L/S167L}$ pachynemas that is stronger in those from $Hsf2bp^{-/-}$ females. Nuclei: $n = 21/20/19$ pachynemas from 2 $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$ embryos (17.5 dpc). Two-tailed Welch's t-test analysis: * $p < 0.05$, **** $p < 0.0001$. (b, d) Double immunolabeling of RPA1 (green) and SYCP3 (red) in (b) spermatocyte and (d) oocyte spreads from $Hsf2bp^{+/+}$, $Hsf2bp^{S167L/S167L}$ and $Hsf2bp^{-/-}$. (b) In males, RPA1 accumulates at early and mid-pachytene in $S167L$ spermatocytes and in the zygotene-like arrested cells from $Hsf2bp^{-/-}$. Plot on the right of the panel represents the quantification. Nuclei: $n = 31/34/37$ leptotemas, $n = 29/29/37$ zygonemas/zygonemas-like from three adult $Hsf2bp^{+/+}$, $Hsf2bp^{S167L/S167L}$ and $Hsf2bp^{-/-}$ mice respectively, $n = 33$ early and 46 mid pachynemas from three adult $Hsf2bp^{+/+}$ and $Hsf2bp^{S167L/S167L}$ mice. Two-tailed Welch's t-test analysis: ns, no significant differences; ** $p < 0.01$, **** $p < 0.0001$. (d) In females, RPA1 labeling is similar in $Hsf2bp^{+/+}$, $Hsf2bp^{S167L/S167L}$ and $Hsf2bp^{-/-}$ oocytes at zygotene and pachytene. Plot on the right of the panel represents the quantification. Nuclei: $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$ $n = 42/41/23$ zygonemas from two embryos (16.5 dpc) and $n = 25/25/24$ pachynemas from two embryos (17.5 dpc). Two-tailed Welch's t-test analysis: ns, no significant differences. Bar in all panels, 10 μ m. Extended panels for RPA1 figures in **Figure 4—figure supplement 1**.

The online version of this article includes the following figure supplement(s) for figure 4:

Figure supplement 1. RPA localization in *Hsf2bp* mutants.

(Brandsma et al., 2019). Given the impossibility to detect endogenous BRCA2 by immunofluorescence in mouse spermatocytes, we carried out co-localization/interaction assays in a heterologous system by transfecting BRCA2-C (i.e. its C-term) and HSF2BP in U2OS/HEK293T. Our results showed that BRCA2-C co-immunoprecipitates with both HSF2BP-WT and HSF2BP-S167L in similar ways (**Figure 6—figure supplement 2a**). In single transfections, HSF2BP localized in the nucleus and cytoplasm whereas BRCA2-C showed nuclear localization. This pattern changed drastically to a nuclear dotted pattern when co-transfected (**Figure 6—figure supplement 2b**). This re-localization was independent of the HSF2BP variant, suggesting that the HSF2BP variant effects are not directly mediated by BRCA2 delocalization.

BRME1, a novel interactor of HSF2BP

In order to further understand the mechanism underlying the pathogenicity of the HSF2BP-S167L variant, we searched for proteins that interact with the murine HSF2BP through a yeast two hybrid (Y2H) screening. The analysis of the clones with putative interactors revealed that 19 out of 98 analyzed clones matched the uncharacterized gene 4930432K21Rik, which corresponds to human C19ORF57, hereby dubbed BRME1 for Break Repair Meiotic recombination recruitment factor 1. This HSF2BP interactor consists of 600 amino acids with a high content of acidic residues, has no recognizable functional domains and is intrinsically disordered. The interaction was validated by transiently transfecting plasmids driving the expression of HSF2BP and BRME1. Both HSF2BP-S167L and WT interacted with BRME1 (**Figure 7a**). We further validated this interaction in vivo by co-immunoprecipitation (co-IP) of both proteins from mouse whole testis extracts (**Figure 7b**). To identify the regions required for this interaction, we split the BRME1 protein into three fragments (N-terminal, central region and C-terminal). We mapped the HSF2BP/BRME1-interacting domain to the C-term fragment of BRME1 (spanning residues 475–600 of the murine protein, **Figure 7c**). In line with this, the $Brme1^{\Delta 142-472/\Delta 142-472}$ mutant mice, expressing the BRME1 protein devoid of its central part, were fertile and did not show defects in chromosome synapsis or an alteration of HSF2BP loading to axes, further indicating that a large fraction of the coding protein of BRME1 is not essential for BRME1/HSF2BP function in vivo (**Figure 7—figure supplement 1**).

We also sought to characterize the involvement of BRME1 in meiosis through immunofluorescence. BRME1 localized to the chromosome axes of WT meiotic cells from zygotene to pachytene with a pattern of discrete foci that mimics the RNs (**Figure 7—figure supplement 2a–b**). In agreement with the yeast two hybrid and co-IP results, BRME1 perfectly co-localized with HSF2BP on the chromosome axes (**Figure 7d** and **Supplementary file 1d** for quantification). This co-localization was verified by super-resolution microscopy (**Figure 7e**). In accordance with the tight association of BRME1 with HSF2BP and with a role in DSB repair, both HSF2BP and BRME1 colocalized with RPA and DMC1 foci. During prophase I, HSF2BP and BRME1 showed higher levels of spatio-temporal colocalization at the RNs with RPA than with DMC1 (**Figure 7—figure supplement 3a–b** and **Supplementary files 1d–e** for quantification). We also analyzed the HSF2BP-dependent localization of BRME1 in $Hsf2bp^{-/-}$ and $Hsf2bp^{S167L/S167L}$ mutants. Immunofluorescence analysis of meiotic cells

Figure 5. The loading of recombinases is compromised in $Hsf2bp^{S167L/S167L}$ mice. Double immunolabeling of (a, c) DMC1 or (b, d) RAD51 (green) and SYCP3 (red) in $Hsf2bp^{+/+}$, $Hsf2bp^{S167L/S167L}$ and $Hsf2bp^{-/-}$ (a, b) spermatocytes and (c, d) oocytes showing a strong reduction ($Hsf2bp^{-/-}$) and mild reduction ($Hsf2bp^{S167L/S167L}$) in the number of foci in comparison with their WT counterparts. Plots on the right of the panels represent the quantification of foci on each genotype and stage. Male nuclei DMC1: $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$, respectively, $n = 43/38/37$ early zygonemas, 41/43/37 late zygonemas/zygonemas-like and 44/37 early pachynemas from two adult mice of each genotype. Male nuclei RAD51: $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$ respectively $n = 39$ early zygonemas from all genotypes, 37/40/43 late zygonemas/zygonema-like and 37/39 early pachynemas from two adult mice of each genotype. Oocyte nuclei DMC1: $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$ $n = 18/18/22$ zygonemas from two embryos and $n = 21/30/23$ pachynemas from two embryos (17.5 dpc). Oocyte nuclei RAD51: $Hsf2bp^{+/+}/Hsf2bp^{S167L/S167L}/Hsf2bp^{-/-}$ $n = 35/35/42$ zygonemas and $n = 42/42/40$ pachynemas from two embryos (17.5 dpc). Two-tailed Welch's t-test analysis: ns, no significant differences, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Bar in all panels, 10 μ m. Extended panels for these figures in **Figure 5—figure supplement 1**.

The online version of this article includes the following figure supplement(s) for figure 5:

Figure supplement 1. Defective loading of recombinases in $Hsf2bp^{S167L/S167L}$ mice.

showed a complete lack of BRME1 staining in the absence of HSF2BP and a significant reduction of foci number in the presence of HSF2BP-S167L variant (35% reduction in males and 72% in females at pachytene; **Figure 7f–g** and **Figure 7—figure supplement 4a**). Western blot analysis also revealed a drastic reduction of BRME1 expression in $Hsf2bp^{S167L/S167L}$ and $Hsf2bp^{-/-}$ spermatocytes, suggesting an HSF2BP-dependent stabilization of BRME1 (**Figure 7h**).

To assess if HSF2BP and/or BRME1 had DNA-binding activity (targeting to DSBs), we carried out an in vitro binding assay using HSF2BP and BRME1 proteins expressed in a transcription and translation coupled reticulocyte system (TNT; **Loregian et al., 2004; Souquet et al., 2013**) in which there

Figure 6. Recombination proficiency is decreased in *Hsf2bp*^{S167L/S167L} mice. (a) Double labeling of SPATA22 (green) and SYCP3 (red) in spermatocyte (upper panel) and oocyte (lower panel) spreads from WT, *Hsf2bp*^{-/-} and *Hsf2bp*^{S167L/S167L} mice. SPATA22 is accumulated in knock-out spermatocytes and oocytes and shows a milder accumulation in the *Hsf2bp*^{S167L/S167L} spermatocytes. *Hsf2bp*^{S167L/S167L} oocytes show a slight but not significant accumulation. Plots on the right of the panel represents the quantification of SPATA22 labeling. Males nuclei: n = 20 cells for each stage from two adult mice of each genotype. Females nuclei: *Hsf2bp*^{+/+}/*Hsf2bp*^{S167L/S167L}/*Hsf2bp*^{-/-} n = 41/40/40 zygonemas from two embryos and n = 40/39/38 pachynemas from two embryos (17.5 dpc). Two-tailed Welch's t-test analysis: ns, no significant differences, **p<0.01, ****p<0.0001. (b) Double immunofluorescence of MLH1 (green) and SYCP3 (red) in spermatocyte spreads from WT, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-}. MLH1 foci are significantly reduced in the *Hsf2bp*^{S167L/S167L} spermatocytes and absent in the knock-out. The plot on the right shows the quantification. See also **Figure 6—figure supplement 1b** for the plot showing the percentage of bivalents without CO. Nuclei: n = 61 for *Hsf2bp*^{+/+}, 89 for *Hsf2bp*^{S167L/S167L} and 60 for *Hsf2bp*^{-/-} from three adult mice of each genotype. Two-tailed Welch's t-test analysis: ****p<0.0001. (c) Double labeling of CDK2 (green) and SYCP3 (red) in oocyte spreads from 17.5 dpc *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} embryos. During meiotic prophase I, CDK2 localizes to the telomeres of chromosomes from leptotene to diplotene. However, around mid-pachytene additional interstitial CDK2 signals appear at CO sites, colocalizing with MLH1. As a measure of COs, just interstitial CDK2 foci (non-telomeric) have been counted. *Hsf2bp*^{-/-} and *Hsf2bp*^{S167L/S167L} females show a high and moderate reduction in the number of COs, respectively. Plot on the right of the panel show the quantification. See also **Figure 6—figure supplement 1d** for the plot showing the percentage of bivalents without CO. Nuclei: *Hsf2bp*^{+/+}/*Hsf2bp*^{S167L/S167L}/*Hsf2bp*^{-/-} n = 79/67/46 from three embryos (17.5 dpc). Two-tailed Welch's t-test analysis: ***p<0.0001, ****p<0.0001. (d) Double immunofluorescence of γH2AX (green) and SYCP3 (red) in spermatocyte spreads from WT and *Hsf2bp*^{S167L/S167L} mice. At pachytene, γH2AX identifies the XY bivalent. The diagram on the right represents the quantification of the pachynemas with unsynapsed sex chromosomes from *Hsf2bp*^{S167L/S167L} and WT mice. Nuclei: n = 150 pachynemas from three adult mice of each genotype. Two-tailed Welch's t-test analysis: *p<0.05. Bar in all panels, 10 μm.

The online version of this article includes the following figure supplement(s) for figure 6:

Figure supplement 1. Meiotic recombination is affected in *Hsf2bp*^{S167L/S167L} mice.

Figure supplement 2. Comparative interaction of HSF2BP-S167L and HSF2BP-WT with BRCA2.

Figure 7. BRME1, a novel HSF2BP interactor that colocalizes to the recombination nodules. (a) HEK293T cells were transfected with Flag-HSF2BP (WT, upper panel; S167L, lower panel) and its novel interactor GFP-BRME1. Protein complexes were immunoprecipitated (IP: green text) with either anti-Flag or anti-EGFP or IgGs (negative control) and analysed by immunoblotting with the indicated antibody (WB: red text). Both HSF2BP variants (WT and S167L) co-immunoprecipitated similarly with BRME1. (b) IP of testis extracts with antibodies against BRME1, HSF2BP and IgGs as a negative control (IP: Figure 7 continued on next page

Figure 7 continued

green text) and western blot with the indicated antibodies (WB: red text) (c) Schematic representation of full-length BRME1 protein and the corresponding deletion (Δ) constructs (filled boxes) generated to decipher the essential BRME1 region for interacting with HSF2BP (green positive interaction and red no interaction). Western blots under the scheme show the Co-IP experiments. HEK293T cells were transfected with GFP-HSF2BP and the different delta constructs of Flag-BRME1. The $\Delta 475$ –600 abolishes the interaction, indicating that the C terminus of BRME1 is the essential region of interaction with HSF2BP. (d) Triple immunofluorescence of BRME1 (green), HSF2BP (red) and SYCP3 (blue) in WT spermatocyte spreads showing high colocalization between BRME1 and HSF2BP at late zygotene and pachytene (See **Supplementary file 1d** for quantification). Bar in panel, 10 μ m. (e) Double immunolabeling of spermatocyte spread preparations with HSF2BP (green) and BRME1 (red) analyzed by Stimulated emission depletion (STED) microscopy. Bar in panel, 5 μ m. (f–g) Double immunofluorescence of BRME1 (green) and SYCP3 (red) in (f) spermatocytes and (g) oocyte spreads from *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} showing a strong reduction of BRME1 staining in the S167L mutant and absence in the *Hsf2bp* knock-out. Plots next to the panel represent the quantification. See also extended **Figure 7—figure supplement 4a**. Male nuclei: *Hsf2bp*^{+/+}/*Hsf2bp*^{S167L/S167L}/*Hsf2bp*^{-/-}; n = 16/16/20 early and 15/15/16 late zygonemas, 16/16/- pachynemas from two adult mice. Female nuclei: n = 18 pachynemas from two embryos (17.5 dpc) of each genotype. Two-tailed Welch's t-test analysis: *p<0.05, ****p<0.0001. Bar in panels, 10 μ m. (h) Western blot analysis of protein extracts from 13 dpp WT, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-} testes using an antibody against BRME1. Tubulin was used as loading control. Graph on the right represents the relative quantification of the immunoblotting. Mice: n = 2 *Hsf2bp*^{+/+}, *Hsf2bp*^{S167L/S167L} and *Hsf2bp*^{-/-}. Two-tailed Welch's t-test analysis: *p<0.05.

The online version of this article includes the following figure supplement(s) for figure 7:

Figure supplement 1. *Brme1* $\Delta 142$ –472 mutants do not show meiotic defects.

Figure supplement 2. C19ORF57/BRME1 localizes at meiotic RNs.

Figure supplement 3. Colocalization analysis of BRME1 and HSF2BP with RPA and DMC1.

Figure supplement 4. BRME1 localization depends on HSF2BP and none of them has DNA-binding abilities.

Figure supplement 5. Generation and genetic characterization of *Rnf212*^{-/-} and *Hei10*^{-/-} mice.

Figure supplement 6. BRME1 loading depends on DSBs generation but not on synapsis.

are no nuclear proteins and chromatin (Melton et al., 1984) and used RPA as positive control. Our results show that both proteins lacked direct DNA-binding abilities, in contrast to the strong activity of RPA (Figure 7—figure supplement 4b–c).

To determine the role of BRME1 in recombination and DNA repair, we analyzed its cytological distribution pattern in different mutants lacking synapsis/recombination-related proteins. These mutants were the meiotic cohesin REC8 (Bannister et al., 2004), the central element protein of the SC SIX6OS1 (Gómez-H et al., 2016), the E3 ligases involved in the stabilization of recombinogenic proteins RNF212 and HEI10 (Qiao et al., 2014), the spermatoproteasomal subunit PSMA8 (Gómez-H et al., 2019), and the nuclease SPO11 required for DSBs generation (*Rnf212*^{-/-}, *Hei10*^{-/-} and *Spo11*^{-/-} mouse mutants are described in this work, see Materials and methods and Figure 7—figure supplement 5 and Figure 7—figure supplement 6a, Baudat et al., 2000). HSF2BP staining was also carried out for a direct comparison. We were able to show that none of the recombination-deficient mutants abrogate BRME1 labeling at zygotene (or the corresponding meiotic stage at which the mutant spermatocytes are arrested), in contrast to its absence of loading in SPO11-deficient mice (Figure 7—figure supplement 6b, left). These results are very similar to those obtained for HSF2BP in these mutants (Figure 7—figure supplement 6b, right) and indicate that SPO11-dependent DSBs are essential for targeting HSF2BP/BRME1 to the RNs, and that the heterocomplex can be positioned at early events soon after DSBs generation.

To functionally analyze the role of BRME1 in mouse fertility, we generated a *Brme1*^{-/-} null mutant by genome editing (Figure 8—figure supplement 1a–d). *Brme1*^{-/-} females, despite being fertile, showed a strong reduction of the follicle pool (Figure 8a). Male *Brme1*^{-/-} mice were infertile, the average size of their testes was severely reduced (76% reduction compared to WT; testis weight/body weight ratio: *Brme1*^{-/-} 0.08% $\pm$ 0.004 (n = 6) vs 0.33% $\pm$ 0.05 for WT controls (n = 14), ****p<0.0001, Figure 8b and Figure 2—figure supplement 2b), and lacked spermatozoa (Figure 8c). Histological analysis showed a meiotic arrest at epithelial stage IV with apoptotic spermatocytes (Figure 8c). Double immunolabeling of SYCP3 and SYCP1 revealed that spermatocytes were partially synapsed and showed a partner-switch phenotype in which synapsis is not restricted to homologous pairs (Figure 8d). The arrest corresponds to a zygotene-like stage though a small fraction of cells (3.7% $\pm$ 1.9; n = 3) were able to escape this blockage reaching early pachytene. *Brme1*^{-/-} oocyte spread analysis revealed the presence of a subset of fully synapsed pachynemas but an increased number of cells with different degree of asynapsis (47.9% $\pm$ 2.2 vs 12% $\pm$ 5.7 in the WT; n = 2 (both genotypes), *p<0.05 Figure 8e and Figure 8—figure supplement 1e). Given the

Figure 8. *Brme1*^{-/-} mice show severe fertility defects. (a) Hematoxylin+eosin stained sections of ovaries from adult *Brme1*^{-/-} females showing a strong depletion of follicles. Plot on the right represents the quantification in 3 months-old females. Ovaries: n = 3 ovaries for each genotype. Two-tailed Welch's t-test analysis: *p<0.05, ***p<0.001, ****p<0.0001. Bar in panels, 50 μ m. (b) Testes from adult *Brme1*^{-/-} males show a strong reduction of the testis size. See quantification of testis weight/body weight at Figure 2—figure supplement 2b. (c) Spermatogenesis is arrested at epithelial stage IV in Figure 8 continued on next page

Figure 8 continued

Brme1^{-/-} as shown in PAS+hematoxylin stained testis sections. Massive apoptosis of spermatocytes is indicated (red asterisks). The spermatogenic arrest leads to empty epididymides and non-obstructive azoospermia. (St) Seminiferous tubules. (Ep) Epididymides. Bar in panels, 10 μ m. (d–e) Double labeling of (d) spermatocyte and (e) oocyte spreads from WT and *Brme1*^{-/-} mice with SYCP3 (red) and SYCP1 (green). *Brme1*^{-/-} spermatocytes arrest in a zygotene-like stage and show synapsis between non-homologous chromosomes. (e) *Brme1*^{-/-} females showed a subset of fully-synapsed pachynemas (18.5 dpc) but increased numbers of synapsis-defective cells. See extended panel for females at **Figure 8—figure supplement 1e**. Bar in panels, 10 μ m. (f–g) Double labeling with HSF2BP (green) and SYCP3 (red) of (f) spermatocyte and (g) oocyte spreads from *Brme1*^{-/-} mice showing faint HSF2BP labeling in spermatocytes and total absence of labeling in oocytes. Plot on the right of (f) panel represents de quantification of HSF2BP foci in *Brme1*^{-/-} spermatocytes. Nuclei: n = 30 zygonemas/zygonemas-like from two adult mice of each genotype (*Brme1*^{+/+} values from **Figure 3a**) Two-tailed Welch's t-test analysis: ****p<0.0001. Bar in panels, 10 μ m. (h) Western blot analysis of protein extracts from 13 dpp WT, *Brme1*^{-/-} and *Hsf2bp*^{-/-} testes with a specific antibody against HSF2BP. Tubulin was used as loading control. Graph on the right represents the relative quantification of the immunoblotting. Mice: n = 2 *Brme1*^{+/+}, *Brme1*^{-/-} and *Hsf2bp*^{-/-}. Two-tailed Welch's t-test analysis: *p<0.05. The online version of this article includes the following figure supplement(s) for figure 8:

Figure supplement 1. Generation and genetic characterization of *Brme1* knock-out mice.

interaction between HSF2BP and BRME1, we tested whether HSF2BP localization depended on BRME1 by immunolabeling of HSF2BP in *Brme1*^{-/-} spermatocytes and oocytes. Our results showed a strong reduction of HSF2BP staining in BRME1-null spermatocytes (**Figure 8f**) and a total absence in oocytes (**Figure 8g**). Western blot analysis of HSF2BP in 13 dpp testis extracts from *Brme1*^{-/-} mice showed a strong reduction in comparison with the WT control (**Figure 8h**), suggesting again that BRME1 is necessary for HSF2BP protein stabilization.

Immunostaining of *Brme1*^{-/-} spermatocytes for γ H2AX, RPA, the recombinases RAD51 and DMC1 or SPATA22 revealed an accumulation of γ H2AX and RPA on zygonema-like spermatocytes (**Figure 9—figure supplement 1a–b**), a drastic reduction of RAD51/DMC1 foci in early and late zygonema (**Figure 9a and c**, **Figure 9—figure supplement 2a and c**) and a strong accumulation of SPATA22 (**Figure 9e and Figure 9—figure supplement 2e**). According to the meiotic arrest at the zygotene-like stage, MLH1 staining revealed a total absence of COs (**Figure 9g**). As in males, *Brme1*^{-/-} oocytes showed an accumulation of γ H2AX (**Figure 9—figure supplement 1c**) and SPATA22 (**Figure 9f and Figure 9—figure supplement 2f**) and a reduced staining of DMC1 and RAD51 leading to a reduced number of COs (measured as interstitial CDK2; **Figure 9b, d and h**, **Figure 9—figure supplement 2b and d**). However, *Brme1*^{-/-} oocytes did not show RPA accumulation (**Figure 9—figure supplement 1d**). These results are similar to the phenotypes described for HSF2BP mutants (**Figures 4, 5 and 6**, **Figure 4—figure supplement 1**, **Figure 5—figure supplement 1**, **Figure 6—figure supplement 1** and (Brandsma et al., 2019) see **Supplementary file 1f** for a complete comparison among mutants and their meiotic alterations). Thus, both HSF2BP and BRME1 mutant mice show a highly similar phenotype including sexual dimorphism.

BRME1 and HSF2BP form a multimeric complex with PALB2 and BRCA2

To further delineate the interactome of BRME1, we immuno-precipitated BRME1 from testis extracts coupled to mass-spectrometry. We identified as expected HSF2BP as the main interactor, but also BRCA2, PALB2, RAD51 and RPA, strongly suggesting that they form a large multimeric complex (**Supplementary files 1g–1h**). For validation, we transfected the corresponding expression plasmids in HEK293T cells for co-IP analysis. BRME1 co-immunoprecipitated with BRCA2 and HSF2BP when they were all co-transfected, but importantly BRME1 alone did not co-immunoprecipitate with BRCA2 (**Figure 10a–b**). The reciprocal co-IP of BRCA2 with HSF2BP and BRME1 was also positive. We also observed modest but positive co-IP of HSF2BP with RPA, PALB2 and RAD51; and of BRME1 with RAD51 and RPA but not with PALB2 (**Figure 10—figure supplement 1a**). These interactions were further analyzed in a cell-free TNT system coupled to co-immunoprecipitation assays. We observed an absence of direct interaction between any of them, with the exception of BRME1 and HSF2BP, as expected from the Y2H analysis (**Figure 10—figure supplement 1b**). These results suggest that these proteins belong to a complex (or complexes) in vivo (likely through BRCA2) and that the HSF2BP-S167L variant could be altering BRME1 interaction with partners of major BRCA2-containing recombination complexes.

Finally, given the interaction of BRME1 and HSF2BP, we analysed their interdependence in U2OS cells. Transfected HSF2BP was localized diffusely in the cytoplasm and the nucleus (**Figure 10c**).

Figure 9. BRME1 is essential for meiotic recombination. (a–b) Double immunofluorescence of DMC1 (green) and SYCP3 (red) in *Brme1*^{+/+} and *Brme1*^{-/-} (a) spermatocytes and (b) oocytes showing a reduction in the number of DMC1 foci. Plots under each panel represent the quantification. See also extended panels on [Figure 9—figure supplement 2a–b](#). Male nuclei for DMC1: *Brme1*^{+/+}/*Brme1*^{-/-} n = 43/50 early and 41/52 late zygonemas/zygonemas like from two adult mice of each genotype (*Brme1*^{+/+} values from [Figure 5a](#)). Female nuclei for DMC1: *Brme1*^{+/+}/*Brme1*^{-/-} n = 18/30. Figure 9 continued on next page

Figure 9 continued

zygonemas and 21/31 pachynemas from two embryos (17.5 dpc) of each genotype (*Brme1*^{+/+} values from **Figure 5c**). Two-tailed Welch's t-test analysis: ****p* < 0.001, *****p* < 0.0001. Bar in panels, 10 μm. (c–d) Double immunofluorescence of RAD51 (green) and SYCP3 (red) in *Brme1*^{+/+} and *Brme1*^{-/-} (c) spermatocytes and (d) oocytes showing a reduction in the number of RAD51 foci in the *Brme1*^{-/-} in comparison to the WT. Plots under each panel represent the quantification. See also extended panels on **Figure 9—figure supplement 2c–d**. Male nuclei for RAD51: *Brme1*^{+/+}/*Brme1*^{-/-} *n* = 39/39 early and 37/45 late zygonemas/zygonemas like from two adult mice of each genotype (*Brme1*^{+/+} values from **Figure 5b**). Female nuclei for RAD51: *Brme1*^{+/+}/*Brme1*^{-/-} *n* = 35/31 zygonemas and 42/40 pachynemas from two embryos (17.5 dpc) of each genotype (*Brme1*^{+/+} values from **Figure 5d**). Two-tailed Welch's t-test analysis: *****p* < 0.0001. (e–f) Quantification of SPATA22 intensity in (e) spermatocytes and (f) oocyte spreads from *Brme1*^{+/+} and *Brme1*^{-/-}. See immunofluorescences in **Figure 9—figure supplement 2e–f**. Nuclei: Males, *n* = 20 cells from two adult mice of each genotype (*Brme1*^{+/+} from **Figure 6a**). Females, *Brme1*^{+/+}/*Brme1*^{-/-} *n* = 41/40 zygonemas and 40/40 pachynemas from two embryos of each genotype (17.5 dpc) (*Brme1*^{+/+} from **Figure 6a**). Two-tailed Welch's t-test analysis: *****p* < 0.0001. (g) Double immunofluorescence of MLH1 (green) and SYCP3 (red) in *Brme1*^{+/+} and *Brme1*^{-/-} spermatocytes showing the absence of MLH1 labeling in the knock-out. (h) Double labeling of CDK2 (green) and SYCP3 (red) in oocyte spreads from 17.5 dpc *Brme1*^{+/+} and *Brme1*^{-/-} embryos. During meiotic prophase I, CDK2 localizes to the telomeres of chromosomes from leptotene to diplotene. However, around mid-pachytene additional interstitial CDK2 signals appear at CO sites, colocalizing with MLH1. As a measure of COs, just interstitial CDK2 foci (non-telomeric) have been counted. *Brme1*^{-/-} females show a strong reduction in the number of COs. Plot under the panel show the quantification. Nuclei: *Brme1*^{+/+}/*Brme1*^{-/-} *n* = 79/49 from three embryos (17.5 dpc) in WT and two embryos in *Brme1*^{-/-} (*Brme1*^{+/+} from **Figure 6c**). Two-tailed Welch's t-test analysis: ****p* < 0.0001. Bar in all panels, 10 μm.

The online version of this article includes the following figure supplement(s) for figure 9:

Figure supplement 1. DSBs are formed but not properly repaired in *Brme1*-deficient mice and mimic the phenotype of *Hsf2bp*-deficient mice.

Figure supplement 2. Altered dynamic of recombinational proteins in the absence of BRME1.

However, when HSF2BP was co-overexpressed with BRME1, its pattern changed to an intense nucleoplasm staining with nuclear invaginations that resemble nucleoplasmic reticulum (**Figure 10c; Malhas et al., 2011**). Interestingly, such invaginations were reduced when BRME1 was co-transfected with HSF2BP-S167L (**Figure 10c**). In addition, the intensity of the fluorescence signal of HSF2BP-S167L was lower than for HSF2BP-WT and both intensities increased when HSF2BP was co-expressed with BRME1 (**Figure 10c**). Western blot analysis indicated a reduced protein stability of the S167L variant (**Figure 10d**), in agreement with the results observed in vivo (*Hsf2bp*^{S167L/S167L} mutant, **Figure 3**). Interestingly, the protein expression level of transfected HSF2BP increased when co-transfected with BRME1 and was partially dependent on proteasome degradation (**Figure 10d**), indicating a role of BRME1 in stabilizing HSF2BP. Taken altogether and given the low protein expression of BRME1 in the *Hsf2bp*^{S167L/S167L}, these results suggest a functional interdependence between BRME1 and HSF2BP that leads to their lower protein stability/expression in mutant meiotic cells, which might induce recombination defects.

Discussion

Using exome sequencing, we identified the S167L missense variant in *HSF2BP* in a consanguineous family with three cases of POI with secondary amenorrhea. All affected family members are homozygous for the variant, and the healthy relatives are heterozygous carriers. The causality of the HSF2BP-S167L variant is supported by the meiotic phenotype and the subfertility observed in *Hsf2bp*^{S167L/S167L} female mice. Furthermore, the DNA repair defects in murine *Hsf2bp*^{S167L/S167L} meiotic cells, displayed by the reduced number of RAD51/DMC1 foci on DSBs and the subsequent reduction in the number of COs, provide evidence that this missense variant alters meiotic recombination. This conclusion was further supported by the comparative analysis of the S167L allele with the *Hsf2bp* null allele, which revealed that the missense variant can be considered as a hypomorphic allele. This is in agreement with the secondary amenorrhea observed in the patients, and the residual (medically-assisted) fertility in one of the affected sisters. Our identification of *HSF2BP* as a gene implicated in POI is in line with recent reports of POI-causing variants in genes that are required for DNA repair and recombination, such as *MCM8*, *MCM9*, *SYCE1*, *MSH4*, *PSMC3IP*, *FANCM* or *NBN* (AlAsiri et al., 2015; Carlosama et al., 2017; de Vries et al., 2014; Fouquet et al., 2017; He et al., 2018; Tenenbaum-Rakover et al., 2015; Tucker et al., 2018; Wood-Trageser et al., 2014; Zangen et al., 2011).

Meiotic mouse mutants often exhibit sexually dimorphic phenotypes (Cahoon and Libuda, 2019). These differences can have a structural basis, given that the organization of the axial elements is known to be different between sexes. This is supported by the difference in length of the axes and

Figure 10. BRME1 forms a complex with BRCA2 and HSF2BP and stabilizes HSF2BP. (a–b) HEK293T cells were co-transfected with GFP-BRME1, Flag-BRCA2-C and HA-HSF2BP. Protein complexes were immunoprecipitated (IP: green text) with either an anti-Flag, anti-EGFP, anti-HA or IgGs, and analyzed by western blot with the indicated antibody (WB: red text). (a) BRME1 does not co-immunoprecipitate with BRCA2-N, BRCA2-M or BRCA2-C. (b) In the presence of HA-HSF2BP (triple co-transfection) BRCA2-C and BRME1 coimmunoprecipitate (co-IPs between HSF2BP and BRCA2-C are shown in [Figure 6—figure supplement 2a](#)). (c) Transfected U2OS cells with plasmids encoding Flag-HSF2BP (WT or S167L) and EGFP-BRME1 alone or together were immuno-detected with antibodies against Flag (red) and EGFP (green). Transfected HSF2BP (WT and S167L) labels the whole cell (S167L less intense) whereas BRME1 shows nuclear localization. When co-expressed, BRME1 and HSF2BP change their patterns and form nuclear invaginations that resemble nucleoplasmic reticulum. This phenotype is milder in the presence of HSF2BP-S167L than with the WT (graph under the panel: quantification of the number of cells showing a nucleoplasmic reticulum pattern). $n > 400$ cells from two independent transfections of each condition. Two-tailed Welch's t-test analysis: $*p < 0.05$. Bar in panel, 20 μm . (d) HEK293T cells were transfected with Flag-HSF2BP (WT and S167L) alone or with GFP-BRME1. Additionally, cells transfected with Flag-HSF2BP were treated with the proteasome inhibitor (MG132, 10 μM) and analyzed by western blot. Cherry was used as transfection efficiency control. HSF2BP-WT was expressed at higher levels than HSF2BP-S167L and their detection (both the WT and the S167L variant) was increased when co-transfected with BRME1. The increase was greater for the HSF2BP-S167L variant in comparison with the WT. Incubation with MG132 increased the detection levels of transfected HSF2BP mimicking the effect of co-transfecting BRME1. $n = 3$ independent transfections for each condition. Two-tailed Welch's t-test analysis: $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. The online version of this article includes the following figure supplement(s) for figure 10:

Figure supplement 1. Co-immunoprecipitations of BRME1, HSF2BP, RPA, RAD51 and PALB2 expressed from transfected HEK293 cells and from TNT assays.

by the essential role that the meiotic cohesin subunit RAD21L plays in males but not in females (Herrán et al., 2011). In general, meiotic recombination mutants appear to proceed further in female than in male, because in males the asynapsis of the sex bivalent leads to a loss of silencing of the Y chromosome, and perhaps also because of the presence of less stringent checkpoints in oogenesis (Hunt and Hassold, 2002). A sexually dimorphic phenotype is also observed here in our HSF2BP-S167L mutant mice. Both sexes show a significant decrease in the number of COs and consequently an increase in the frequency of meiocytes showing bivalents without CO. This leads to a significant reduction of spermatozoa in the epididymis, while the number of oocytes and their distribution in the follicular pool are not affected in females. Our failure to detect any significant impact of the variant on male fertility could be due to the high variability in litter frequency and size in our study. However, it is known that a strong reduction of the spermatozoa count (up to 60%) does not affect male mouse fertility (Schürmann et al., 2002), which would explain the normal fertility of male mice bearing the HSF2BP-S167L variant despite the presence of overt meiotic alterations. By contrast, female mice with the HSF2BP-S167L variant show a mild sub-fertility phenotype with a reduction of litter frequency. This can be due to the very much fewer gametes available for fertilization in females in comparison to males but also to molecular differences in the meiotic recombination process in both sexes (Cahoon and Libuda, 2019), as displayed by the absence of RPA accumulation or the more pronounced decrease in DMC1 foci observed in HSF2BP-S167L oocytes.

Although we cannot exclude that a POI-like phenotype would appear over time in HSF2BP-S167L female mice, the sub-fertility observed in HSF2BP-S167L females appears to be milder in comparison to the phenotype of the human patients. This could be explained by a lower sensitivity of mice to hypomorphic alleles compared to humans, in a similar manner to the known lower gene dosage sensitivity of the former in the context of genes that are haploinsufficient in human (Veitia, 2003). In addition, the initial events of human female meiosis appear to be more error-prone than in mice, or even than human males, as evidenced by the increased incidence of synaptic defects in the human oocytes or the fact that MLH1 foci appear much earlier in prophase I (Hassold et al., 2007). Furthermore, it has recently been shown that human oocytes exhibit a specific CO maturation inefficiency (Wang et al., 2017). Indeed, despite a higher total number of COs in women, the frequency of bivalents without CO is paradoxically higher in women than in men. Altogether, these observations could explain the stronger phenotype of the human POI patients compared to the *Hsf2bp*^{S167L/S167L} female mice.

The absence of homozygous male carriers in the consanguineous family studied here prevents the direct comparison of the impact of the HSF2BP-S167L variant on fertility phenotypes between human and mouse males. Although future studies might identify infertile men homozygous for the S167L variant, it is interesting to note that another variant of HSF2BP (G224*) was shown to affect recombination rate in males and that two siblings homozygous for this HSF2BP variant in the analyzed Icelandic population were healthy but without descendants, suggesting they were infertile (Halldorsson et al., 2019). This reinforces a conserved function of HSF2BP in human male fertility.

We have shown through a biochemical analysis that BRME1 immunoprecipitates mainly with its partner HSF2BP (constituting or belonging to a complex) but also with PALB2, RAD51, RPA and BRCA2 (in testis extracts). These interactions could possibly be mediated by the multidomain hub protein BRCA2 (Siaud et al., 2011) as HSF2BP interacts directly with BRCA2 (Brandsma et al., 2019) and with BRME1 (this work). In addition, BRCA2 also directly interacts with the DSBs recruiter PALB2, with the recombinases RAD51 and DMC1 (through different specific domains), and with DNA (Siaud et al., 2011). This BRCA2-containing complex participates in the orderly orchestration of events at DSBs such as the initial binding of RPA to the resected DNA, the exchange of RPA by RAD51/DMC1, and the loading of the MEIOB-SPATA22 complex to the RPA complexes (Martinez et al., 2016; Zhao et al., 2015). Interestingly, genes with recently identified variants in POI patients are implicated in the repair of induced DSBs at the early stages of meiosis and encode BRCA2-interacting factors, such as MEIOB, DMC1 or BRCA2 itself (Caburet et al., 2019a; Caburet et al., 2020; Caburet et al., 2019b; He et al., 2018). This highlights the crucial importance and the high sensitivity of this particular meiotic step, and the hub role of BRCA2 as a tightly regulated platform for correct meiotic recombination.

We have also shown by several complementary approaches that the proteins HSF2BP/BRME1 constitute in vivo a functional complex in which both subunits are essential for meiotic recombination and for their mutual protein expression and/or stability in vivo. Accordingly, the genetic depletion of

BRME1 or HSF2BP leads to similar if not identical phenotypes in which oogenesis is altered with severe defects in chromosome synapsis that promotes premature loss of ovarian follicles and spermatogenesis is arrested at zygotene-like stage resulting in a lack of spermatozoa. These meocytes are not able to load the recombinases RAD51/DMC1, impairing the proper repair of DSBs leading to the generation of no COs or very few in males and females, respectively. As a consequence, zygonema-like spermatocytes accumulate the single strand binding proteins SPATA22 and RPA, whereas oocytes accumulates only SPATA22. During the course of the reviewing of this work, three *Brme1* knockouts have been described (Zhang et al., 2020; Shang et al., 2020; Takemoto et al., 2020). All the described male mutants show strong fertility defects and similar molecular alterations although with different severity. In females, the two works that address their analysis (Shang et al. and Takemoto et al.), describe normal fertility which is in contrast with the strong reduction in the follicle pool and meiotic defects observed in our *Brme1*^{-/-} females. The higher severity of our male and female mutants could be explained on the basis of the different genetic background of the mice given that all of them are apparently similar.

The S167L human recessive POI variant behave as a hypomorphic allele in mice, which results in a reduction of the protein expression/stability of itself and of its partner BRME1 in vivo and in transfected cells. As a consequence, both male and female *Hsf2bp*^{S167L/S167} mice show a similar but milder phenotype than that of the *Hsf2bp*^{-/-} or *Brme1*^{-/-}, consisting in a reduction in the spermatozoa count while being fertile and a subtle reduction in female fertility. Molecularly, the reduction observed in the meocytes of the mutant *Hsf2bp*^{S167L/S167L} mice of RAD51/DMC1, the reduction of COs (both in males and females), the accumulation of RPA (only in males) and SPATA22 (in males and females) are also weaker than in the null mutant. The observed accumulation of RPA in males is likely to occur at the early stages of recombination because SPATA22 loading to the DSBs is also increased in the mutants of HSF2BP and BRME1. RPA, as part of a trimeric replication protein complex (RPA1-RPA2-RPA3), binds and stabilizes ssDNA intermediates that form during DNA repair. In meiosis, RPA is also forming a complex with two other essential meiotic players MEIOB (homologue of RPA) and SPATA22. However, the loading of this complex to DSBs is RPA-independent (Shi et al., 2019). It has been postulated that RPA functions in meiosis at two different stages; (i) during the early recombination stages when the DSBs ends are resected by the MRN complex and (ii) during the strand invasion into the homologous duplex that is carried out by RAD51/DMC1 and ssDNA is generated at the displacement loops (Shi et al., 2019). The observed lack of DNA-binding ability of HSF2BP/BRME1 points towards a model in which the absence of the complex HSF2BP/BRME1 through a direct interaction with BRCA2 impairs the replacement of RPA by RAD51/DMC1 in the foci that form on the DSBs of the spermatocytes. Similarly, the reduced expression at the protein level of HSF2BP/BRME1 as a consequence of the POI variant, which does not affect their heterodimerization, would make them less proficient in replacing RPA in the spermatocytes by the recombinases RAD51/DMC1 leading to a lower frequency of COs. Given the unknown function that RPA plays in vivo during oogenesis (Shi et al., 2019), it is tempting to speculate that the role of RPA in mediating the replacement of RAD51/DMC1 in female meiosis would be carried out by another protein complex such as SPATA22/MEIOB in a HSF2BP/BRME1-dependent manner.

Very recently, a high-resolution genome-wide recombination map revealed novel loci involved in the control of meiotic recombination and highlighted genes involved in the formation of the SC (SYCE2, RAD21L, SYCP3, SIX6OS1) and the meiotic machinery itself as determinants of COs (Halldorsson et al., 2019). Within the second category, variants of the SUMO ligase RNF212 and the ubiquitin ligase HEI10 have been largely documented as genetic determinants of the recombination rate in humans and, importantly, so were variants of HSF2BP. Consequently, gene dosage of RNF212 and HEI10 affects CO frequency through their activity in CO designation and maturation (Lake and Hawley, 2013; Reynolds et al., 2013). We found that both BRME1 and HSF2BP localization are unaffected in the loss-of-function mouse mutants of *Rnf212* and *Hei10* (*Ccnd1ip1*). This observation together with the proper co-localization of BRME1/HSF2BP with RPA allows us to map these proteins upstream in the recombination pathway.

It is worth noting that some of the genes affecting the recombination rate have also been described as 'fertility genes', such as SYCP3, HFM1 and HSF2BP (Geisinger and Benavente, 2017; Primary Ovarian Insufficiency Collaboration et al., 2014 and this work). Altogether, we propose that different variants of the same meiotic gene (alleles responsible for mild or strong phenotypes) can give rise to either an altered genome-wide recombination rate with no detrimental effect, or

cause infertility when the decreased recombination rate falls below the lower limit of one COs per bivalent. In the present POI family, the S167L variant in HSF2BP seems to be below that limit. To our knowledge, HSF2BP is one of the very few human genes with variants known to affect both the genome-wide recombination rate in the human population and meiotic chromosome missegregation (fertility) through a reduction of the recombination rate (Halldorsson *et al.*, 2019). Along similar lines, it is conceivable that variants with additive effects (Schimenti and Handel, 2018) can lead to a genome-wide reduction of the recombination rate and thus to aneuploidy and infertility. Specifically, variants in genes involved in meiotic recombination and SC constituents could be responsible for a large fraction of genetic infertilities. These variants should be under purifying selection and would be removed or substantially reduced from the population. However, this is not the case for genes with sexual phenotypic dimorphism (Gershoni and Pietrokovski, 2014) as is apparent for a wide number of meiotic genes (Cahoon and Libuda, 2019), including HSF2BP and BRME1, where individuals of one of the sexes are fertile carriers.

In summary, we describe for the first time a human family where POI co-segregates with a genetic variant in HSF2BP (S167L) in a Mendelian fashion. Humanized mice reveal that the HSF2BP variant is a hypomorphic allele that promotes the lower protein expression and/or stability of the HSF2BP/BRME1 complex and phenocopy in a milder manner the meiotic defects observed in mice lacking either HSF2BP or its direct interactor BRME1.

Materials and methods

Whole exome sequencing

Written informed consent was received from participants prior to inclusion in the study and the institutions involved. Genomic DNA was extracted from blood samples by standard protocols.

For individuals III-3 and III-10, library preparation, exome capture, sequencing and initial data processing were performed by Beckman Coulter Genomics (Danvers, USA). Exon capture was performed using the hsV5UTR kit target enrichment kit. Libraries were sequenced on an Illumina HiSeq instrument as paired-end 100 bp reads. For individual III-2, library preparation, exome capture, sequencing and data processing were performed by IntegraGen SA (Evry, France) according to their in-house procedures. Target capture, enrichment and elution were performed according to manufacturer's instructions and protocols (SureSelect Human All Exon Kits Version CRE, Agilent). The library was sequenced on an Illumina HiSeq 2500 as paired-end 75 bp reads. Image analysis and base calling was performed using Illumina Real Time Analysis (RTA 1.18.64) with default parameters.

Bioinformatic analysis

For the three individuals, sequence reads were mapped onto the human genome build (hg38/GRCh38) using the Burrows-Wheeler Aligner (BWA) tool. Duplicated reads were removed using sam-bamba tools. Whole exome sequencing metrics are provided in *Supplementary file 1a*. Variant calling, allowing the identification of SNV (Single Nucleotide Variations) and small insertions/deletions (up to 20 bp) was performed via the Broad Institute GATK Haplotype Caller GVCf tool (3.7). Ensembl VEP (Variant Effect Predictor, release 87) program was used for initial variant annotation. This tool considers data available in dbSNP (dbSNP147), the 1000 Genomes Project (1000G_phase3), the Exome Variant Server (ESP6500SI-V2-SSA137), the Exome Aggregation Consortium (ExAC r3.0), and IntegraGen in-house databases. Additional annotation data was retrieved using dbNSFP (version 3.5, <https://sites.google.com/site/jpopgen/dbNSFP>) and Varsome (<https://varsome.com/>). Minor allele frequencies were manually verified on GnomAD (<http://gnomad.broad-institute.org/>), ISB Kaviar (<http://db.systemsbiology.net/kaviar/>), and Great Middle Eastern variant database GME Variome (<http://igm.ucsd.edu/gme/>).

Variant filtering was performed on the following criteria:

- minimum depth at variant position of 10,
- correct segregation in the family, on the basis of homozygosity by descent: variants should be homozygous in both affected sisters III-2 and III-3, and heterozygous or homozygous for Reference allele in the fertile sister III-10,
- absence in unrelated in-house fertile controls,

- Minor Allele Frequency (MAF) below 1% in global and in each population in the GnomAD database,
- presence in the coding sequence (i.e not in UTRs, introns, intergenic,)
- high predicted functional impact on the protein. Impact was evaluated based on the predictors included in dbNSFP3.5 (Cahoon and Libuda, 2019; Carlosama et al., 2017; Cox and Mann, 2008) (considered as pathogenic when the majority of the predictors agreed).

The number of variants fulfilling those criteria is provided in **Supplementary file 1b**. Visual inspection of the variant was performed using the IGV viewer.

Sanger sequencing analysis

To confirm the presence and segregation of the variant, direct genomic Sanger DNA sequencing of *HSF2BP* was performed in the patients, the parents and non-affected siblings using specific primers: HSF2BP-ex6F: 5'-ctagaatcttctgtatcctgca-3' and HSF2BP-ex6R2: 5'-ggctctggaagcaaacaggcaa-3'. The resulting chromatograms are shown in **Figure 1—figure supplement 1**.

Predictions of pathogenicity and sequence conservation

The S167L variant was predicted to be pathogenic or deleterious and highly conserved by 11 out of the 18 pathogenicity predictors available in dbNSFP 3.5 (**Supplementary file 1c**). Upon verification, it appears that the conflicting interpretation of this variant might stem from the single occurrence of a Leu at this position in zebrafish. As the change in zebrafish is the variant that we have in the human family, we checked all the available sequences (Ensembl Release 99, January 2020, removing the one-to-many relationships). Ser167 is very highly conserved in mammals, birds and reptiles and fish and is present in 208 of 212 orthologous sequences (**Figure 1—figure supplement 2** and **Figure 1—figure supplement 3**).

Generation of CRISPR/Cas9-edited mice

For developing all the mutant mice models (*Hsf2bp*^{-/-}, *Hsf2bp*^{S167L/S167L}, *Brme1*^{Δ142-472/Δ 142-472}, *Brme1*^{-/-}, *Spo11*^{-/-}, *Rnf212*^{-/-} and *Hei10*^{-/-}) the different crRNAs were predicted at https://eu.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM. The crRNAs, the tracrRNA and the ssODNs were produced by chemical synthesis at IDT (crRNAs and ssODNs sequences are listed in **Supplementary files 1i-1j**). For the *Hsf2bp*^{S167L} we introduced a mutation in the mouse counterpart residue (p.Ser171Leu) of the POI mutation found in the clinical case (p.Ser167Leu). However, for the sake of simplicity, on this manuscript we refer to the mutant allele by the acronym of the human mutation (S167L). The ssODN contains the mutation on the corresponding position of the mouse sequence (c.512C > T, p.Ser171Leu, see character in red in **Supplementary file 1j**) and the PAM mutations avoiding amino acid changes (see characters in bold in the **Supplementary file 1j**). For the *Spo11*^{-/-} mice generation, the ssODN contains the mutations in the active site (TACTAC > TTCTTC p.YY137-138FF, see **Supplementary file 1j**) and the PAM mutations (bold characters in **Supplementary file 1j**). In all cases the crRNA and tracrRNA were annealed to obtain the mature sgRNA. A mixture containing the sgRNAs, recombinant Cas9 protein (IDT) and the ssODN (30 ng/μl Cas9, 20 ng/μl of each annealed sgRNA and 10 ng/μl ssODN) were microinjected into B6/CBA F2 zygotes (hybrids between strains C57BL/6J and CBA/J) (Singh et al., 2015) at the Transgenic Facility of the University of Salamanca. Edited founders were identified by PCR amplification (Taq polymerase, NZYtech) with primers flanking the edited region (see **Supplementary file 1k** for primer sequences). The PCR products were direct sequenced or subcloned into pBlueScript (Stratagene) followed by Sanger sequencing, selecting the founders carrying the desired alleles. The selected founders were crossed with wild-type mice to eliminate possible unwanted off-targets. Heterozygous mice were re-sequenced and crossed to give rise to edited homozygous. Genotyping was performed by analysis of the PCR products produced from genomic DNA extracted from tail biopsies. The primers and the expected amplicon sizes are listed in the **Supplementary file 1k**. Mouse mutants for *Rec8*, *Six6os1* and *Psmα8* have been previously described (Bannister et al., 2004; Gómez-H et al., 2019; Gómez-H et al., 2016).

Ethics statement

All the experiments were approved by the Ethics Committee for Animal Experimentation of the University of Salamanca (USAL) and the Ethics committee of the Spanish Research Council (CSIC) under protocol #00–245. Accordingly, all the mouse protocols used in this work have been approved by the Animal Experimentation committees mentioned above. Specifically, mice were always housed in a temperature-controlled facility (specific pathogen free, spf) using individually ventilated cages, standard diet and a 12 hr light/dark cycle, according to EU law (63/2010/UE) and the Spanish royal law (53/2013) at the “Servicio de Experimentación Animal, SEA. In addition, animal suffering was always minimized, and we made every effort to improve animal welfare during the life of the animals. The mice analysed were between 2 and 4 months of age, except in those experiments where the age is indicated.

Histology

For histological analysis, after the necropsy of the mice their testes or ovaries were removed and fixed in Bouin’s fixative or formol 10%, respectively. They were processed into serial paraffin sections and stained with haematoxylin-eosin (ovaries) or Periodic acid–Schiff (PAS) and hematoxylin (testes). The samples were analysed using a microscope OLYMPUS BX51 and images were taken with a digital camera OLYMPUS DP70. For TUNEL assay, sections were deparaffinized and apoptotic cells were detected with the In Situ Cell Death Detection Kit (Roche) and counterstained with DAPI.

Follicle counting

The inner third of each ovary was serially sliced into 5 µm thick sections and follicles were counted every five sections and classified into four stages (primordial, primary, secondary and antral). Only those follicles in which the nucleus of the oocyte was clearly visible were counted.

Epididymal sperm count

The epididymides were removed, minced and incubated in 1,5 ml of KSOM for 30 min at 37°C to release sperm into the medium. The suspension was incubated for 10 min at 60°C and the total sperm count was quantified by using a hemacytometer.

Fertility assessment

Hsf2bp^{+/+} and *Hsf2bp*^{S167L/S167L} males and females (8 weeks old) were mated with WT females and males, respectively, over the course of 4–12 months. six mice per genotype (seven mice for *Hsf2bp*^{S167L/S167L} females) were crossed. The presence of copulatory plug was examined daily and the number of pups per litter was recorded.

Immunocytology and antibodies

Testes were detunicated and processed for spreading using a conventional ‘dry-down’ technique or for squashing (Gómez-H et al., 2016). Oocytes from fetal ovaries (E16.5, E17.5 and E19.5 embryos) were digested with collagenase, incubated in hypotonic buffer, disaggregated and fixed in paraformaldehyde. Rabbit polyclonal antibodies against HSF2BP and BRME1 were developed by Proteintech™ against a fusion protein of poly-His with full length HSF2BP or BRME1 (pUC57 vector) of mouse origin. Two antibodies (named R1 and R2) were generated against each protein (HSF2BP or BRME1) by immunization of two different host rabbits. Rabbit polyclonal antibody against DMC1 was developed by Proteintech™ against a DMC1 peptide (EESGFQDDEESLFQDIDLLQKHGINMA-DIKKLKSVGICTIKG). The primary antibodies used for immunofluorescence were rabbit αHSF2BP R2 (1:30, Proteintech™), rabbit αBRME1 R2 (1:100, Proteintech™), mouse αSYCP3 IgG sc-74569 (1:100, Santa Cruz), rabbit αSYCP3 serum K921 (provided by Dr. José Luis Barbero, Centro de Investigaciones Biológicas, Spain), rabbit αSYCP1 IgG ab15090 (1:200, Abcam), rabbit anti-γH2AX (ser139) IgG #07–164 (1:500, Millipore), mouse αMLH1 51-1327GR (1:20, BD Biosciences), mouse αCDK2 (1:20; Santa Cruz Sc-6248) rabbit αRAD51 PC130 (1:50, Calbiochem), rabbit αRPA1 serum ‘Molly’ (1:30, provided by Dr. Edyta Marcon, Medical Research University of Toronto, Canada), rat αRPA2 2208S (1:100, Cell Signaling), rabbit αDMC1 (1:500, Proteintech™), rabbit αSPATA22 16989–1-AP (1:60, Proteintech), mouse αFlag IgG (1:100; F1804, Sigma-Aldrich).

Image acquisition and analysis

Slides were visualized at room temperature using a microscope (Axioplan 2; Carl Zeiss, Inc) with 63 × objectives with an aperture of 1.4 (Carl Zeiss, Inc). Images were taken with a digital camera (ORCA-ER; Hamamatsu) and processed with OPENLAB 4.0.3 and Photoshop (Adobe). The slides from the different genotypes used for comparative analyses were all freshly prepared in parallel and immunofluorescence were also carried out in parallel with the same freshly prepared cocktail of antibodies. Slides were not frozen to avoid differences in the background and antigen reactivity. All the images acquired were taken with constant exposure times for comparison. Quantification of foci and fluorescence intensity were performed using Image J software. Only the axis-associated foci were counted. For colocalization analysis, the same nucleus was quantified without rotation (experiment) and after rotating 90 degrees one of the images. This condition allows to determine non-specific colocalization (random). Background was subtracted for intensity quantification. Squashed preparations were visualized with a Delta vision microscopy station. Stimulated emission depletion (STED) microscopy (SP8, Leica) was used to generate the super-resolution images. Secondary antibodies for STED imaging were conjugated to Alexa 555 and 488 (Invitrogen) and the slides were mounted in Prolong Antifade Gold without DAPI.

Generation of plasmids

Full-length cDNAs encoding HSF2BP, BRME1 (full length and delta constructs), RPA1, BRCA2 (N, M and C constructs), PALB2, RAD51, and PSMA8 were RT-PCR amplified from murine testis RNA. The cDNAs were cloned into the EcoRV pcDNA3-2XFlag, SmaI pcDNA3-2XHA or SmaI pEGFP-C1 expression vectors under the CMV promoter. In frame cloning was verified by Sanger sequencing.

Y2H assay and screening

Y2H assay was performed using the Matchmaker Gold Yeast Two-Hybrid System (Clontech) according to the manufacturers' instructions. Mouse *Hsf2bp* cDNA was subcloned into the vector pGBKT7 and was used as bait to screen a mouse testis Mate and Plate cDNA library (Clontech Laboratories Inc). Positive clones were initially identified on double dropout SD (synthetic dropout)/-Leu /- Trp/X- α -Gal/Aureobasidin A plates before further selection on higher stringency quadruple dropout SD /- Ade /- His /- Leu /- Trp/X- α -Gal/Aureobasidin A plates. Prey plasmids were extracted from the candidate yeast clones and transformed into *Escherichia coli*. The plasmids from two independent bacteria colonies were independently grown, extracted and Sanger sequenced.

DNA pull-down assay

ssDNA/dsDNA pull down assays were performed using the protocol previously described by [Souquet et al., 2013](#). A HPLC-purified biotinylated oligonucleotide was used for the DNA pull down assays: ss60-mer F: 5'-GAT CTG CACGACGCACACCGGACGTATCTGCTATCGCTCATG TCAACCGCTCAAGCTGC/3'BiotinTEG/ (IDT) and ss60-mer R (No biotinylated): 5'- GCAGC TTGAGCGGTTGACATGAGCGATAGCAGATACGTCCGGTGTGCGTCGTGCAGATC-3'. Double-stranded DNA annealing was carried out in 50 mM NaCl, 25 mM Tris-HCl, pH 7.5 buffer with complementary sequences at molecular equivalence by a denaturing step (5 min at 95°C) and a slow return to room temperature. DNA was immobilized onto Dynabeads M-280 Streptavidin (Dyna) following the manufacturer instructions (0.2 pmol per 1 μ g of beads). Protein extracts were obtained from in vitro coupled transcription/translation systems (TNT T7 Coupled Reticulocyte Lysate Systems, Promega) according to manufacturer's protocol. 15 μ l of Flag-tagged proteins from TNT assays were pre-incubated on ice for 10 min in modified DBB (DBB: 50 mM Tris HCl, 100 mM NaCl, 10% (w/v) glycerol, Complete Protease inhibitor, 1 mM 2-mercaptoethanol pH 7.4 modified with 25 mM Tris-HCl, 1 mM EDTA plus 5 mg/ml BSA). After this preincubation 500 μ g Dynabeads with immobilized ss- or ds-DNA were added and incubated for 1 hr at 4°C under agitation. Then the beads were washed three times (5 min rotating at RT) in 700 μ l of modified DBB without BSA, before being washed once in 700 μ l of rinsing buffer (modified DBB with 150 mM NaCl). Finally, DNA-binding proteins were eluted by resuspending the beads in 30 μ l of Laemmli buffer boiling the samples for 5 min. The samples were analyzed by western blot.

Cell lines and transfections

HEK293T and U2OS cell lines were obtained from the ATCC and transfected with Jetpei (PolyPlus) according to the manufacturer protocol. Cell lines were tested for mycoplasma contamination using the Mycoplasma PCR ELISA (Sigma).

Immunoprecipitation and western blotting

HEK293T cells were transiently transfected and whole cell extracts were prepared in a 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100 buffer supplemented with protease inhibitors. Those extracts were cleared with protein G Sepharose beads (GE Healthcare) for 1 hr. The corresponding antibodies were incubated with the extracts for 2 hr and immunocomplexes were isolated by adsorption to protein G-Sepharose beads o/n. After washing, the proteins were eluted from the beads with 2xSDS gel-loading buffer 100 mM Tris-HCl (pH 7), 4% SDS, 0.2% bromophenol blue, 200 mM β -mercaptoethanol and 20% glycerol, and loaded onto reducing polyacrylamide SDS gels. The proteins were detected by western blotting with the indicated antibodies. Immunoprecipitations were performed using mouse α Flag IgG (5 μ g; F1804, Sigma-Aldrich), mouse α GFP IgG (4 μ g; CSB-MA000051M0m, Cusabio), ChromPure mouse IgG (5 μ g/1 mg prot; 015-000-003). Primary antibodies used for western blotting were rabbit α Flag IgG (1:2000; F7425 Sigma-Aldrich), goat α GFP IgG (sc-5385, Santa Cruz) (1:3000), rabbit α Myc Tag IgG (1:3000; #06-549, Millipore), rabbit α HSF2BP R2 (1:2000, ProteintechTM), rabbit α BRME1 R1 (1:3000, ProteintechTM), rat α RPA2 (1:1000, Cell Signaling (Cat 22085)). Secondary horseradish peroxidase-conjugated α -mouse (715-035-150, Jackson ImmunoResearch), α -rabbit (711-035-152, Jackson ImmunoResearch), α -goat (705-035-147, Jackson ImmunoResearch) or α -rat (712-035-150, Jackson ImmunoResearch) antibodies were used at 1:5000 dilution. Antibodies were detected by using Immobilon Western Chemiluminescent HRP Substrate from Millipore. Secondary DyLight conjugated α -mouse (DyLight 680, 35518 Thermo-Scientific) and α -rabbit (DyLight 800, 35571 Thermo-Scientific) were used at 1:10,000 dilution. Antibodies were detected using a LI-COR Odyssey fluorescent Imager.

Testis immunoprecipitation

Testis extracts were prepared in 50 mM Tris-HCl (pH8), 500 mM NaCl, 1 mM EDTA 1% Triton X100. 4 mg of protein were incubated with 10 μ g of the specific antibody against the protein to be immunoprecipitated for 2 hr at 4°C rotating. Then 50 μ l of sepharose beads (GE Healthcare) were added to the protein-Ab mixture and incubated overnight at 4°C with rotation. After that, the protein-bound beads were washed four times with 500 μ l of the extraction buffer by centrifugating 1 min at 10,000 rpm and 4°C. Finally, the co-immunoprecipitated proteins were eluted from the beads by resuspending the beads in 50 μ l Laemmli buffer and boiling for 5 min. The samples were analyzed by western blot.

Testis immunoprecipitation coupled to mass spectrometry analysis

200 μ g of antibodies R1 and R2 against BRME1 (two independent IPs) and IgG from rabbit (negative control) were crosslinked to 100 μ l of sepharose beads slurry (GE Healthcare). Testis extracts were prepared in 50 mM Tris-HCl (pH8), 500 mM NaCl, 1 mM EDTA 1% Triton X100. 20 mg of protein extracts were incubated o/n with the sepharose beads. Protein-bound beads were packed into columns and washed in extracting buffer for three times. Proteins were eluted in 100 mM glycine pH3 and analysed by LC-MS/MS shotgun in LTQ Velos Orbitrap at the Proteomics facility of Centro de Investigación del Cáncer (CSIC/University of Salamanca).

Mass spectrometry data analysis

Raw data were analysed using MaxQuant v 1.6.2.6 (Cox and Mann, 2008) against SwissProt Mouse database (UP000000589, Oct, 2019) and MaxQuant contaminants. All FDRs were of 1%. Variable modifications taken into account were oxidation of M and acetylation of the N-term, while fixed modifications included considered only carbamidomethylation of C. The maximum number of modifications allowed per peptide was of 5. Proteins were quantified using iBAQ (Schwanhäusser et al., 2011). Potential contaminants, reverse decoy sequences and proteins identified by site were removed. Proteins with less than two unique peptides in the R1 and R2 groups were not considered for ulterior analysis. Proteins with less than two unique peptides in the control group and more than

two in both groups R1 and R2 were selected as high-confidence candidates (group R1 and R2 only). An additional group of putative candidates was selected for those proteins with two or more unique peptides in one of the R1 or R2 groups and no unique peptides in the control sample (groups R1 only and R2 only, respectively).

Statistics

In order to compare counts between genotypes, we used the Two-tailed Welch's t-test (unequal variances t-test), which was appropriate as the count data were not highly skewed (i.e., were reasonably approximated by a normal distribution) and in most cases showed unequal variance. We applied a two-sided test in all the cases. Asterisks denote statistical significance: *p-value<0.05, **p-value<0.01, ***p-value<0.001 and ****p-value<0.0001.

Acknowledgements

We thank Dr. Barbero and Dr. Edyta Marcon, for providing antibodies against SYCP3 and RPA, respectively, Dr. Emmanuelle Martini for helpful advice with the DNA binding assay, Dr. Fabien Fauchereau for the genetic analysis of the family and Dr. Alex N Zelensky for helpful discussion. This study was supported by Université Paris Diderot and the Fondation pour la Recherche Médicale (Labelisation Equipes DEQ20150331757, SC, A-LT and RAV). This work was supported by MINECO (BFU2017-89408-R) and by Junta de Castilla y León (CSI239P18). NFM, FSS and LGH are supported by European Social Fund/JCyLe grants (EDU/310/2015, EDU/556/2019 and EDU/1083/2013). YBC is funded by a grant from MINECO (BS-2015-073993). The proteomic analysis was performed in the Proteomics Facility of Centro de Investigación del Cáncer, Salamanca, Grant PRB3(IPT17/0019 - ISCIII-SGEFI/ERDF). CIC-IBMCC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia cofunded by the Castilla-León autonomous government and the European Regional Development Fund (CLC-2017-01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Additional information

Funding

Funder	Author
Ministerio de Economía y Competitividad	Alberto M Pendás

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Natalia Felipe-Medina, Resources, Formal analysis, Supervision, Investigation, Methodology, Writing - review and editing; Sandrine Caburet, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review and editing; Fernando Sánchez-Sáez, Yazmine B Condezo, Laura Gómez-H, Paloma Duque, Formal analysis, Methodology; Dirk G de Rooij, Anne Laure Todeschini, Formal analysis, Investigation; Rodrigo Garcia-Valiente, Formal analysis; Manuel Adolfo Sánchez-Martin, Resources, Methodology; Stavitsky A Shalev, Resources; Elena Llano, Conceptualization, Formal analysis, Investigation, Writing - original draft, Writing - review and editing; Reiner A Veitia, Conceptualization, Funding acquisition, Investigation, Writing - original draft, Writing - review and editing; Alberto M Pendás, Conceptualization, Formal analysis, Supervision, Funding acquisition, Investigation, Methodology, Writing - original draft, Writing - review and editing

Author ORCIDs

Natalia Felipe-Medina <https://orcid.org/0000-0001-6975-2524>

Sandrine Caburet <http://orcid.org/0000-0002-7404-8213>

Rodrigo Garcia-Valiente <http://orcid.org/0000-0003-0444-5587>

Reiner A Veitia <https://orcid.org/0000-0002-4100-2681>

Alberto M Pendás <https://orcid.org/0000-0001-9264-3721>

Ethics

Animal experimentation: All the experiments were approved by the Ethics Committee for Animal Experimentation of the University of Salamanca (USAL) and the Ethics committee of the Spanish Research Council (CSIC) under protocol #00-245. Accordingly, all the mouse protocols used in this work have been approved by the above mentioned Animal Experimentation committees. Specifically, mice were always housed in a temperature-controlled facility (specific pathogen free, spf) using individually ventilated cages, standard diet and a 12 h light/dark cycle, according to EU law (63/2010/UE) and the Spanish royal law (53/2013) at the "Servicio de Experimentación Animal, SEA. In addition, animal suffering was always minimized and we made every effort to improve animal welfare during the life of the animals.

Decision letter and Author response

Decision letter <https://doi.org/10.7554/eLife.56996.sa1>

Author response <https://doi.org/10.7554/eLife.56996.sa2>

Additional files

Supplementary files

- Source data 1. Raw data for all figures and figure supplements.
- Supplementary file 1. Supplementary data and tables including quantification data. Supplementary file 1a. shows the whole exome sequencing and mapping metrics for the three genomic samples. Supplementary file 1b. shows the numbers of variants from the WES analysis and passing the various filters. Supplementary file 1c. shows the predictions of pathogenicity and conservations by 18 computational predictors. Supplementary files 1d and 1e. show the quantification of the colocalization between HSF2BP, BRME1, RPA1 and DMC1. Supplementary file 1f. shows the comparative alterations between all the mutants. Associated to **Figures 4, 5, 6 and 9**. Supplementary files 1g and 1h. show the putative BRME1 interactors identified by mass spectrometry. Supplementary files 1i and 1j. show respectively the crRNAs and the ssODN employed in the generation of the different mouse models. Supplementary file 1k. shows the primers and expected product sizes for genotyping mouse models. Supplementary file 1l. shows a summary of all the main and supplementary figures and their relationship.
- Transparent reporting form

Data availability

All data generated or analysed during this study are included in the manuscript and supporting files.

References

- Ahmed EA, de Rooij DG. 2009. Staging of mouse seminiferous tubule cross-sections. *Methods in Molecular Biology* **558**:263–277. DOI: https://doi.org/10.1007/978-1-60761-103-5_16, PMID: 19685330
- AlAsiri S, Basit S, Wood-Trageser MA, Yatsenko SA, Jeffries EP, Surti U, Ketterer DM, Afzal S, Ramzan K, Faiyaz-Ul-Haque M, Jiang H, Trakselis MA, Rajkovic A. 2015. Exome sequencing reveals MCM8 mutation underlies ovarian failure and chromosomal instability. *Journal of Clinical Investigation* **125**:258–262. DOI: <https://doi.org/10.1172/JCI78473>, PMID: 25437880
- Baker SM, Plug AW, Prolla TA, Bronner CE, Harris AC, Yao X, Christie DM, Monell C, Arnheim N, Bradley A, Ashley T, Liskay RM. 1996. Involvement of mouse Mlh1 in DNA mismatch repair and meiotic crossing over. *Nature Genetics* **13**:336–342. DOI: <https://doi.org/10.1038/ng0796-336>, PMID: 8673133
- Bannister LA, Reinholdt LG, Munroe RJ, Schimenti JC. 2004. Positional cloning and characterization of mouse mei8, a disrupted allele of the meiotic cohesin Rec8. *Genesis* **40**:184–194. DOI: <https://doi.org/10.1002/gene.20085>, PMID: 15515002
- Baudat F, Manova K, Yuen JP, Jasin M, Keeney S. 2000. Chromosome Synapsis defects and sexually dimorphic meiotic progression in mice lacking Spo11. *Molecular Cell* **6**:989–998. DOI: [https://doi.org/10.1016/S1097-2765\(00\)00098-8](https://doi.org/10.1016/S1097-2765(00)00098-8), PMID: 11106739

- Baudat F, Imai Y, de Massy B. 2013. Meiotic recombination in mammals: localization and regulation. *Nature Reviews Genetics* **14**:794–806. DOI: <https://doi.org/10.1038/nrg3573>, PMID: 24136506
- Brandtsma I. 2006. *Balancing Pathways in DNA Double Strand Break Repair*. Erasmus Rotterdam University.
- Brandtsma I, Sato K, van Rossum-Fikkert SE, van Vliet N, Sleddens E, Reuter M, Odijk H, van den Tempel N, Dekkers DHW, Bezstarosti K, Demmers JAA, Maas A, Lebbink J, Wyman C, Essers J, van Gent DC, Baarends WM, Knipscheer P, Kanaar R, Zelensky AN. 2019. HSF2BP interacts with a conserved domain of BRCA2 and is required for mouse spermatogenesis. *Cell Reports* **27**:3790–3798. DOI: <https://doi.org/10.1016/j.celrep.2019.05.096>
- Caburet S, Arboleda VA, Llano E, Overbeek PA, Barbero JL, Oka K, Harrison W, Vaiman D, Ben-Neriah Z, García-Tuñón I, Fellous M, Pendás AM, Veitia RA, Vilain E. 2014. Mutant cohesin in premature ovarian failure. *New England Journal of Medicine* **370**:943–949. DOI: <https://doi.org/10.1056/NEJMoa1309635>, PMID: 24597867
- Caburet S, Heddar A, Dardillac E, Creux H, Lambert M, Messiaen S, Misrahi M. 2019a. A homozygous hypomorphic BRCA2 variant causes primary ovarian insufficiency without cancer or Fanconi anemia traits. *bioRxiv*. DOI: <https://doi.org/10.1101/751644>
- Caburet S, Todeschini AL, Petrillo C, Martini E, Farran ND, Legois B, Livera G, Younis JS, Shalev S, Veitia RA. 2019b. A truncating MEIOB mutation responsible for familial primary ovarian insufficiency abolishes its interaction with its partner SPATA22 and their recruitment to DNA double-strand breaks. *EBioMedicine* **42**: 524–531. DOI: <https://doi.org/10.1016/j.ebiom.2019.03.075>, PMID: 31000419
- Caburet S, Heddar A, Dardillac E, Creux H, Lambert M, Messiaen S, Misrahi M. 2020. Homozygous hypomorphic. *Journal of Medical Genetics* **43**:909–912. DOI: <https://doi.org/10.1093/hmg/ddz199>
- Cahoon CK, Hawley RS. 2016. Regulating the construction and demolition of the synaptonemal complex. *Nature Structural & Molecular Biology* **23**:369–377. DOI: <https://doi.org/10.1038/nsmb.3208>, PMID: 27142324
- Cahoon CK, Libuda DE. 2019. Leagues of their own: sexually dimorphic features of meiotic prophase I. *Chromosoma* **131**:199–214. DOI: <https://doi.org/10.1007/s00412-019-00692-x>
- Carlosama C, Elzaia M, Patiño LC, Mateus HE, Veitia RA, Laissue P. 2017. A homozygous donor splice-site mutation in the meiotic gene MSH4 causes primary ovarian insufficiency. *Human Molecular Genetics* **26**:3161–3166. DOI: <https://doi.org/10.1093/hmg/ddx199>, PMID: 28541421
- Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature Biotechnology* **26**:1367–1372. DOI: <https://doi.org/10.1038/nbt.1511>, PMID: 19029910
- de Rooij DG, de Boer P. 2003. Specific arrests of spermatogenesis in genetically modified and mutant mice. *Cytogenetic and Genome Research* **103**:267–276. DOI: <https://doi.org/10.1159/000076812>, PMID: 15051947
- de Vries L, Behar DM, Smirin-Yosef P, Lagovsky I, Tzur S, Basel-Vanagaite L. 2014. Exome sequencing reveals SYCE1 mutation associated with autosomal recessive primary ovarian insufficiency. *The Journal of Clinical Endocrinology & Metabolism* **99**:E2129–E2132. DOI: <https://doi.org/10.1210/jc.2014-1268>, PMID: 25062452
- Fouquet B, Pawlikowska P, Caburet S, Guigon C, Mäkinen M, Tanner L, Hietala M, Urbanska K, Bellutti L, Legois B, Bessieres B, Gougeon A, Benachi A, Livera G, Rosselli F, Veitia RA, Misrahi M. 2017. A homozygous FANCM mutation underlies a familial case of non-syndromic primary ovarian insufficiency. *eLife* **6**:e30490. DOI: <https://doi.org/10.7554/eLife.30490>, PMID: 29231814
- Geisinger A, Benavente R. 2017. Mutations in genes coding for synaptonemal complex proteins and their impact on human fertility. *Cytogenetic and Genome Research* **150**:77–85. DOI: <https://doi.org/10.1159/000453344>
- Gershoni M, Pietrovski S. 2014. Reduced selection and accumulation of deleterious mutations in genes exclusively expressed in men. *Nature Communications* **5**:4438. DOI: <https://doi.org/10.1038/ncomms5438>, PMID: 25014762
- Gómez-H L, Felipe-Medina N, Sánchez-Martín M, Davies OR, Ramos I, García-Tuñón I, de Rooij DG, Dereli I, Tóth A, Barbero JL, Benavente R, Llano E, Pendas AM. 2016. C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nature Communications* **7**:13298. DOI: <https://doi.org/10.1038/ncomms13298>, PMID: 27796301
- Gómez-H L, Felipe-Medina N, Condezo YB, García-Valiente R, Ramos I, Suja JA, Barbero JL, Roig I, Sánchez-Martín M, de Rooij DG, Llano E, Pendas AM. 2019. The PSMA8 subunit of the spermatoproteasome is essential for proper meiotic exit and mouse fertility. *PLOS Genetics* **15**:e1008316. DOI: <https://doi.org/10.1371/journal.pgen.1008316>, PMID: 31437213
- Guo T, Zhao S, Zhao S, Chen M, Li G, Jiao X, Wang Z, Zhao Y, Qin Y, Gao F, Chen ZJ. 2017. Mutations in MSH5 in primary ovarian insufficiency. *Human Molecular Genetics* **26**:1452–1457. DOI: <https://doi.org/10.1093/hmg/ddx044>, PMID: 28175301
- Halldorsson BV, Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP, Gunnarsson B, Oddsson A, Halldorsson GH, Zink F, Gudjonsson SA, Frigge ML, Thorleifsson G, Sigurdsson A, Stacey SN, Sulem P, Masson G, Helgason A, Gudbjartsson DF, Thorsteinsdottir U, et al. 2019. Characterizing mutagenic effects of recombination through a sequence-level genetic map. *Science* **363**:eaau1043. DOI: <https://doi.org/10.1126/science.aau1043>, PMID: 30679340
- Handel MA, Schimenti JC. 2010. Genetics of mammalian meiosis: regulation, dynamics and impact on fertility. *Nature Reviews Genetics* **11**:124–136. DOI: <https://doi.org/10.1038/nrg2723>, PMID: 20051984
- Hassold T, Hall H, Hunt P. 2007. The origin of human aneuploidy: where we have been, where we are going. *Human Molecular Genetics* **16**:R203–R208. DOI: <https://doi.org/10.1093/hmg/ddm243>, PMID: 17911163
- He WB, Tu CF, Liu Q, Meng LL, Yuan SM, Luo AX, He FS, Shen J, Li W, Du J, Zhong CG, Lu GX, Lin G, Fan LQ, Tan YQ. 2018. DMC1 mutation that causes human non-obstructive azoospermia and premature ovarian

- insufficiency identified by whole-exome sequencing. *Journal of Medical Genetics* **55**:198–204. DOI: <https://doi.org/10.1136/jmedgenet-2017-104992>, PMID: 29331980
- Herrán Y, Gutiérrez-Caballero C, Sánchez-Martín M, Hernández T, Viera A, Barbero JL, de Álava E, de Rooij DG, Suja JA, Llano E, Pendás AM. 2011. The cohesin subunit RAD21L functions in meiotic Synapsis and exhibits sexual dimorphism in fertility. *The EMBO Journal* **30**:3091–3105. DOI: <https://doi.org/10.1038/emboj.2011.222>, PMID: 21743440
- Hunt PA, Hassold TJ. 2002. Sex matters in meiosis. *Science* **296**:2181–2183. DOI: <https://doi.org/10.1126/science.1071907>, PMID: 12077403
- Hunter N. 2015. Meiotic recombination: the essence of heredity. *Cold Spring Harbor Perspectives in Biology* **12**:a016618. DOI: <https://doi.org/10.1101/cshperspect.a016618>
- Isaksson R, Tiitinen A. 2004. Present concept of unexplained infertility. *Gynecological Endocrinology* **18**:278–290. DOI: <https://doi.org/10.1080/0951359042000199878>, PMID: 15346664
- Lake CM, Hawley RS. 2013. RNF212 marks the spot. *Nature Genetics* **45**:228–229. DOI: <https://doi.org/10.1038/ng.2559>, PMID: 23438588
- Loidl J. 2016. Conservation and variability of meiosis across the eukaryotes. *Annual Review of Genetics* **50**:293–316. DOI: <https://doi.org/10.1146/annurev-genet-120215-035100>, PMID: 27686280
- Loregian A, Appleton BA, Hogle JM, Coen DM. 2004. Specific residues in the connector loop of the human Cytomegalovirus DNA polymerase accessory protein UL44 are crucial for interaction with the UL54 catalytic subunit. *Journal of Virology* **78**:9084–9092. DOI: <https://doi.org/10.1128/JVI.78.17.9084-9092.2004>, PMID: 15308704
- Malhas A, Goulbourne C, Vaux DJ. 2011. The nucleoplasmic reticulum: form and function. *Trends in Cell Biology* **21**:362–373. DOI: <https://doi.org/10.1016/j.tcb.2011.03.008>, PMID: 21514163
- Martínez JS, von Nicolai C, Kim T, Ehlén Å, Mazin AV, Kowalczykowski SC, Carreira A. 2016. BRCA2 regulates DMC1-mediated recombination through the BRC repeats. *PNAS* **113**:3515–3520. DOI: <https://doi.org/10.1073/pnas.1601691113>, PMID: 26976601
- Matzuk MM, Lamb DJ. 2008. The biology of infertility: research advances and clinical challenges. *Nature Medicine* **14**:1197–1213. DOI: <https://doi.org/10.1038/nm.f.1895>, PMID: 18989307
- Melton DA, Krieg PA, Rebagliati MR, Maniatis T, Zinn K, Green MR. 1984. Efficient *in vitro* synthesis of biologically active RNA and RNA hybridization probes from plasmids containing a bacteriophage SP6 promoter. *Nucleic Acids Research* **12**:7035–7056. DOI: <https://doi.org/10.1093/nar/12.18.7035>, PMID: 6091052
- Primary Ovarian Insufficiency Collaboration, Wang J, Zhang W, Jiang H, Wu BL. 2014. Mutations in HFM1 in recessive primary ovarian insufficiency. *The New England Journal of Medicine* **370**:972–974. DOI: <https://doi.org/10.1056/NEJMc1310150>, PMID: 24597873
- Qiao H, Prasada Rao HB, Yang Y, Fong JH, Cloutier JM, Deacon DC, Nagel KE, Swartz RK, Strong E, Holloway JK, Cohen PE, Schimenti J, Ward J, Hunter N. 2014. Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nature Genetics* **46**:194–199. DOI: <https://doi.org/10.1038/ng.2858>, PMID: 24390283
- Reynolds A, Qiao H, Yang Y, Chen JK, Jackson N, Biswas K, Holloway JK, Baudat F, de Massy B, Wang J, Höög C, Cohen PE, Hunter N. 2013. RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis. *Nature Genetics* **45**:269–278. DOI: <https://doi.org/10.1038/ng.2541>, PMID: 23396135
- Rossetti R, Ferrari I, Bonomi M, Persani L. 2017. Genetics of primary ovarian insufficiency. *Clinical Genetics* **91**:183–198. DOI: <https://doi.org/10.1111/cge.12921>, PMID: 27861765
- Schimenti JC, Handel MA. 2018. Unpackaging the genetics of mammalian fertility: strategies to identify the "reproductive genome". *Biology of Reproduction* **99**:1119–1128. DOI: <https://doi.org/10.1093/biolre/iox133>, PMID: 29878059
- Schürmann A, Koling S, Jacobs S, Saftig P, Krauss S, Wennemuth G, Kluge R, Joost HG. 2002. Reduced sperm count and normal fertility in male mice with targeted disruption of the ADP-ribosylation factor-like 4 (ArL4) gene. *Molecular and Cellular Biology* **22**:2761–2768. DOI: <https://doi.org/10.1128/MCB.22.8.2761-2768.2002>, PMID: 11909968
- Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M. 2011. Global quantification of mammalian gene expression control. *Nature* **473**:337–342. DOI: <https://doi.org/10.1038/nature10098>, PMID: 21593866
- Shang Y, Huang T, Liu H, Liu Y, Liang H, Yu X, Li M, Zhai B, Yang X, Wei Y, Wang G, Chen Z, Wang S, Zhang L. 2020. MEIOK21: a new component of meiotic recombination bridges required for spermatogenesis. *Nucleic Acids Research* **48**:6624–6639. DOI: <https://doi.org/10.1093/nar/gkaa046>, PMID: 32463460
- Sharan SK, Pyle A, Coppola V, Babus J, Swaminathan S, Benedict J, Swing D, Martin BK, Tessarollo L, Evans JP, Flaws JA, Handel MA. 2004. BRCA2 deficiency in mice leads to meiotic impairment and infertility. *Development* **131**:131–142. DOI: <https://doi.org/10.1242/dev.00888>, PMID: 14660434
- Shi B, Xue J, Yin H, Guo R, Luo M, Ye L, Shi Q, Huang X, Liu M, Sha J, Wang PJ. 2019. Dual functions for the ssDNA-binding protein RPA in meiotic recombination. *PLOS Genetics* **15**:e1007952. DOI: <https://doi.org/10.1371/journal.pgen.1007952>, PMID: 30716097
- Siaud N, Barbera MA, Egashira A, Lam I, Christ N, Schlacher K, Xia B, Jasin M. 2011. Plasticity of BRCA2 function in homologous recombination: genetic interactions of the PALB2 and DNA binding domains. *PLOS Genetics* **7**:e1002409. DOI: <https://doi.org/10.1371/journal.pgen.1002409>, PMID: 22194698
- Singh P, Schimenti JC, Bolcun-Filas E. 2015. A mouse geneticist's practical guide to CRISPR applications. *Genetics* **199**:1–15. DOI: <https://doi.org/10.1534/genetics.114.169771>, PMID: 25271304

- Souquet B, Abby E, Hervé R, Finsterbusch F, Tourpin S, Le Bouffant R, Duquenne C, Messiaen S, Martini E, Bernardino-Sgherri J, Toth A, Habert R, Livera G. 2013. MEIOB targets single-strand DNA and is necessary for meiotic recombination. *PLOS Genetics* **9**:e1003784. DOI: <https://doi.org/10.1371/journal.pgen.1003784>, PMID: 24068956
- Takemoto K, Tani N, Takada-Horisawa Y, Fujimura S, Tanno N, Yamane M, Okamura K, Sugimoto M, Araki K, Ishiguro KI. 2020. Meiosis-Specific C19orf57/4930432K21Rik/BRME1 modulates localization of RAD51 and DMC1 to DSBs in mouse meiotic recombination. *Cell Reports* **31**:107686. DOI: <https://doi.org/10.1016/j.celrep.2020.107686>, PMID: 32460033
- Tenenbaum-Rakover Y, Weinberg-Shukron A, Renbaum P, Lobel O, Eideh H, Gulsuner S, Dahary D, Abu-Rayyan A, Kanaan M, Levy-Lahad E, Bercovich D, Zangen D. 2015. Minichromosome maintenance complex component 8 (MCM8) gene mutations result in primary gonadal failure. *Journal of Medical Genetics* **52**:391–399. DOI: <https://doi.org/10.1136/jmedgenet-2014-102921>, PMID: 25873734
- Tucker EJ, Grover SR, Robevska G, van den Bergen J, Hanna C, Sinclair AH. 2018. Identification of variants in pleiotropic genes causing "isolated" premature ovarian insufficiency: implications for medical practice. *European Journal of Human Genetics* **26**:1319–1328. DOI: <https://doi.org/10.1038/s41431-018-0140-4>, PMID: 29706645
- Veitia RA. 2003. A sigmoidal transcriptional response: cooperativity, synergy and dosage effects. *Biological Reviews of the Cambridge Philosophical Society* **78**:149–170. DOI: <https://doi.org/10.1017/S1464793102006036>, PMID: 12620064
- Wang S, Hassold T, Hunt P, White MA, Zickler D, Kleckner N, Zhang L. 2017. Inefficient crossover maturation underlies elevated aneuploidy in human female meiosis. *Cell* **168**:977–989. DOI: <https://doi.org/10.1016/j.cell.2017.02.002>, PMID: 28262352
- Webster A, Schuh M. 2017. Mechanisms of aneuploidy in human eggs. *Trends in Cell Biology* **27**:55–68. DOI: <https://doi.org/10.1016/j.tcb.2016.09.002>, PMID: 27773484
- Wood-Trageser MA, Gurbuz F, Yatsenko SA, Jeffries EP, Kotan LD, Surti U, Ketterer DM, Matic J, Chipkin J, Jiang H, Trakselis MA, Topaloglu AK, Rajkovic A. 2014. MCM9 mutations are associated with ovarian failure, short stature, and chromosomal instability. *The American Journal of Human Genetics* **95**:754–762. DOI: <https://doi.org/10.1016/j.ajhg.2014.11.002>, PMID: 25480036
- Yoshima T, Yura T, Yanagi H. 1998. Novel testis-specific protein that interacts with heat shock factor 2. *Gene* **214**:139–146. DOI: [https://doi.org/10.1016/S0378-1119\(98\)00208-X](https://doi.org/10.1016/S0378-1119(98)00208-X), PMID: 9651507
- Zangen D, Kaufman Y, Zeligson S, Perlberg S, Fridman H, Kanaan M, Abdulhadi-Atwan M, Abu Libdeh A, Gussow A, Kisslov I, Carmel L, Renbaum P, Levy-Lahad E. 2011. XX ovarian dysgenesis is caused by a PSMC3IP/HOP2 mutation that abolishes coactivation of estrogen-driven transcription. *The American Journal of Human Genetics* **89**:572–579. DOI: <https://doi.org/10.1016/j.ajhg.2011.09.006>, PMID: 21963259
- Zhang J, Fujiwara Y, Yamamoto S, Shibuya H. 2019. A meiosis-specific BRCA2 binding protein recruits recombinases to DNA double-strand breaks to ensure homologous recombination. *Nature Communications* **10**:722. DOI: <https://doi.org/10.1038/s41467-019-08676-2>, PMID: 30760716
- Zhang J, Gurusaran M, Fujiwara Y, Zhang K, Echbarhi M, Vorontsov E, Guo R, Pendlebury DF, Alam I, Livera G, Emmanuelle M, Wang PJ, Nandakumar J, Davies OR, Shibuya H. 2020. The BRCA2-MEILB2-BRME1 complex governs meiotic recombination and impairs the mitotic BRCA2-RAD51 function in Cancer cells. *Nature Communications* **11**:2055. DOI: <https://doi.org/10.1038/s41467-020-15954-x>, PMID: 32345962
- Zhao W, Vaithiyalingam S, San Filippo J, Maranon DG, Jimenez-Sainz J, Fontenay GV, Kwon Y, Leung SG, Lu L, Jensen RB, Chazin WJ, Wiese C, Sung P. 2015. Promotion of BRCA2-Dependent homologous recombination by DSS1 via RPA targeting and DNA mimicry. *Molecular Cell* **59**:176–187. DOI: <https://doi.org/10.1016/j.molcel.2015.05.032>, PMID: 26145171
- Zickler D, Kleckner N. 2015. Recombination, pairing, and Synapsis of homologs during meiosis. *Cold Spring Harbor Perspectives in Biology* **7**:a016626. DOI: <https://doi.org/10.1101/cshperspect.a016626>, PMID: 25986558

ARTICLE

Received 23 May 2016 | Accepted 19 Sep 2016 | Published 31 Oct 2016

DOI: 10.1038/ncomms13298

OPEN

C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility

Laura Gómez-H^{1,*}, Natalia Felipe-Medina^{1,*}, Manuel Sánchez-Martín^{2,3}, Owen R. Davies⁴, Isabel Ramos¹, Ignacio García-Tuñón¹, Dirk G. de Rooij⁵, Ihsan Dereli⁶, Attila Tóth⁶, José Luis Barbero⁷, Ricardo Benavente⁸, Elena Llano^{1,9} & Alberto M. Pendas¹

Meiotic recombination generates crossovers between homologous chromosomes that are essential for genome haploidization. The synaptonemal complex is a 'zipper'-like protein assembly that synapses homologue pairs together and provides the structural framework for processing recombination sites into crossovers. Humans show individual differences in the number of crossovers generated across the genome. Recently, an anonymous gene variant in *C14ORF39/SIX6OS1* was identified that influences the recombination rate in humans. Here we show that *C14ORF39/SIX6OS1* encodes a component of the central element of the synaptonemal complex. Yeast two-hybrid analysis reveals that SIX6OS1 interacts with the well-established protein synaptonemal complex central element 1 (SYCE1). Mice lacking SIX6OS1 are defective in chromosome synapsis at meiotic prophase I, which provokes an arrest at the pachytene-like stage and results in infertility. In accordance with its role as a modifier of the human recombination rate, SIX6OS1 is essential for the appropriate processing of intermediate recombination nodules before crossover formation.

¹Instituto de Biología Molecular y Celular del Cáncer (CSIC-Universidad de Salamanca), 37007 Salamanca, Spain. ²Departamento de Medicina, Universidad de Salamanca, 37007 Salamanca, Spain. ³Transgenic Facility, Nucleus platform, Universidad de Salamanca, 37007 Salamanca, Spain. ⁴Institute for Cell and Molecular Biosciences, Newcastle University, Newcastle upon Tyne NE2 4HH, UK. ⁵Reproductive Biology Group, Division of Developmental Biology, Department of Biology, Faculty of Science, Utrecht University, 3584CM Utrecht, The Netherlands. ⁶Institute of Physiological Chemistry, Medical Faculty of TU Dresden, Fiedlerstrasse 42, 01307 Dresden, Germany. ⁷Departamento de Biología Celular y Molecular, Centro de Investigaciones Biológicas (CSIC), Madrid 28040, Spain. ⁸Department of Cell and Developmental Biology, Biocenter, University of Würzburg, D-97074 Würzburg, Germany. ⁹Departamento de Fisiología y Farmacología, Universidad de Salamanca, 37007 Salamanca, Spain. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to A.M.P. (email: amp@usal.es) or to E.L. (email: ellano@usal.es).

During meiosis, two successive rounds of chromosome segregation occur following a single round of replication, resulting in the formation of haploid gametes from diploid progenitors¹. This ploidy reduction is achieved through a series of meiosis-specific events, including pairing, synapsis, crossover formation between homologues, suppression of sister centromere separation during the first (reductional) division and separation of sister chromatids during the second (equational) division. Homologous chromosomes become tethered together through numerous recombination events between homologous non-sister chromatids, which are triggered by double-strand break induction. Through resolution, a subset of recombination events mature into crossovers (chiasmata) that maintain the physical tethering between homologues until the onset of anaphase I (ref. 1).

In humans, the number of crossovers occurring across the genome differs between individuals. Through exploitation of data resources in Iceland, Kong *et al.*² recently analysed over two million recombination events and putative variants from 2,261 whole genome-sequenced individuals to identify variants that influence the global recombination rate. Among the new variants, several coding SNPs in very well-known meiotic players were identified, including the histone methyltransferase PRDM9 and the meiotic cohesin RAD21L, the latter of which has been the focus of our previous meiotic studies^{3–5}.

In addition to their well-established role in mediating sister chromatid cohesion through ring structure formation, cohesin complexes are also responsible for the assembly of the synaptonemal complex (SC)^{5–7}. The SC is a proteinaceous structure that holds homologous chromosome pairs in synapsis during prophase I, from zygonema to pachynema. It consists of two parallel axial elements (AEs) that bind sister chromatids together, and which become known as lateral elements (LEs) upon chromosome pairing. It also contains transverse filaments, which connect (synapse) the two LEs together. Transverse filament proteins are recruited to LEs and undergo zipper-like assembly, bridging between LEs through the formation of the midline central element (CE), and thereby generating the tripartite structure of the SC⁸. To date, only seven protein structural components of the SC have been identified in mammals, namely LE proteins SYCP2 and SYCP3, transverse filament protein SYCP1, and CE proteins SYCE1, SYCE2, SYCE3 and TEX12 (ref. 9). The location of CE-specific proteins is, by definition, restricted to the synapsed regions of the chromosomes from zygotene to diplotene⁹. The SC provides the structural framework for synapsis, double-strand break (DSB) repair and exchange between homologues^{10,11}. It is known from mouse mutants and through human genetic analysis of families with non-obstructive azoospermia and premature ovarian failure, that alterations in these genes (that is, meiosis-specific cohesin subunit STAG3, and SYCE1) can result in meiotic arrest and human infertility^{12,13}.

To gain further insight into the biological processes affecting recombination rates across the human genome, we have investigated the list of genes that were recently identified as having coding variants². We focus our analysis on the anonymous *C14ORF39/SIX6OS1* gene (herein *SIX6OS1*) based on its restricted pattern of transcription and expression. Here, we show that *C14ORF39/SIX6OS1* encodes a component of the CE of the SC. Yeast two-hybrid analysis reveals that *SIX6OS1* interacts with SYCE1. In addition, mice lacking *SIX6OS1* are defective in chromosome synapsis at meiotic prophase I, which provokes an arrest at the pachytene-like stage and results in mouse infertility. In accordance with its role as a modifier of the human recombination rate, *SIX6OS1* is essential for the appropriate processing of intermediate recombination nodules before crossover formation in mice.

Results

C14ORF39/SIX6OS1 is a protein of the mammalian SC. The sequence variants identified by Kong *et al.*² include known genes functioning in meiotic recombination such as *RNF212* (refs 14,15), *RAD21L* (ref. 4), *PRDM9* (ref. 16), *MSH4* (ref. 17) and *CCNB1P1* (ref. 18). They also include an anonymous open reading frame, containing a nonsynonymous SNV with unknown function (rs1254319, p.Leu524Phe). This gene, named *SIX6OS1*, is also annotated as coding for a natural antisense transcript (NAT) that is associated with the eye transcription factor *SIX6* (ref. 19). However, and in contrast to most natural antisense transcripts, *SIX6OS1* shows a high degree of sequence similarity between mouse (*4930447C04Rik*) and human (*C14ORF39*), and contains large theoretical conserved open reading frames encoding putative proteins of 587 and 574 residues in mouse and human, respectively (Supplementary Fig. 1). Phylogenetic analysis indicates that *SIX6OS1* is a unique gene that appeared firstly in the genomes of cartilaginous fish, and it can be clearly identified from lobed fin fish to mammals (Supplementary Fig. 1). Interestingly, in the variant rs1254319 (p.Leu524Phe), the phenylalanine residue is very well conserved in all genomes (including several other primates, for example, Orangutan and Baboon) except humans (Supplementary Fig. 1).

Analysis of *Six6os1* mRNA expression in mouse tissues by RT-qPCR (Fig. 1a) revealed that it is most abundantly expressed in testis (in agreement with GTEx database²⁰).

The *Six6os1* open reading frame predicts a protein of around 70 kDa, in agreement with our western blot analysis (Supplementary Fig. 3a). Sequence analysis reveals the presence of an evolutionarily conserved region of high helical content within its N terminus (corresponding to amino acids 1–261), including a short stretch of predicted coiled-coil structure towards the C-terminal end of this region (Supplementary Fig. 2a). These features are typical of SC proteins, which commonly contain a high percentage of helical content and adopt homo- or hetero-oligomeric helical bundle or coiled-coil structures^{21,22}. The presence of conserved proline residues between predicted helices suggests that the structure includes helix–turn–helix motifs, rather than adopting an extended helical conformation such as that observed in the crystal structure of SYCP3 (ref. 22). This feature is in common with SC central element proteins SYCE1 and SYCE3, but contrasts with the elongated helical structure predicted and observed in solution for central element complex SYCE2–TEX12 and transverse filament protein SYCP1 (ref. 21). We therefore predict that this N-terminal helical region could mediate interactions with structural proteins of the SC. The remainder of the *SIX6OS1* sequence is predicted to be largely unstructured, but importantly contains patches of evolutionary conservation towards its C-terminal end (Supplementary Fig. 2b). These features are characteristic of flexible sequences that interact with globular proteins at specific peptide motifs through induced fit, and thereby mediate the assembly of macromolecular complexes. The unstructured C-terminal region further contains numerous predicted phosphorylation sites, including four conserved S/TP potential CDK phosphorylation sites (Supplementary Fig. 2c), which may function in regulating the timely assembly of such macromolecular complexes during the first meiotic division.

To explore the localization of *SIX6OS1*, we *in vivo* electroporated an expression plasmid encoding *SIX6OS1*–GFP²³ into mouse testis. After 48 h, *SIX6OS1*–GFP co-localized with SYCP3 along the synapsed LEs at pachynema (spermatocytes in which homologues are fully synapsed) (Fig. 1b). In addition, we carried out a detailed analysis of mouse spermatocytes and oocytes spreads through double labelling with specific antibodies against *SIX6OS1* (which were intensively validated, Fig. 5c;

Figure 1 | Transcriptional analysis and distribution of *SIX6OS1* in mouse meiocytes. (a) Relative transcription of *Six6os1* and *Rad21l* (ref. 4) mRNA by quantitative reverse transcription PCR (RT-qPCR) in mouse tissues. β -Actin transcription was used to normalize the expression (mean $\pm$ s.d., three replicates). (b) Immunolabelling of *in vivo* electroporated *SIX6OS1*-GFP in mouse testis. *SIX6OS1* was detected with anti-GFP (green) and endogenous *SYCP3* was detected using mouse anti-*SYCP3* (red). DNA was stained with DAPI (blue). During pachytene, *SIX6OS1* colocalizes with *SYCP3* along synapsed lateral elements (LEs) including the pseudoautosomic region (PAR) of the XY bivalent (spermatocytes). In diplotene and late diplotene, *SIX6OS1* localizes at the still synapsed LEs. (c) Double immunolabelling of endogenous *SIX6OS1* (green) and *SYCP3* (red) in spermatocytes. DNA was stained with DAPI (blue). During pachynema, *SIX6OS1* is located at the synapsed autosomal LEs and at the PAR of the sex XY bivalent. (d, e) Co-labelling of spermatocytes spread preparations with *SIX6OS1* (green) and *SYCP1*, *SYCE1*, *SYCE3*, *SYCE2* or *TEX12* (red), showing that *SIX6OS1* localizes to the synapsed LEs but best mirrors *SYCE1* localization. (f) Immunoelectron microscopy of frozen mouse testis sections marked with goat anti-*SIX6OS1* antibody. Left panel corresponds to an autosomal chromosome and right panel to the XY bivalent in which the PAR is shown. Gold particles 6 nm. Scale bar in b–e, 10 μ m. PAR is indicated with an asterisk in b and c.

Supplementary Fig. 3) and SYCP3 or SYCP1 (Fig. 1c,d; Supplementary Fig. 4). SIX6OS1 was detected from zygonema to pachynema, co-localizing with SYCP1 along synapsed LEs, but with diminished co-localization at telomeres (Fig. 1d). On the XY bivalent, the pseudoautosomal synapsed region labelled positively for SIX6OS1 (Fig. 1c; Supplementary Fig. 4b). As desynapsis

progressed through diplonema, SIX6OS1 (together with SYCP1) was not observed at the desynapsed regions of spermatocytes and oocytes (Fig. 1d; Supplementary Fig. 4a). Thus, SIX6OS1 partially overlaps the distribution of SYCP1 at the synapsed axes.

We next measured and compared the fluorescence profile of SIX6OS1 along the chromosome axes with those of CE

Figure 2 | Co-localization profile of SIX6OS1 with central element proteins. (a) Double immunostaining of SIX6OS1 (green) and SYCP1, SYCE1, SYCE3, SYCE2 or TEX12 (red). Immunofluorescence signal levels were measured on synapsed chromosome axes. Left plots represent normalized signal intensity profiles of SIX6OS1 with each CE protein. Right plots show regression analysis of the correlation between each pair. **(b)** Plot of the mean correlation between SIX6OS1 and SYCP1, SYCE1, SYCE3, SYCE2 or TEX12 ($n=38$ axial elements (AEs), mean $\pm$ s.d.). The best correlation value was obtained with SYCE1. Scale bar, 2.5 μ m.

proteins SYCP1, SYCE1, SYCE2, SYCE3 and TEX12 (Figs 1e, 2a). This revealed that SIX6OS1 localization is more similar to the continuous pattern of SYCP1 and SYCE1/3 (best regression with SYCE1) than to the more punctate pattern of SYCE2 and TEX12 (ref. 24) (Fig. 2b). However, the SIX6OS1 localization pattern is not strictly identical to SYCP1, especially at the telomeres, where SIX6OS1 stained more weakly (Figs 1d and 2a).

In addition, we performed immuno-gold electron microscopy on testis sections using the same SIX6OS1 antibody. The gold particle distribution agrees with those previously reported for CE proteins SYCE1, SYCE2, TEX12 and SYCE3 (refs 24,25) thus supporting the localization of SIX6OS1 at the CE (Fig. 1f). Taken together, these results demonstrate that SIX6OS1 is a meiotic protein that is located at the CE of the SC.

SIX6OS1 interacts with SYCE1. To understand the role of SIX6OS1 in meiosis, we searched for proteins that interact with mouse SIX6OS1 through yeast two hybrid (Y2H) screening (see Methods). Of the 6.1 million independent clones screened, 90 colonies containing interacting bait and prey fusion proteins grew under the highest stringency conditions. Analysis of the positively interacting clones (Methods) revealed that they all encode SYCE1, a well-known protein of the CE of the SC²⁶. To confirm and validate this interaction, we made use of heterologous HEK 293T cells by transiently transfecting expression plasmids encoding GFP-SIX6OS1 and Myc-SYCE1. SIX6OS1 was found to co-immunoprecipitate (co-IP) reciprocally with SYCE1 (Fig. 3a).

Through biochemical, biophysical and crystallographic studies, all SC proteins studied to date have proven to exist as homo- and/or hetero-oligomers. To explore the possible self-association of SIX6OS1, we co-transfected *Six6os1* tagged with two different epitopes (GFP and Flag) and found that they co-immunoprecipitate (Fig. 3b), suggesting that it exists as a homo-oligomer.

Next, we adopted a candidate gene approach to identify additional putative interactors of SIX6OS1. We co-transfected *Six6os1* with cDNAs encoding each of the known central element proteins (SYCE1, SYCE2, SYCE3 and TEX12), transverse filament protein SYCP1, LE protein SYCP3, and the meiotic cohesins REC8 and Sororin (a recently identified cohesin subunit localized to synapsed regions²⁷) (Fig. 3a; Supplementary Fig. 5a,b). As positive controls we used the well-known interaction between SYCE2 and TEX12 (ref. 21), and between SYCE3 and SYCE1 (ref. 28; Supplementary Fig. 5c). We detected co-immunoprecipitation only between SIX6OS1 and SYCE1.

Finally, we used truncated forms of SIX6OS1 to show that the N-terminal half (1–286), but not the C-terminal half (287–574), is able to interact with SYCE1 in isolation (Fig. 3c). Together, these results indicate that the interaction between SIX6OS1 and SYCE1 occurs in a very specific manner through the N-terminal half of SIX6OS1.

Polycomplex formation of SIX6OS1. SYCP1 and SYCP3 form filamentous structures, so-called polycomplexes, in the cytoplasm of transfected cells. Thus, co-expression of an interacting partner with SYCP1 or SYCP3 may lead to its recruitment to polycomplexes^{25,29}. To analyse this, we transfected the cDNA encoding SIX6OS1 in combination with SYCP1 alone or in different combinations with SYCE1, SYCE2, SYCE3 and TEX12, and studied their distribution by immunofluorescence. In absence of SYCP1, single transfections of SYCE1, SYCE3, SYCE2, TEX12 and SIX6OS1 produced different distributions (cytoplasmic aggregates, whole cell, cytoplasmic and nuclear, respectively), but in all cases without the appearance of self-assembled higher order structures (Fig. 3d; Supplementary Fig. 6a).

When transfected in combination with SYCP1, SIX6OS1 was not recruited to the filamentous structures, and its cellular localization was not modified (Fig. 3d). We then tested whether the distribution pattern of transfected *Six6os1* in COS7 cells was altered by its co-transfection with *Syce1*, *Syce3*, *Syce2* or *Tex12*. This revealed that the SIX6OS1 distribution is drastically affected only in the presence of SYCE1 (from diffuse pattern to punctate, Fig. 3d; Supplementary Fig. 6b). Moreover, when COS7 cells were transfected with cDNAs encoding SYCP1, SYCE1, SYCE3 and SIX6OS1 simultaneously, all components co-localized in speckled cytoplasmic aggregates (Fig. 3e). This pattern was only altered when SYCE1 was absent (Supplementary Fig. 6c–f). We further validated the interaction between SYCE1 and SIX6OS1 in transfected COS7 cells by proximity ligation assay (PLA) (Supplementary Fig. 7). In summary, these results further support the findings of the Y2H and co-IP experiments by showing that SIX6OS1 interacts specifically and exclusively with SYCE1 in transfected COS7 cells.

SIX6OS1 loading is dependent on synapsis. To investigate the possible dependence of SIX6OS1 localization on the presence of other SC proteins, we analysed spermatocytes of mice deficient for *Syce3* (ref. 30), *Sycp1* (ref. 31) and meiotic cohesins *Rad21l* (ref. 4), *Rec8* (ref. 32) and *Stag3* (ref. 33). These meiotic mutants display different synaptic defects, from mild to more severe. In the absence of RAD21L or REC8, double labelling of SIX6OS1 and SYCP3 shows that SIX6OS1 is localized to synapsed-like regions (Fig. 4a). Interestingly, in *Rec8* mutants, where there is no synapsis between homologues but instead the AEs of 40 univalents are decorated with SYCP1 as a result of ‘synapsis-like’ events between sister chromatids³⁴, SIX6OS1 is also present at these atypical synapsed-like regions (Fig. 4b). In *Stag3*-deficient mice, in which spermatocytes show almost no synapsis and very short AEs, SIX6OS1 also mimics SYCP1 localization. Finally, in mice lacking the central element proteins SYCE3 and SYCP1, in which AEs completely fail to synapse in a pachytene-like stage^{30,31}, SIX6OS1 was not detected despite the presence of a weak discontinuous pattern of SYCP1 deposition in the *Syce3* mutant (Fig. 4; Supplementary Fig. 9c)³⁰. These results, obtained through immunofluorescence analysis, allow a precise comparison of the different CE-mutant phenotypes (compare Fig. 4 with Fig. 7a by Schramm *et al.*³⁰), and thus provide a global picture of the biology of the CE proteins. In this regard, we predict that *Syce1* mutants will also be defective in SIX6OS1 loading since SYCE3 deficiency leads to failure in loading of both SYCE1 (ref. 30) and SIX6OS1 onto the LEs (Fig. 4).

Together, our results indicate that SIX6OS1 is a new protein of the CE, and its loading is consequently dependent on the assembly of the tripartite SC structure that occurs upon synapsis between homologous chromosomes or, interestingly, even between sister chromatids.

Mice lacking SIX6OS1 are infertile. To investigate the function of SIX6OS1 we generated a mutation of the murine *Six6os1* gene by CRISPR/Cas9 genome editing (Fig. 5a). The *a priori* most suitable null mutation was chosen by PCR sequencing of the targeted region of the murine *Six6os1* gene (Fig. 5b). A founder line was crossed with wild-type C57BL/6J and the resulting heterozygotes were interbred. Spermatocytes from homozygous targeted mice showed no SIX6OS1 protein expression by immunofluorescence when analysed using two independent polyclonal antibodies (Fig. 5c; Supplementary Fig. 3c). These results indicate that the mutation is a null allele of the *Six6os1* gene (herein *Six6os1*^{−/−}).

Figure 3 | SIX6OS1 interacts specifically with SYCE1. (a–c) HEK 293T cells were transfected or co-transfected with the indicated expression vectors. Protein complexes were immunoprecipitated overnight with either an anti-Flag, anti-EGFP or anti-Myc antibody, and were analysed by immunoblotting with the indicated antibody. (a) SIX6OS1 co-immunoprecipitates (co-IP) with SYCE1 (as well as in the reciprocal IP) but not with either SYCP1, SYCE3, SYCE2 or TEX12. (b) SIX6OS1-Flag co-immunoprecipitates with SIX6OS1-GFP, suggesting that it is able to form at least dimers. (c) SYCE1 co-immunoprecipitates with the SIX6OS1 N-terminal half (1–286) but not with the C-terminal half (287–574). Immunoprecipitation of SYCE1 and full length SIX6OS1 was used as positive control. (d) COS7 cells were transfected with *Six6os1* alone (left panel) or in combination with *Sycp1* and *Syce1* (right panel). SIX6OS1 localization drastically changed in the presence of SYCE1 but not with SYCP1. (e) *Sycp1*, *Syce3*, *Syce1* and *Six6os1* were simultaneously co-transfected in COS7 cells and found to co-localize in the cytoplasm in the punctate pattern of SYCE1. The experiments were reproduced three times. Scale bars, 15 μ m.

Mice lacking SIX6OS1 did not display any obvious abnormalities but were sterile. Consistent with this, testes size from *Six6os1*^{−/−} mice was only 30% of wild-type mice, and their epididymides exhibited complete absence of spermatozoa (Fig. 6a,b). Histological analysis of adult *Six6os1*^{−/−} testes revealed seminiferous tubules that lacked postmeiotic cell types. The presence of spermatogonia, spermatocytes and Sertoli and Leydig cells was not altered (Fig. 6b). By identifying groups of associated germ cell types in seminiferous tubule sections, the twelve stages of the epithelial cycle can be distinguished. Following these criteria, mutant adult mice showed an arrest at stage IV of the epithelial cycle. Spermatogenesis proceeds

apparently normally in these mice up to prophase I, and then at stage IV, there is a massive apoptosis of spermatocytes (Fig. 6b). At 18 days of age, extensive apoptosis was also detected (Fig. 6c), suggesting that SIX6OS1 deficiency already affects spermatocytes during the first wave of meiosis. Thus, we conclude that SIX6OS1 is essential for spermatogenesis and its deficiency leads to non-obstructive azoospermia and consequently to infertility.

Histological analysis of whole ovaries of *Six6os1*^{−/−} female mice at 4 months of age showed a lack of oocytes and a dense stroma (Fig. 6d). To investigate when this ovarian failure occurred, we histologically analysed ovaries from 6 day old

Figure 4 | SIX6OS1 loading is dependent on synapsis but not on AE proteins. (a) Double labelling of SIX6OS1 (green) and SYCP3 (red) or (b) SYCP1 (red) in *Rad21*^{-/-}, *Rec8*^{-/-}, *Stag3*^{-/-}, *Syce3*^{-/-} and *Sycp1*^{-/-} showing that loading of SIX6OS1 is dependent on synapsis. SIX6OS1 is detected in the synapsed LEs of meiotic cohesin mutants but is absent from unsynapsed AEs in *Syce3*^{-/-} and *Sycp1*^{-/-} spermatocytes. Scale bar, 10 μ m.

females (6 d.p.p.), a time point at which all oocytes are arrested in dictyate and present a large number of primordial follicles (outer cortex) and growing oocytes (inner cortex) (Fig. 6d). At 6 d.p.p., ovaries of *Six6os1*^{-/-} mice are already depleted of follicles and show a severe ovarian dysgenesis (Fig. 6d) that is responsible for the absence of oogenesis and consequently for the severe premature ovarian failure.

SIX6OS1 is essential for chromosome synapsis. To characterize the meiotic defect in detail, *Six6os1*-deficient meocytes were analysed using spread preparations from males as well as from fetal females. They were initially stained for AEs proteins (that is, SYCP3), revealing that mutant spermatocytes have AEs of normal morphology and composition (Fig. 7). Further, cohesins SMC3, REC8, STAG3, RAD21L and SMC1B are all present in AEs together with SYCP3 in *Six6os1*-deficient mice (Supplementary Fig. 8). As expected, in wild-type spermatocytes homologues were aligned in close juxtaposition during zygotene, and full synapsis was achieved at pachynema (Fig. 7a; Supplementary Fig. 9a). However, in both male and female *Six6os1*-deficient mice, synapsis failed to develop between homologues and all meocytes were arrested in a pachytene-like stage, in most cases with their AEs properly aligned (A-type). However, a subset of meocytes, more frequently observed in oocytes than in spermatocytes, showed poorly or even completely unaligned chromosome pairs (U-type; 17.6 $\pm$ 3.7% in spermatocytes; n = 3 and 79.06 $\pm$ 18.9% in oocytes; n = 3,

Fig. 7a,b; Supplementary Fig. 9a). The lack of synapsis, and the absence of breaks in unsynapsed AEs, were further analysed by counting the number of centromeres (ACA staining, 21 versus 40 in spermatocytes and 20 versus 40 in oocytes, Supplementary Fig. 10a) and telomeres (RAP1 marker, 40 versus 80, Supplementary Fig. 10b) in arrested meocytes. This confirmed complete desynapsis but with the full integrity of AEs (all of the AEs have two RAP1 signals at their ends). Finally, and to refine the stage of the blockade, we immunolabelled *Six6os1*^{-/-} spermatocytes with the mid pachytene-specific histone variant H1t (ref. 35). The positive staining for H1t (Supplementary Fig. 10c) indicates that arrested spermatocytes reach the mid-pachytene stage.

To gain further insight into the synaptic defects, we double immunolocalized SYCP3 and SYCP1. In contrast to other CE mutants such as *Syce3*, and even more so for *Syce2* (ref. 36) and *Tex12* (ref. 37), *Six6os1*-deficient spermatocytes have reduced levels of SYCP1 labelling (93.70% reduction in *Six6os1*^{-/-} versus 54.36% reduction in *Syce3*^{-/-}). Mutant oocytes, however, show a slightly weaker reduction of SYCP1 staining (79.28% reduction, Fig. 7a,b; Supplementary Fig. 9c for quantification). We next double immunolocalized SYCP3 with SYCE1, SYCE3, SYCE2 and TEX12, revealing the absence of staining of all CE components in *Six6os1*^{-/-} spermatocytes (Fig. 7c) and oocytes (Supplementary Fig. 11). Similarly, the regulatory cohesin subunit Sororin, which is located at the CE²⁷, is also lacking in *Six6os1*-deficient spermatocytes (Supplementary Fig. 8).

Figure 5 | Generation and genetic characterization of *Six6os1*-deficient mice. (a) Schematic representation of the wild-type locus (WT) and the genome editing strategy at the *Six6os1* locus, showing the sgRNAs, the corresponding coding exons (light grey) and non-coding exons (open boxes). Thin (non-coding) and thick (coding sequences) lines under exons represent the expected transcript derived from WT and *Six6os1* edited allele. ATG, initiation codon; TGA, stop codon. The nucleotide sequence of the 124 base pair deletion derived from PCR amplification of DNA from the *Six6os1*^{edited/edited} is indicated. Primers are represented by arrows. (b) PCR analysis of genomic DNA from three littermate progeny of *Six6os1*^{+/-} heterozygote crosses. The PCR amplification with primers F and R (indicated by arrows) revealed 413 and 289 bp fragments for wild-type and disrupted alleles, respectively. (+/+), (+/-) and (-/-) designate wild-type, heterozygous and homozygous knockout animals, respectively. (c) Double immunofluorescence of spermatocytes at pachytene stage obtained from *Six6os1*^{+/+} and *Six6os1*^{-/-} mice using SYCP3 (red) and a goat polyclonal antibody against SIX6OS1 (green). Scale bar, 10 μ m.

To establish a direct causal role of SIX6OS1 deficiency in the observed phenotype, we analysed mice during the first almost synchronous wave of spermatogenesis at 18 d.p.p. We find that the meiotic arrest observed at this stage mimics that observed in adult males. Interestingly, the arrest is more homogeneous, with a lack of SYCP1 labelling in all AEs, despite the presence of both A- and U-type AEs (U-type $35.37 \pm 2.3\%$; Supplementary Fig. 7b). This suggests that the weak SYCP1 staining observed in the adult mutants could be a byproduct of a longer arrest.

In *Sycp1*-deficient mice, SIX6OS1 (Fig. 4), SYCE1-3 and TEX12 are not recruited to the SC³⁰. Surprisingly, the lack of any of these central element proteins also leads to the aberrant deposition of SYCP1 in a weak discontinuous pattern, with the severest phenotype occurring in SIX6OS1 deficiency (weakest staining). This mutual interdependence, in addition to the fact that biochemical reactions (that is, DSB processing) take place in the three-dimensional (3D) mesh of the SC, make it difficult to distinguish cause and effect when analyzing mutant mice such as

Figure 6 | The absence of SIX6OS1 provokes azoospermia and ovarian failure. (a) Genetic ablation of *Six6os1* leads to a reduction of the testis size ($n=8$ wild-type and knock out, Welch's t-test analysis: $P<0.0001$), and (b) a complete arrest of spermatogenesis in epithelial stage IV as shown in hematoxylin-eosin stained testis sections. Massive apoptosis of spermatocytes is indicated (asterisks). The spermatogenic arrest leads to empty epididymides and azoospermia. Scale bar in upper panels, 100 μm and in lower panels, 5 μm . (St) Seminiferous tubules. (Ep) Epididymides. (c) Tubule degeneration in juvenile mice (13 d.p.p. and 18 d.p.p.) lacking SIX6OS1 and spermatogenic arrest before pachytene studied by histology. At 13 d.p.p. spermatogenesis has reached late zygotene; at 18 d.p.p. it has reached late pachytene. Spermatocyte degeneration (apoptosis is indicated by asterisks) was first seen in 18 d.p.p. *Six6os1*^{-/-}. (d) Ovaries from *Six6os1*-deficient mice show atrophy with fibrosis and depletion of follicles. Comparative histological analysis of ovaries from *Six6os1*^{-/-} and wild-type mice at 6 days (6 d.p.p.), and 4 months (4 m) of age. Asterisks indicate corpora lutea. Scale bars represent 100 μm in 4 m, and 20 μm in 6 d.p.p.

those of the SC. Based on recent progress in elucidating the organization of the SC³⁸, and on the specific interaction of SIX6OS1 with SYCE1, it seems most plausible that SYCP1 recruits CE proteins, and the nascent CE then stabilizes SYCP1 assembly. Thus, absence of CE proteins disrupts the full accumulation of SYCP1, leading to weak/discontinuous staining patterns.

Defective DSB processing in *Six6os1*^{-/-} meiotic cells. During leptotene, DSBs are generated by SPO11 and are then resected to form ssDNA ends that invade into the homologous chromosome. DSBs are marked by the presence of phosphorylated H2AX (γ -H2AX)³⁹, which is formed through phosphorylation by the kinase ATR following its recruitment by BRCA1 (ref. 37). Thus, we monitored the formation of DSBs by analyzing the presence of γ -H2AX. While γ -H2AX distribution in mutant spermatocytes resembles that of wild-type cells in early prophase I (leptonema, zygotene) (Supplementary Table 1), γ -H2AX is not restricted to sex chromosomes during pachynema (Fig. 8a). In contrast, γ -H2AX shows a moderate labelling on the chromatin of AEs in mutant pachytene-like spermatocytes (WT 23.40 ± 3.2 ; U-type 26.02 ± 5.0 ; A-type 30.74 ± 3.6 ; see Supplementary Table 2). In females, the distribution of γ -H2AX is slightly different. Oocytes at pachytene-like stage show a similar overall pattern of γ -H2AX labelling as spermatocytes (WT 20.85 ± 3.5 ; U-type

28.14 ± 9.8 ; A-type 27.01 ± 10.9 ; see Supplementary Table 2), but it is more strictly localized to their AEs (Fig. 8b). The distribution of γ -H2AX-labelling during early prophase I, and its persistence in meiotic cells during the pachynema-like stage, suggest that DSBs are generated in *Six6os1*^{-/-} meiotic cells but are not properly repaired. To better understand the processing of DSBs, we explored the kinetics and distribution of proteins involved in DSB recombination and repair. After DSBs are induced, the recombinase RAD51 is recruited to early recombination nodules and promotes homologous strand invasion⁴⁰. In wild-type zygotene spermatocytes, RAD51 assembles on numerous foci along the AEs/LEs, which are substituted by the single strand binding protein RPA and finally disappear towards pachytene, with the exception of the unsynapsed sex AEs (Fig. 8a). During early stages (leptonema), RAD51 distribution in mutants was similar to wild-type controls (Supplementary Table 1). However, in *Six6os1*^{-/-} spermatocytes at zygotene and pachytene-like stage, both RAD51 and RPA remained partially associated with the AEs (Fig. 8a; see Supplementary Tables 1 and 2). We obtained similar results when we analysed spreads from *Six6os1*^{-/-} oocytes (Fig. 8b; Supplementary Table 2).

Next, we analysed the presence of MSH4 and MLH1 foci in mutant spermatocytes. MSH4 mediates the transition after synapsis from initial to late recombination nodules. MLH1 is a component of the post-replicative mismatch repair system and

Figure 7 | *Six6os1*^{-/-} meiotic defects are not able to synapse. (a) Double labelling of spermatocyte spreads of wild-type pachytene, and *Six6os1*^{-/-} and *Syce3*^{-/-} arrested pachytene-like spermatocytes with SYCP3 (red) and SYCP1 (green). In *Six6os1*^{-/-} spermatocytes, SYCP1 does not localize to the unaligned-type (U-type) AEs but shows a very weak staining in spermatocytes with more aligned AEs (aligned-type, A-type). By direct comparison, in *Syce3*^{-/-} arrested spermatocytes SYCP1 localizes in a discontinuous pattern along AEs independent of whether or not they are closely aligned. **(b)** Double labelling of spreads of wild-type pachytene and *Six6os1*^{-/-} pachytene-like oocytes (aligned and unaligned) with SYCP3 and SYCP1. **(c)** Double labelling of spreads of wild-type pachytene and *Six6os1*^{-/-} pachytene-like spermatocytes of SYCP3 (red) and SYCE1, SYCE3, SYCE2 or TEX12 (green) (see also extended Supplementary Fig. 11 for staining in oocyte spreads). All proteins are completely absent from the AEs in *Six6os1*-deficient mice. Scale bar, 10 μm.

marks sites of future chiasmata^{41,42}. During early stages, MSH4 foci in mutants resemble those in wild-type spermatocytes. However, these foci persisted in pachytene-like arrested spermatocytes from *Six6os1*-deficient mice (Supplementary Fig. 12). Lastly, MLH1 foci were absent in *Six6os1*^{-/-} pachytene-like chromosomes (Fig. 9a), while one/two MLH1 foci per bivalent were observed in wild-type spermatocytes. Similar results were obtained in oocytes lacking SIX6OS1 (Fig. 9a), suggesting a direct function of SIX6OS1 in homologous recombination rather than in the elimination of arrested spermatocytes at the so-called pachytene checkpoint of males. To further validate this, and in light of the late arrest at mid-pachytene-like stage (H1t positive), we exposed mutant spermatocytes to the PP2A inhibitor okadaic acid to allow *in vitro* transition from pachytene to metaphase-like I (ref. 43). After okadaic acid treatment, there was a rapid induction of chromosome condensation, leading to 20 bivalents that stain for SYCP3, with the formation of at least one chiasma in the wild type (Fig. 9b). In contrast, okadaic acid-treated *Six6os1*^{-/-} spermatocytes displayed 40 free univalents, with characteristic labelling for SYCP3 (Fig. 9b). Together, our data strongly suggest that processing of recombination intermediates into MLH1-marked late recombination nodules (chiasmata) is critically dependent on SIX6OS1.

X-Y chromosome behaviour and sex body formation. In *Six6os1* mutant spermatocytes, the X and Y chromosomes are aligned in only 25.49 ± 0.06% of cells (Fig. 10a). In contrast, the degree of alignment of the sex bivalent in *Syce3* null mutants is 44.10 ± 3.17%. In mutant spermatocytes that lack aligned sex chromosomes, the sex body is not formed (see below H2AX staining Fig. 10b). The remaining fraction of *Six6os1*^{-/-} spermatocytes with aligned XY chromosomes show apparent synapsis at the PAR, but without staining for SYCP1, in contrast to the positive SYCP1 labelling of the PAR in *Syce3*^{-/-} spermatocytes (Fig. 10a). These results suggest that whilst the sex body is not formed in either mutant, the synapsis defect in the absence of SIX6OS1 is more severe than in the *Syce3* knockout.

The X and Y chromosomes show homology only along the distal pseudoautosomal region⁴⁴ of their chromosome lengths, and the remaining unsynapsed parts are subjected to meiotic sex chromosome inactivation. This is a meiotic specific process that uses the DNA damage response to recognize unsynapsed regions and reconfigure their chromatin to a silent epigenetic domain named the sex body. The act of silencing is itself dependent upon phosphorylation of histone H2AX (γ-H2AX) by ATR in a BRCA1-dependent manner⁴⁵. We performed γ-H2AX labelling of mutant spermatocytes and found moderate staining of the X and Y chromosomes in those cells showing aligned sex

Figure 8 | DSBs are generated but defectively repaired in *Six6os1*-deficient meocytes. (a) Double immunolabelling of γ -H2AX (green) with SYCP3 (red) in wild-type and *Six6os1*^{-/-} spermatocytes (left panel). In wild-type pachytene, γ -H2AX labels intensely the chromatin of the unsynapsed sex bivalent. In *Six6os1*^{-/-} pachytene-like spermatocytes γ -H2AX labelling remains in the chromatin. Double immunolabelling of SYCP3 (red) and RAD51 (green) (central panel) or RPA (green) (right panel). Both RAD51 and RPA remain associated to the AEs in *Six6os1*^{-/-} pachytene-like spermatocytes, showing a higher number of foci than wild-type pachytene. (b) Immunostaining of spread preparations of wild-type pachytene and *Six6os1*^{-/-} pachytene-like oocytes for γ -H2AX (green), RAD51 (green) and RPA (green) together with SYCP3 (red). γ -H2AX labelling in *Six6os1*^{-/-} arrested oocytes is more restricted to the AEs than in spermatocytes. Plots under each image panel represent the quantification of intensity or number of foci from wild-type and pachytene-like arrested meocytes. Welch's *t*-test analysis: **P* < 0.01; ***P* < 0.001; ****P* < 0.0001. (See numeric data at Supplementary Table 2). Scale bar, 10 μ m.

Figure 9 | SIX6OS1 is essential for the processing of intermediate recombination nodules. (a) Double immunolabelling of SYCP3 (red) with MLH1 (green). MLH1 foci are absent at the AEs/LEs of *Six6os1*^{-/-} meocytes whereas at least one focus is present along each autosomal SC in wild-type pachytene meocytes. (b) Immunostaining of SYCP3 (red) and ACA (green) in wild type and *Six6os1*^{-/-} spermatocytes. Okadaic acid-induced metaphase I plates of wild-type spermatocytes give rise to 20 bivalents each, with two opposed centromere signals (ACA) and positive staining for SYCP3, whereas *Six6os1*^{-/-} spermatocytes lead to 40 free univalents, each with an ACA signal and SYCP3 labelling the centromeric and interchromatid domain. Scale bar, 10 μ m.

chromosomes. This is in contrast with the strong labelling observed in the sex body chromatin of control spermatocytes at pachynema (Fig. 10b). Given the interplay between synapsis and DNA damage response, we directly analysed the status of 53BP1, a component of the DNA damage response that collaborates with BRCA1 in the sex body formation. In contrast to its accumulation on the unsynapsed XY⁴⁵ in wild type, 53BP1 signals were not observed in *Six6os1* mutant spermatocytes (Fig. 10c). Together, these results indicate that SIX6OS1 deficiency, similar to most asynaptic mice mutants, impedes sex body formation⁴⁶.

Discussion

Synapsis of homologous chromosomes is essential for the completion of meiosis and thus for fertility. The SC provides the structural framework for synapsis and for the processing of recombination intermediates into crossovers. Recently, the gene variant rs1254319 (p.Leu524Phe) in the anonymous *C14ORF39*/SIX6OS1 gene was identified as an influencing polymorphism affecting the human recombination rate². In addition, the same rs1254319 (p.Leu524Phe) variant has been associated with age at menarche, an indirect fertility trait⁴⁷, in a meta-analysis of 32

genome-wide association studies in 87,802 women of European descent⁴⁸. Accordingly, we show that this coding variant of *SIX6OS1* lies in a conserved residue of the SIX6OS1 protein. Cytological analysis revealed that SIX6OS1 is a new component of the CE of the SC that co-localizes with SYCE1 and SYCE3 at the synapsed chromosome axes (Fig. 1e).

By comparison of the cytological localization of CE proteins, their protein–protein interaction network, and the phenotypes obtained from their knockout mice, it has been suggested that there are two discernible subdomains within the central element. One domain, formed by SYCP1, SYCE1 and SYCE3, would act in concert through a network of interactions, specifically between SYCP1 and SYCE3, and between SYCE3 and SYCE1 (refs 49,50). The other, more inner domain of TEX12 and SYCE2 would form a separate complex by themselves. The SYCE2–TEX12 complex is an equimolar hetero-octamer, formed by the association of a SYCE2 tetramer and two TEX12 dimers²¹, and their corresponding mutant mice show some degree of synapsis with small but intense foci of SYCP1, SYCE1 and SYCE3 between their aligned AEs^{24,30,51}. Mutant spermatocytes for *Syce3* show an intermediate phenotype, with defective recruitment of CE proteins but a weak discontinuous pattern of SYCP1

Figure 10 | X-Y chromosome behaviour. (a) Double immunolabelling of SYCP3 (red) and SYCP1 (green) in wild-type pachytene, and *Six6os1*^{-/-} and *Syce3*^{-/-} pachytene-like arrested spermatocytes. Yellow letters indicate aligned/unaligned sex chromosomes in mutant spermatocytes. (b) Co-labelling of SYCP3 (red) and γ-H2AX (green) in *Six6os1*^{-/-} pachytene-like arrested spermatocytes, showing moderate staining in XY chromosomes. (c) Double immunolabelling of SYCP3 (red) and 53BP1 (green). 53BP1 signal is not observed in the XY chromosomes of mutant spermatocytes, in contrast to the strong labelling of the sex body in wild type. Scale bar, 10 μm.

staining³⁰. SIX6OS1 deficiency produces a more severe phenotype, with weaker discontinuous SYCP1 loading and also lack of recruitment of CE proteins (Fig. 7). Thus, SIX6OS1 would belong to the first subdomain (together with SYCP1, SYCE1 and SYCE3), in which mutant mice display aligned homologues and normally assembled AEs, but no CE structures at all. From a cytological perspective, the pattern of SIX6OS1 distribution along LEs in pachynema (revealed through co-localization curves and regression coefficients) is also in agreement with the continuous distribution of SYCE1 and SYCP1 along synapsed LEs^{24,30} (Fig. 2). This localization of SIX6OS1 is also congruent with the Y2H, co-immunoprecipitation and PLA results we have obtained showing its specific ability to interact and co-localize with SYCE1. Taken together with the presence of more unaligned chromosome

pairs in *Six6os1*^{-/-} than in *Syce3*^{-/-} spermatocytes, and the interdependent loading of SIX6OS1 and SYCE3 (Figs 4a,b and 7c), we suggest that SIX6OS1 is required at a similar hierarchy level (neither upstream nor downstream) to SYCE3, and downstream of SYCP1.

Despite the recent advances in reconstructing the 3D molecular organization of the mammalian SC with isotropic resolution through super-resolution imaging⁵², several gaps still remain in the net of interactors and partners involved in the assembly of the SC tripartite structure. In this regard, it has been postulated that SYCE1 stabilizes SYCP1 N-terminal interactions in the CE²⁵, suggesting that SYCE1 can act alongside SYCP1 with SYCE3. However, this view is neither validated by the phenotype of *Syce3*^{-/-} spermatocytes, in which SYCP1 stains weakly and

SYCE2/TEX12 is absent from AEs³⁰, nor by Y2H studies²¹. Recently, a direct interaction between SYCP1 and SYCE3, but not with SYCE1, SYCE2 or TEX12, has been shown by co-IP experiments and biochemical studies⁵⁰, which is in closer agreement with the genetic depletion phenotype. Similarly, with the present knowledge of interactions and components of the SC, we have no explanation for the weaker SYCP1 staining in adult *Six6os1* mutants in comparison with the *Syce3* mutant (which also lacks SIX6OS1). In this sense, the appearance of a new player in the CE family of proteins such as SIX6OS1, which is essential for the stabilization of the central region and for synapsis, deepens the complexity of the multilayered structure of the CE and suggests that unknown players could help to elucidate several open questions.

It has been shown through mouse mutants lacking CE-specific proteins that assembly of the SC central region is essential for recombination progression and chiasmata formation⁹. Similarly, *Six6os1*-deficient meiotic cells showed an arrest in the processing of recombination intermediates into MLH1-marked late recombination nodules (chiasmata). Together, these observations raise the possibility that interaction between components of the CE and recombination machinery would be critical for meiotic recombination. In this context, interactions have been described between RAD51 and both SYCP1 and SYCE2 (ref. 26). Based on sequence analysis, we predict that SIX6OS1 contains a highly helical structural region within its N terminus, followed by a flexible linker and then a C-terminal flexible protein docking sequence that could recruit multiple globular proteins to induce macromolecular protein complex. In accordance with other SC proteins, self-association and SYCE1-binding are likely mediated by the helical N-terminal domain, suggesting that this region may function in the structural assembly of the CE. The predicted unstructured nature of the remainder of the sequence suggests it may act as a flexible linker between the N-terminal structural domain and protein-protein interactions mediated by conserved patches of residues within the C terminus of the molecule. It is tempting to speculate that the predicted protein-protein interaction motifs of the C-terminal region may be responsible for the recruitment and/or stabilization of components of recombination nodules necessary for proper recombination progression. Consequently, subtle variations in the protein sequence of human SIX6OS1 (that is, rs1254319, p.Leu524Phe), could act by modifying the CO/NCO ratio, which is ultimately responsible for the observed number of recombination events genome wide. Interestingly, allele A in rs1254319 p.Leu524Phe is associated with higher recombination rate in women (53 cM) but not in men². This observation fits well with the observed sexual dimorphism in several cellular aspects of *Six6os1*-deficient mice, such as differences in the deposition of SYCP1 (Fig. 7a,b; Supplementary Fig. 9c) and difference in the frequency of U-type AEs between mutant oocytes and spermatocytes.

In summary, we have identified the biological pathway by which the SNV identified in *SIX6OS1* affects the recombination rate in humans. Our functional data show how this protein of the SC is dispensable for the generation of DSBs, but is required for the appropriate processing of intermediate recombination nodules immediately before reciprocal recombination and CO formation, and is thus essential for chromosome synapsis and fertility.

Methods

Histology. For histological analysis of adult testes, mice were perfused and their testes/ovaries were processed into serial paraffin sections and stained with hematoxylin-eosin. For histological studies of 13 and 18 day mice, testes were fixed in Bouin's fixative.

Immunocytology and antibodies. Testes were decapsulated and processed for spreading using the 'dry-down' technique. Oocytes from fetal ovaries (E17.5 embryos) were digested with collagenase, incubated in hypotonic buffer, disaggregated, fixed in paraformaldehyde and incubated with the indicated antibodies for immunofluorescence. Goat polyclonal antibodies against C14ORF39/SIX6OS1 were developed by Santa Cruz (sc-245304) and used for the immunofluorescence analysis. This antibody was raised against a conserved internal region of human SIX6OS1. Rabbit polyclonal antibodies against SIX6OS1 were developed by Proteintech (22664-1-AP) against a fusion protein of GST with SIX6OS1 (C-350 aa) of human origin (see Supplementary Fig. 3 for validation) and was used to validate the immunofluorescence results obtained with the goat polyclonal antibody against C14ORF39/SIX6OS1 developed by Santa Cruz. The primary antibodies used for immunofluorescence were rabbit α SMC3 serum K987 (1:20), rabbit α SMC1 β serum K974 (1:20), rabbit α STAG3 serum K403, α REC8 serum K1019, rabbit α RAD21 IgG K854 (1:5)⁴⁵, mouse α SYCP3 IgG sc-74569 (1:100), rabbit α RAD51 sc-8349 (1:30) and PC130 (1:50), rabbit α SYCP1 IgG ab15090 (1:200) (Abcam), rabbit anti- γ H2AX (ser139) IgG #07-164 (1:200) (Millipore), ACA or purified human α -centromere proteins IgG 15-235 (1:5, Antibodies Incorporated), mouse α MLH1 51-1327GR (1:5, BD Biosciences), rabbit α 53BP1 sc-22760 (1:20), rabbit α RAP1 IgG (1:400, provided by Dr Titia de Lange, The Rockefeller University, USA), and rabbit α RPA IgG (1:300, provided by Dr E. Marcon, Toronto University, Canada), rabbit α TEX12 IgG (1:100) and guinea pig α SYCE3(1:20) (provided by Dr R. Benavente, University of Würzburg, Germany), guinea pig α SYCE1 (1:100), rabbit α SYCE1 (Proteintech), guinea pig α SYCE2 (1:50) (provided by C. Höög, Karolinska Institutet, Sweden) and guinea pig α H1t (Provided by MA Handel). The secondary antibodies used were TRITC α -mouse 115-095-146/ α -rabbit 111-025-144 and FITC α -mouse 115-095-146/ α -rabbit 111-095-045 (Jackson ImmunoResearch) (all 1:100). Slides were visualized at room temperature using a microscope (Axioplan 2; Carl Zeiss, Inc.) with 63 $\times$ objectives with an aperture of 1.4 (Carl Zeiss, Inc.). Images were taken with a digital camera (ORCA-ER; Hamamatsu) and processed with OPENLAB 4.0.3 and Photoshop (Adobe). Quantification of γ H2AX and SYCP1 fluorescence signals was performed using Image J software. Chromosome counts of A-type and U-type cells were performed on at least 100 pachytene-like spermatocytes and oocytes from three individuals.

In vivo electroporation. Testes were freed from the abdominal cavity and 10 μ l of DNA solution (50 μ g) mixed with 1 μ l of 10 $\times$ FastGreen (Sigma Aldrich F7258) was injected in the rete testis with a DNA embryo microinjection tip. After a period of 1 h following the injection, testes were held between a pair of electrodes and electric pulses were applied four times (35 V for 50 ms each pulse) using a CUY21 BEX electroporator²³.

Electron microscopy. For immunoelectron microscopy, 10 μ m cryosections of mouse testis were fixed with acetone for 10 min at -20°C and air dried. Incubation with primary antibodies was carried out in a humidified box for 4 h at room temperature. After rinsing twice in PBS, sections were fixed for 10 min in 2% formaldehyde and blocked with 50 mM NH_4Cl . Secondary antibodies conjugated to 6 nm gold particles were incubated overnight at 4°C , and samples were subsequently washed in PBS. Samples were fixed for 30 min in 2.5% glutaraldehyde and postfixed in 2% osmium tetroxide. After rinsing three times with H_2O , samples were dehydrated in an ethanol series and embedded in Epon. Ultrathin sections were stained with uranyl acetate and lead citrate according to standard procedures³⁰.

Okadaic acid assay. Testes were dissected into a Petri dish containing ice cold sterile medium (4 mM L-glutamine, 10% fetal calf serum, and 25 mM Hepes in Dulbecco's Modified Eagle's medium) and cell suspensions (5×10^6 cells per ml) were exposed to 5 μM okadaic acid (Sigma-Aldrich) for 5 h at 32°C and 5% CO_2 before spreading the cells by the dry down procedure⁴.

Generation of plasmids. Full-length cDNAs encoding SIX6OS1, TEX12, SYCE1, SYCE2, SYCE3, SYCP1 and SYCP3 were RT-PCR amplified from murine testis RNA. Full-length cDNAs were cloned into the pcDNA3, pcDNA3 x2Flag, pCEFL HA or pcDNA3.1 Myc-His (-) or pEGFP-C1 mammalian expression vectors.

Cell lines and transfections. HEK 293T and COS7 cell lines were transfected with Lipofectamine (Invitrogen) or Jetpei (PolyPlus) and obtained from the ATCC. Cell lines were tested for mycoplasma contamination (Mycoplasma PCR ELISA, Sigma).

Immunoprecipitation and proximity ligation assay. HEK 293T cells were transiently transfected and whole cell extracts were prepared and cleared with protein G Sepharose beads (GE Healthcare) for 1 h. The antibody was added for 2 h and immunocomplexes were isolated by adsorption to protein G Sepharose beads for 1 h. After washing, beads were loaded onto reducing 10% polyacrylamide SDS gels and proteins were detected by western blotting with the indicated antibodies. Immunoprecipitations were performed using mouse α Flag IgG (5 μ g; F1804, Sigma-Aldrich), rabbit α Myc Tag IgG (4 μ g; #06-549, Millipore), mouse α HA.11

IgG MMS- (5 μ l, approx. 10 μ g per 1 mg prot; 101R, Covance), goat α GFP IgG (4 μ g; sc-5385, Santa Cruz), ChromPure mouse IgG (5 μ g/1 mg prot; 015-000-003), ChromPure rabbit IgG (5 μ g per 1 mg prot; 011-000-003, Jackson ImmunoResearch), ChromPure goat IgG (5 μ g per 1 mg prot; 005-000-003, Jackson ImmunoResearch). Primary antibodies used for western blotting were mouse α Flag IgG (P1804, Sigma-Aldrich) (1:10,000), rabbit α HA IgG (H6908, Sigma-Aldrich) (1:1,000), rabbit α Flag IgG (1:800; F7425 Sigma-Aldrich), mouse α Myc obtained from hybridoma cell myc-1-9E10.2 ATCC (1:1,000). Secondary horseradish peroxidase-conjugated α -mouse (NA931V, GE Healthcare), α -rabbit (#7074, Cell Signaling), or α -goat (A27014, Thermo Scientific) antibodies were used at 1:10,000, 1:3,000 or 1:10,000 dilution, respectively. Antibodies were detected by using Immobilon Western Chemiluminescent HRP Substrate from Millipore. Proximity Ligation Assay was performed using goat α SIX6OS1 (sc-5385) and rabbit α SYCE1, with the corresponding anti-goat PLA Probe PLUS and anti-rabbit PLA probe MINUS, following the manufacturer's instructions (Duolink Using PLA Technology, SIGMA).

The uncropped versions of western blots in Fig. 3 are shown in Supplementary Fig. 13.

Production of CRISPR/Cas9-edited mice. *Six6os1*-gRNAs (G68 5'-CACCGATCTGTTTGTTCAGTTTGGAC-3' and 5'-AAACGTCCAACTGACAAAC AGATC-3' and G75 5'-CACCCTACTTATGTCTT GTCATAC-3' and 5'-AAACGTATGACAAAGACATAAGTAC-3' targeting exon 2 and exon 3 were predicted at crispr.mit.edu. *Six6os1*-sgRNAs were produced by cloning annealed complementary oligos at the BbsI site of pX330 (#42230, Addgene), generating PCR products containing a T7 promoter sequence that were purified (NZYtech), and then performing *in vitro* transcription using the MEGAshortscript T7 Transcription Kit (Life Technologies). The plasmid pST1374-NLS-flag-linker-Cas9 (#44758; Addgene) was used for generating Cas9 mRNA after linearization with AgeI. *In vitro* transcription and capping were performed using the mMESSAGE mMACHINE T7 Transcription Kit (AM1345; Life Technologies). Products were purified using the RNeasy Mini Kit (Qiagen). RNA (100 ng μ l⁻¹ Cas9 and 50 ng μ l⁻¹ each guide RNA) was microinjected into zygotes (F1 hybrids between strains C57BL/6J and CBA/J)⁵³. Edited founders were identified by PCR amplification (Taq polymerase, NZYtech) with primers flanking exons 2 and 3 (Primer F 5'-CACTTACATTTTCTCTTTAAGAATGC-3' and R 5'-CCCTCTCATACATCAAGTTGC-3') and subcloned into pBlueScript (Stratagene) followed by standard Sanger sequencing. The length of the corresponding wild-type and mutant allele were 413 and 289 bp, respectively. The selected founder was crossed with wild-type C57BL/6J to eliminate possible unwanted off-targets and to generate pure heterozygous. *Six6os1*^{+/-} heterozygous mice were sequenced again by Sanger sequencing and crossed to give rise to *Six6os1*^{-/-} homozygous. Genotyping was performed by agarose gel electrophoresis analysis of PCR products produced from DNA isolated from tail biopsy specimens. Mouse mutants for *Rec8*, *Rad21*, *Sycp3*, *Sycp1* and *Stag3* have been previously developed^{4,13,30-32}.

Mice were housed in a temperature-controlled facility (specific pathogen free, spf) using individually ventilated cages, standard diet and a 12h light-dark cycle, according to European Union regulations at the 'Servicio de Experimentación Animal, SEA'. Mouse protocols were approved by the Ethics Committee for Animal Experimentation of the University of Salamanca (USAL). We made every effort to minimize suffering and to improve animal welfare. Blinded experiments were not possible since the phenotype was very obvious between wild-type and *Six6os1*-deficient mouse for all of the experimental procedures used. No randomization methods were applied since the animals were not divided in groups/treatments. The minimum size used for each analysis was three animals/genotype. The mice analysed were between 2 and 4 months of age, except in those experiments where is indicated.

Quantitative PCR. Total RNA was isolated from various tissues of wild-type adult mice. To analyse the expression of *Six6os1* and *Rad21* mRNAs, equal amounts of cDNA were synthesized using SuperScript II Reverse Transcriptase (Invitrogen, Life Technologies) and Oligo (dT). qPCR was performed using FastStart Universal SYBR Green Master Mix (ROX) (Roche) and specific forward and reverse primers: qSIX6OS1_F 5'-GCTGAATGTGGAGATAAAGAG-3' and qSIX6OS1_R 5'-AGGAGTTTCAGGAGTTTGAGG-3'; qRAD21L_F 5'-TTGCAGCTCACTGGGAGAGA-3' and qRAD21L_R 5'-AGTCCTGGGGCGAAATGTCATC-3'. All qPCR reactions were performed at 95 °C for 10 min, and then 40 cycles of 95 °C for 15 s and 62 °C for 1 min on the iQ5 Thermal Cycler (Bio-Rad). β -Actin was amplified as a housekeeping gene with the primers q β -actin_F 5'-GGCACCACACCTTCTACAATG-3' and q β -actin_R 5'-GTGGTGGTGAAGCTGTAGCC-3'.

Y2H assay and screening. Y2H assay was performed using the Matchmaker Gold Yeast Two-Hybrid System (Clontech) according to the manufacturers' instructions. Mouse *Six6os1* cDNA encoding the N terminus (1-138) was subcloned into the vector pGBKT7 and was used as bait to screen a mouse testis Mate & Plate cDNA library (Clontech Laboratories Inc.). Positive clones were initially identified on double dropout SD (synthetic dropout)/-Leu/-Trp/X- α -Gal/Aureobasidin A plates before further selection on higher stringency quadruple dropout SD/-Ade/-His/-Leu/-Trp/X- α -Gal/Aureobasidin A plates. Prey plasmids were extracted from the candidate yeast clones and transformed into *Escherichia coli*. The plasmids from two

independent bacteria colonies were independently grown, extracted and Sanger sequenced. Southern blotting was also used for plasmid screening.

Sequence analysis. Protein sequences were extracted from the UniProt database and analysed using Jalview 2 (ref. 54). Multiple sequence alignments and secondary structure predictions were performed using MUSCLE (EBI)⁵⁵ and Jpred 4 (ref. 56), respectively.

Co-localization profile. SIX6OS1 and either SYCP1, SYCE1, SYCE3, SYCE2 or TEX12 were stained on spreads of wild-type spermatocytes. Images were captured with identical camera settings. Fluorescence signals were measured along the 19 autosomal AEs of pachytene cells using the 'Plot profile' tool of ImageJ. Signal intensities were standardized, acquiring values between -1 and 1, and the overlay profiles of SIX6OS1 and other CE proteins were plotted. Regression analysis for each pair of proteins was performed to determine the correlation between their profiles. The values of the coefficients of determination R^2 are shown in the scatter plots.

Statistics. To compare counts between genotypes at different stages, we used the Welch's *t*-test (unequal variances *t*-test), which was appropriate as the count data were not highly skewed (that is, were reasonably approximated by a normal distribution) and in most cases showed unequal variance. Asterisks denote statistical significance: **P* value <0.01, ***P* value <0.001 and ****P* value <0.0001.

Data availability. Genomic DNA sequences of *H. sapiens* (human, 317761), *M. musculus* (mouse, 75801) are available on GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>). Amino acid sequences of *H. sapiens* (Q8N1H7), *M. musculus* (NP_083381), *P. troglodytes* (Chimp, H2Q8E6), *S. charissii* (Tasmanian devil, G3WQ57), *O. anatinus* (Latypus, F6ZZ02), *P. sinensis* (Chinese turtle, K7GAG2), *G. gallus* (Chick, E1C952) and *L. chalumnae* (West India coelacanth, M3XIB0) were obtained from the UniProt database (<http://www.uniprot.org/>). All remaining data generated in this study are available in the Article and Supplementary Information files or available from the authors upon request from the authors.

References

- Handel, M. A. & Schimenti, J. C. Genetics of mammalian meiosis: regulation, dynamics and impact on fertility. *Nat. Rev. Genet.* **11**, 124–136 (2010).
- Kong, A. *et al.* Common and low-frequency variants associated with genome-wide recombination rate. *Nat. Genet.* **46**, 11–16 (2014).
- Gutierrez-Caballero, C. *et al.* Identification and molecular characterization of the mammalian alpha-kleisin RAD21L. *Cell Cycle* **10**, 1477–1487 (2011).
- Herran, Y. *et al.* The cohesin subunit RAD21L functions in meiotic synapsis and exhibits sexual dimorphism in fertility. *EMBO J.* **30**, 3091–3105 (2011).
- Llano, E. *et al.* Meiotic cohesin complexes are essential for the formation of the axial element in mice. *J. Cell Biol.* **197**, 877–885 (2012).
- Hartsuiker, E., Vaessen, E., Carr, A. M. & Kohli, J. Fission yeast Rad50 stimulates sister chromatid recombination and links cohesion with repair. *EMBO J.* **20**, 6660–6671 (2001).
- Klein, P. *et al.* A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis. *Cell* **98**, 91–103 (1999).
- Zickler, D. & Kleckner, N. Recombination, pairing, and synapsis of homologs during meiosis. *Cold Spring Harb. Perspect. Biol.* **7**, 1–26 (2015).
- Fraune, J., Schramm, S., Alsheimer, M. & Benavente, R. The mammalian synaptonemal complex: protein components, assembly and role in meiotic recombination. *Exp. cell Res.* **318**, 1340–1346 (2012).
- Henderson, K. A. & Keeney, S. Synaptonemal complex formation: where does it start? *BioEssays* **27**, 995–998 (2005).
- Baudat, F., Imai, Y. & de Massy, B. Meiotic recombination in mammals: localization and regulation. *Nat. Rev. Genet.* **14**, 794–806 (2013).
- de Vries, L. *et al.* Exome sequencing reveals SYCE1 mutation associated with autosomal recessive primary ovarian insufficiency. *J. Clin. Endocrinol. Metab.* **99**, E2129–E2132 (2014).
- Caburet, S. *et al.* Mutant cohesin in premature ovarian failure. *N. Engl. J. Med.* **370**, 943–949 (2014).
- Shinohara, M., Oh, S. D., Hunter, N. & Shinohara, A. Crossover assurance and crossover interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nat. Genet.* **40**, 299–309 (2008).
- Reynolds, A. *et al.* RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis. *Nat. Genet.* **45**, 269–278 (2013).
- Baudat, F. *et al.* PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* **327**, 836–840 (2010).
- Kneitz, B. *et al.* MutS homolog 4 localization to meiotic chromosomes is required for chromosome pairing during meiosis in male and female mice. *Genes Dev.* **14**, 1085–1097 (2000).
- Ward, J. O. *et al.* Mutation in mouse hei10, an e3 ubiquitin ligase, disrupts meiotic crossing over. *PLoS Genet.* **3**, e139 (2007).

19. Alfano, G. *et al.* Natural antisense transcripts associated with genes involved in eye development. *Hum. Mol. Genet.* **14**, 913–923 (2005).
20. Consortium, G.T. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**, 648–660 (2015).
21. Davies, O. R., Maman, J. D. & Pellegrini, L. Structural analysis of the human SYCE2-TEX12 complex provides molecular insights into synaptonemal complex assembly. *Open Biol.* **2**, 120099 (2012).
22. Syrjänen, J. L., Pellegrini, L. & Davies, O. R. A molecular model for the role of SYCP3 in meiotic chromosome organisation. *eLife* **3**, e02963 (2014).
23. Shibuya, H., Morimoto, A. & Watanabe, Y. The dissection of meiotic chromosome movement in mice using an *in vivo* electroporation technique. *PLoS Genet.* **10**, e1004821 (2014).
24. Hamer, G. *et al.* Characterization of a novel meiosis-specific protein within the central element of the synaptonemal complex. *J. Cell Sci.* **119**, 4025–4032 (2006).
25. Costa, Y. *et al.* Two novel proteins recruited by synaptonemal complex protein 1 (SYCP1) are at the centre of meiosis. *J. Cell Sci.* **118**, 2755–2762 (2005).
26. Bolcun-Filas, E. *et al.* Mutation of the mouse Syce1 gene disrupts synapsis and suggests a link between synaptonemal complex structural components and DNA repair. *PLoS Genet.* **5**, e1000393 (2009).
27. Gomez, R. *et al.* Sororin loads to the synaptonemal complex central region independently of meiotic cohesin complexes. *EMBO Rep.* **17**, 695–707 (2016).
28. Lu, J. *et al.* Structural insight into the central element assembly of the synaptonemal complex. *Sci. Rep.* **4**, 7059 (2014).
29. Winkler, K., Alsheimer, M., Ollinger, R. & Benavente, R. Protein SYCP2 provides a link between transverse filaments and lateral elements of mammalian synaptonemal complexes. *Chromosoma* **118**, 259–267 (2009).
30. Schramm, S. *et al.* A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility. *PLoS Genet.* **7**, e1002088 (2011).
31. de Vries, F. A. *et al.* Mouse Sycp1 functions in synaptonemal complex assembly, meiotic recombination, and XY body formation. *Genes Dev.* **19**, 1376–1389 (2005).
32. Bannister, L. A., Reinholdt, L. G., Munroe, R. J. & Schimenti, J. C. Positional cloning and characterization of mouse mei8, a disrupted allele of the meiotic cohesin Rec8. *Genesis* **40**, 184–194 (2004).
33. Llano, E. *et al.* STAG3 is a strong candidate gene for male infertility. *Hum. Mol. Genet.* **23**, 3421–3431 (2014).
34. Xu, H., Beasley, M. D., Warren, W. D., van der Horst, G. T. & McKay, M. J. Absence of mouse REC8 cohesin promotes synapsis of sister chromatids in meiosis. *Dev. Cell* **8**, 949–961 (2005).
35. Inselman, A., Eaker, S. & Handel, M. A. Temporal expression of cell cycle-related proteins during spermatogenesis: establishing a timeline for onset of the meiotic divisions. *Cytogenet. Genome Res.* **103**, 277–284 (2003).
36. Hamer, G. *et al.* Progression of meiotic recombination requires structural maturation of the central element of the synaptonemal complex. *J. Cell Sci.* **121**, 2445–2451 (2008).
37. Turner, J. M. *et al.* Silencing of unsynapsed meiotic chromosomes in the mouse. *Nat. Genet.* **37**, 41–47 (2005).
38. Schucker, K., Holm, T., Franke, C., Sauer, M. & Benavente, R. Elucidation of synaptonemal complex organization by super-resolution imaging with isotropic resolution. *Proc. Natl Acad. Sci. USA* **112**, 2029–2033 (2015).
39. Rogakou, E. P., Pilch, D. R., Orr, A. H., Ivanova, V. S. & Bonner, W. M. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J. Biol. Chem.* **273**, 5858–5868 (1998).
40. Mimitou, E. P. & Symington, L. S. Nucleases and helicases take center stage in homologous recombination. *Trends Biochem. Sci.* **34**, 264–272 (2009).
41. Santucci-Darmanin, S., Vidal, P., Scimeca, J. C., Turc-Carel, C. & Paquis-Flucklinger, V. Family of SRY/Sox proteins is involved in the regulation of the mouse Msh4 (MutS Homolog 4) gene expression. *Mol. Reprod. Dev.* **60**, 172–180 (2001).
42. Moens, P. B., Marcon, E., Shore, J. S., Kochakpour, N. & Spyropoulos, B. Initiation and resolution of interhomolog connections: crossover and non-crossover sites along mouse synaptonemal complexes. *J. Cell Sci.* **120**, 1017–1027 (2007).
43. Wiltshire, T., Park, C., Caldwell, K. A. & Handel, M. A. Induced premature G2/M-phase transition in pachytene spermatocytes includes events unique to meiosis. *Dev. Biol.* **169**, 557–567 (1995).
44. Ferguson-Smith, M. A. X-Y chromosomal interchange in the aetiology of true hermaphroditism and of XX Klinefelter's syndrome. *Lancet* **2**, 475–476 (1966).
45. Broering, T. J. *et al.* BRCA1 establishes DNA damage signaling and pericentric heterochromatin of the X chromosome in male meiosis. *J. Cell Biol.* **205**, 663–675 (2014).
46. Burgoyne, P. S., Mahadevaiah, S. K. & Turner, J. M. The consequences of asynapsis for mammalian meiosis. *Nat. Rev. Genet.* **10**, 207–216 (2009).
47. van Noord, P. A., Dubas, J. S., Dorland, M., Boersma, H. & te Velde, E. Age at natural menopause in a population-based screening cohort: the role of menarche, fecundity, and lifestyle factors. *Fertil. Steril.* **68**, 95–102 (1997).
48. Elks, C. E. *et al.* Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nat. Genet.* **42**, 1077–1085 (2010).
49. Fraune, J., Wiesner, M. & Benavente, R. The synaptonemal complex of basal metazoan hydra: more similarities to vertebrate than invertebrate meiosis model organisms. *J. Genet. Genomics* **41**, 107–115 (2014).
50. Hernandez-Hernandez, A. *et al.* The central element of the synaptonemal complex in mice is organized as a bilayered junction structure. *J. Cell Sci.* **129**, 2239–2249 (2016).
51. Bolcun-Filas, E. *et al.* SYCE2 is required for synaptonemal complex assembly, double strand break repair, and homologous recombination. *J. Cell Biol.* **176**, 741–747 (2007).
52. Fraune, J. *et al.* Hydra meiosis reveals unexpected conservation of structural synaptonemal complex proteins across metazoans. *Proc. Natl Acad. Sci. USA* **109**, 16588–16593 (2012).
53. Singh, P., Schimenti, J. C. & Bolcun-Filas, E. A mouse geneticist's practical guide to CRISPR applications. *Genetics* **199**, 1–15 (2015).
54. Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009).
55. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
56. Drozdetskiy, A., Cole, C., Procter, J. & Barton, G. J. JPred4: a protein secondary structure prediction server. *Nucleic Acids Res.* **43**, W389–W394 (2015).

Acknowledgements

We wish to express our sincere thanks to Drs Lu, Höög, Titia de Lange, Schimenti and Handel for providing antibodies and reagents (mice), and to Dr JL de la Pompa for reviewing the MS and Dr Jessberger for useful help. This work was supported by BFU 2014-59307-R, MEIONet and JCyLe (CSI052U16). LGH and NFM are supported by European Social Fund/JCyLe grants (EDU/1083/2013 and EDU/310/2015). ORD is a Sir Henry Dale Fellow jointly funded by the Wellcome Trust and Royal Society (Grant Number 104158/Z/14/Z). RB is funded by DFG (grant Be1168/8-1). AT and ID were supported by DFG grants TO421/8-2 and TO421/6-1, respectively.

Author contributions

L.G.H. and N.F.M. performed the characterization of the mutant mice including the cytological and biochemical analysis. M.S.M. carried out the Cas9 injections. O.R.D. carried out the protein analysis. I.R. carried out infertility phenotyping of mutant mice. I.G.T. contributed with the initial co-IP experiments. DdR performed the staging of the seminiferous tubules. J.L.B. contributed with reagents and discussion. R.B. performed the E.M. work and contributed with the Syce3 KO samples and discussion of the results. E.L.C. performed the Y2H analysis. I.D. and A.T. provided spermatocytes spreads from Sycp1 KO. A.M.P. and E.L.C. designed the experiments and wrote the paper with the input of the remaining authors.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Gómez-H, L. *et al.* C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nat. Commun.* **7**, 13298 doi: 10.1038/ncomms13298 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016

RESEARCH ARTICLE

The PSMA8 subunit of the spermatoproteasome is essential for proper meiotic exit and mouse fertility

Laura Gómez-H¹, Natalia Felipe-Medina¹, Yazmine B. Condezo¹, Rodrigo Garcia-Valiente¹, Isabel Ramos¹, José Angel Suja², José Luis Barbero³, Ignasi Roig⁴, Manuel Sánchez-Martín⁵, Dirk G. de Rooij^{6,7}, Elena Llano^{1,8*}, Alberto M. Pendas^{1*}

1 Molecular Mechanisms Program, Centro de Investigación del Cáncer and Instituto de Biología Molecular y Celular del Cáncer (CSIC-Universidad de Salamanca), Salamanca, Spain, **2** Unidad de Biología Celular, Universidad Autónoma de Madrid, Madrid, Spain, **3** Centro de Investigaciones Biológicas (CSIC), Madrid, Spain, **4** Genome Integrity and Instability Group, Institut de Biotecnologia i Biomedicina, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, Spain, **5** Departamento de Medicina, Universidad de Salamanca, Salamanca, Spain, **6** Reproductive Biology Group, Division of Developmental Biology, Department of Biology, Faculty of Science, Utrecht University, Utrecht, The Netherlands, **7** Center for Reproductive Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands, **8** Departamento de Fisiología y Farmacología, Universidad de Salamanca, Salamanca, Spain

* ellano@usal.es (EL); amp@usal.es (AMP)

OPEN ACCESS

Citation: Gómez-H L, Felipe-Medina N, Condezo YB, Garcia-Valiente R, Ramos I, Suja JA, et al. (2019) The PSMA8 subunit of the spermatoproteasome is essential for proper meiotic exit and mouse fertility. *PLoS Genet* 15(8): e1008316. <https://doi.org/10.1371/journal.pgen.1008316>

Editor: Mary Ann Handel, The Jackson Laboratory, USA, UNITED STATES

Received: February 21, 2019

Accepted: July 17, 2019

Published: August 22, 2019

Copyright: © 2019 Gómez-H et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Amino acid sequences of PSMA8 from *H. sapiens* (Q8TAA3) and *M. musculus* (Q9CWH6) and PSMA7 from *H. sapiens* (O14818) and from *M. musculus* (Q9Z2U0) were obtained from the UniProt database (<http://www.uniprot.org/>). Searches were generated versus the *Mus musculus* proteome (UP000000589, May 2017 release) and Maxquant contaminants. PSMA8 expression was obtained at <https://gtexportal.org/home/gene/PSMA8>. All

Abstract

The ubiquitin proteasome system regulates meiotic recombination in yeast through its association with the synaptonemal complex, a ‘zipper’-like structure that holds homologous chromosome pairs in synapsis during meiotic prophase I. In mammals, the proteasome activator subunit PA200 targets acetylated histones for degradation during somatic DNA double strand break repair and during histone replacement during spermiogenesis. We investigated the role of the testis-specific proteasomal subunit $\alpha 4s$ (PSMA8) during spermatogenesis, and found that PSMA8 was localized to and dependent on the central region of the synaptonemal complex. Accordingly, synapsis-deficient mice show delocalization of PSMA8. Moreover, though *Psm8*-deficient mice are proficient in meiotic homologous recombination, there are alterations in the proteostasis of several key meiotic players that, in addition to the known substrate acetylated histones, have been shown by a proteomic approach to interact with PSMA8, such as SYCP3, SYCP1, CDK1 and TRIP13. These alterations lead to an accumulation of spermatocytes in metaphase I and II which either enter massively into apoptosis or give rise to a low number of aberrant round spermatids that apoptose before histone replacement takes place.

Author summary

Proteins within the cells that are unnecessary or damaged are degraded by a large protein complex named the proteasome. The proteins to be degraded are marked by a small protein called ubiquitin. The addition of a small modification (acetyl group) to some proteins also promotes their degradation by the proteasome. Proteasomal degradation of proteins

remaining data generated in this study are available in the Article and the Supporting Information files.

Funding: This work was supported by MINECO (BFU2017-89408-R) and by Junta de Castilla y León (CSI239P18). LGH and NFM are supported by European Social Fund/JCyLe grants (EDU/1083/2013 and EDU/310/2015) and YBC by a FPI grant from the MINECO (BS-2015-073993). IR was supported by MINECO (BFU2016-80370-P). JAS was supported by MINECO (BFU2014-53681-P). We appreciate the help of M^a Luz Sánchez García for the FACs analysis. The proteomic analysis was performed in the Proteomics Facility of Centro de Investigación del Cáncer, Salamanca, Grant PRB3 (IPT17/0019 - ISCIII-SGEFI / ERDF). CIC-IBMCC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia cofunded by the Castilla-León autonomous government and the European Regional Development Fund (CLC-2017-01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

is an essential mechanism for many developmental programs including gametogenesis, a process whereby a diploid cell produces a haploid cell or gamete (sperm or egg). The mechanism by which this genome reduction occurs is called meiosis. Here, we report the study of a protein, named PSMA8 that is specific for the testis proteasome in vertebrates. Using the mouse as a model, we show that loss of PSMA8 leads to infertility in males. By co-immunoprecipitation-coupled mass spectroscopy we identified a large list of novel PSMA8 interacting proteins. We focused our functional analysis on several key meiotic proteins which were accumulated such as SYCP3, SYCP1, CDK1 and TRIP13 in addition to the known substrate of the spermatoproteasome, the acetylated histones. We suggest that the altered accumulation of these important proteins causes a disequilibrium of the meiotic division that produces apoptotic spermatocytes in metaphase I and II and also early spermatids that die soon after reaching this stage.

Introduction

Intracellular protein content is controlled through the balance between the rates of their synthesis and degradation. In eukaryotic cells, the bulk of the degradation is carried out by the ubiquitin-proteasome system (UPS). The proteasome is a multi-subunit complex that eliminates proteins, typically labeled with ubiquitin, by ATP-driven proteolysis [1]. Proteasome complexes comprise a cylindrical catalytic core particle (CP, 20S) and different regulatory particles (RPs, 19S) that regulate the access to the CP by capping it at either end [2]. The CP is composed of seven α -type subunits and seven β -type subunits arranged as a cylinder of four rings (α 1–7, β 1–7, β 1–7, α 1–7) [1, 3]. RPs are composed of 20 subunits and their association with the CP is ATP-dependent. There are four additional activators, the 11S regulator PA28 α / β / γ and the ubiquitous PA200 (*Psme4*) regulator that stimulates protein degradation independently of ubiquitin [4] and plays a main role in acetylation-dependent degradation of somatic core histones during DNA repair and spermiogenesis [5, 6]. Hybrid proteasomes enclosing a RP at one end and an activator at the other end are also possible [7]. In addition, there are paralogs for three β -genes that are expressed only in the immunological system, which constitutes the immunoproteasome [8], and one β 5t gene expressed exclusively in the thymus, which constitutes the thymoproteasome [9]. Finally, there is a meiotic paralog of the α 4 subunit (*Psma7*), named α 4s (*Psma8*) [10], which might provide substrate specificity and heterogeneity to the α 4s-containing proteasome.

The proteolytic activity of the proteasome is regulated by the rate of protein ubiquitylation, but also by its association with E3 ubiquitin ligases and deubiquitinating enzymes that edit their potential substrates [11, 12]. The classical targets of the UPS are misfolded or damaged proteins and/or short-lived regulatory proteins, whose concentration is regulated by fine-tuning of their synthesis and degradation kinetics [13, 14]. Typical examples of the latter proteins are cyclins [15, 16]. More recently, it has been hypothesized but not proven that the ZMM complex (also known as the synapsis initiation complex) involved in meiotic homologous recombination is similarly regulated in the mouse [17, 18].

Meiosis is a fundamental process in sexually reproducing species that ensures the production of genetic diversity and the generation of haploid gametes from diploid progenitors [19]. This reduction in genome content is achieved by the physical connections between homologs by chiasmata [20], which are mediated by the repair of self-induced DNA double-strand breaks (DSBs) as crossing-overs (COs). Meiotic recombination takes place on proteinaceous core structures or axial elements (AEs) that scaffold the chromosomal DNA content and

physically connect (synapse) homologs through the assembly of the synaptonemal complex (SC) during prophase I [21].

The UPS regulates meiotic recombination in yeast and mouse *via* its physical association to AEs [17, 22]. Given the unknown function that the α 4s-containing proteasome plays during spermatogenesis, we explored its function in the mouse. In this study, we show that PSMA8 is localized to and dependent on the central element of the SC, and promotes the assembly of the proteasome activator PA200. Accordingly, synapsis-deficient mice show delocalization of PSMA8. Also, *Psm8*-deficient mice are proficient in meiotic homologous recombination, but show alterations in the proteostasis of several key meiotic players including acetylated histones, SYCP3, SYCP1, CDK1 and TRIP13, which in turn leads to an aberrant meiotic exit, accumulation of apoptotic spermatocytes in metaphase I and II, and finally early spermatid arrest long before histone replacement takes place.

Results

PSMA8 is expressed in spermatocytes and its localization to the SC is dependent on synapsis

Psm8 mRNA expression in mouse tissues is almost exclusively restricted to the testis (GTEx database [23] and previous studies [10]). To elucidate the cell type in which PSMA8 is expressed, we examined by western blotting testis extracts at various postnatal ages during the first wave of spermatogenesis, which progresses more synchronously than in adult mice. PSMA8 expression (using a specific antibody against the PSMA8 C-terminus [10], see Fig 1A) was first detected at P12 and increased from P14 to P20. We also used a PSMA8-R2 antibody raised against the entire recombinant PSMA8 protein, which detected the expression of both PSMA7 (already apparent at P8, before meiosis has started) and PSMA8 (Fig 1A and S1 Fig). Analysis of testis cell lines (including spermatogonium GC1-spg, Leydig cell TM3, and Sertoli cell TM4 lines), revealed the expression of PSMA7 but not PSMA8 (Fig 1A). These results indicate that its expression is restricted to cells undergoing meiosis.

To explore the subcellular localization of PSMA8, we employed the R2 antibody (PSMA7/8) since the PSMA8 C-terminus antibody did not produce any specific labeling. Double immunolabeling of PSMA8 with the AE protein SYCP3 or with SYCP1, the transverse filament protein essential for synapsis (Fig 1B and S2 Fig), revealed PSMA7/8 presence at the central region of the synaptonemal complex (super resolution imaging, Fig 1B). We validated this localization by *in vivo* electroporating [24] an expression plasmid encoding GFP-PSMA8 in the testis (Fig 1C). These results agree with the recent localization of the proteasome to the chromosome axes [17].

To investigate the possible dependence of PSMA8 localization on synapsis, we analyzed synaptic mutants with mild (*Rec8*^{-/-} [25]) and severe (*Six6os1*^{-/-} [24]) phenotypes. Mutants for the meiotic cohesin REC8 show pseudo-synapsis between sister chromatids [25], and PSMA8 was detected at these atypical synapsed-like regions (Fig 1D). In mice lacking the novel central element protein SIX6OS1, in which AEs are physically separated and unsynapsed at pachynema [24], PSMA8 signal was not restricted to their AEs and showed a broader and more disperse labeling (Fig 1D). These results indicate that PSMA8 localization to the SC central region is consequently dependent on the assembly of the SC.

Male mice lacking PSMA8 are infertile

To study the role of PSMA8, we generated a targeted mutation in exon 1-intron 1 of the murine *Psm8* gene by CRISPR/Cas9 genome editing (S3A and S3B Fig). Homozygous mutant

Fig 1. Expression analysis and localization of PSMA8 in the mouse. (A) Western blot analysis of protein extracts from mouse testis (from P8 to adult) and cell lines (TM3, TM4 and GC1) with a specific antibody against the C-terminal ($\alpha 4s$) and whole recombinant PSMA8 protein (PSMA8-R2). β -Actin was used as loading control. The corresponding bands to PSMA8 and PSMA7 are indicated in the right of the panel. Note that from P16 to adult the intensity of both PSMA8 and PSMA7 bands impedes its independent observation. (B) Double immunolabeling of spermatocyte spread preparations with PSMA8 (green) and SYCP3 (red) by Stimulated emission depletion (STED) microscopy, showing that PSMA8 localizes to the central region of the SC. PAR (pseudo-autosomal region) of the XY bivalent is indicated with an arrow. (C) Immunolocalization of PSMA8 in mouse testis after *in vivo* electroporation of a plasmid encoding a protein fusion of PSMA8 with GFP (GFP-PSMA8). PSMA8 was detected with anti-GFP antibody (green) and endogenous SYCP3 was detected using mouse anti-SYCP3 (red). (D) Triple labeling of PSMA8 (green), SYCP3 (blue) and SYCP1 (red) in *Rec8*^{-/-} and *Six6os1*^{-/-}. PSMA8 is detected in the pseudosynapsed AEs of the meiotic *Rec8* cohesin mutant but is absent from the unsynapsed AEs in *Six6os1*^{-/-} spermatocytes. Bar in panels, 5 μ m (B, upper panel), 1 μ m (B, lower panel) and 10 μ m (C, D).

<https://doi.org/10.1371/journal.pgen.1008316.g001>

testes showed no PSMA8 protein expression by western blotting when analyzed using two independent polyclonal antibodies (S3C Fig). Immunofluorescence analysis of PSMA8 expression (R2 antibody, S3D Fig) revealed a weaker signal in the SC of the mutant spermatocytes than in WT spermatocytes (51% less; 4.22 ± 1.9 WT vs 2.05 ± 1.7 KO), likely representing PSMA7 detected by the R2 antibody (also observed in the western blot; S3C Fig). These results indicate that the generated mutation is a null allele of the *Psm8* gene (herein termed *Psm8*^{-/-}).

Mice lacking PSMA8 did not display any somatic abnormalities; however, male but not female mice were sterile (S1 Table). Indeed, *Psm8* mutation resulted in a reduction of the testis weight (63.09% decrease; N = 6) and the absence of spermatozoa in the epididymides (Fig 2A and 2B). Histological analysis of adult *Psm8*^{-/-} testes revealed the presence of apparently normal numbers of spermatogonia, spermatocytes, Sertoli cells and Leydig cells (Fig 2B).

Fig 2. PSMA8 deficiency leads to azoospermia. (A-B) Genetic ablation of *Psm8* leads to a reduction of the testis size (A) ($n = 6$, WT and KO; Welch's t -test analysis: $p < 0.0001$), and (B) the accumulation of metaphase I (black asterisks), apoptotic meiotic division (red asterisks), round spermatids entering apoptosis (arrowheads), and apoptotic round spermatids (blue asterisks) in PAS stained testis sections. The spermatogenic arrest leads to empty epididymides and azoospermia. Bar in upper panels 100 μ m, lower panels 200 μ m and in right panels, 5 μ m. (St) Seminiferous tubules and (Ep) Epididymides. (C) Immunofluorescence analysis of p-ser10-H3 (green) of paraffin sections of *Psm8*^{+/+} and *Psm8*^{-/-} tubules. Nuclei were counterstained with DAPI. Bar represents 10 μ m. The diagram represents the quantification of the fraction of tubules showing the indicated number of metaphase I/II. Number of tubules counted for each genotype is expressed in [S2B Table](#). (D) Low magnification view of a

representative squash preparation of seminiferous tubules showing the accumulation of metaphases I and metaphases II in knock-out *Psm8* in comparison with a representative wild-type view. The identity of metaphases I /metaphases II (asterisks) was confirmed by the immunolabeling of SYCP3 (red) in squash preparations. Chromosomes were counterstained with DAPI (blue). The diagram represents the percentage of spermatocytes at metaphase I and II (normal and apoptotic) in relation with the total number of spermatocytes from *Psm8*^{+/+} and *Psm8*^{-/-} tubules (right). Quantification and number of cells analyzed are described in S2C Table. Welch's *t*-test analysis: * *p*<0.01; ** *p*<0.001; *** *p*<0.0001.

<https://doi.org/10.1371/journal.pgen.1008316.g002>

Mouse seminiferous tubules can be classified from epithelial stage I to XII by determining the groups of associated germ cell types in histological sections. Following these criteria, we found that spermatogenesis in the mutant testes proceeded normally up to diplotene in epithelial stage XI. However, the proportion of tubules at stage XII was more than 2-fold increased in the mutant sections (12.5% in mutants versus 5.4% in WT, S2A Table). Given that spermatocytes in meiotic divisions were seen to occur at epithelial stage XII, we used p-ser10-H3 (pH3) staining to analyze the number of metaphase I and II cells present in these tubules, finding an increase in the mutant (Fig 2C and S2B Table). Quantitative analysis of seminiferous tubules in squashed preparations confirmed the increase in the number of metaphase I and metaphase II cells as compared with WT testes (77% and 89% respectively, Fig 2D and S2C Table). Moreover, a large proportion of these metaphases were positive for Caspase-3 and TUNEL indicating apoptosis (Figs 2D, 3A and 3B and S2C Table).

Together with the accumulation of apoptotic meiotic divisions, other apoptotic cells could be also observed that, from their size and molecular markers of the acrosome and chromatoid body, were round spermatids (Fig 3C and S4 Fig). Indeed, seminiferous tubules in PSMA8-deficient testes sometimes contained a few surviving round spermatids. However, these round spermatids were unable to form a proper acrosome but did accumulate some PAS positive material. Apoptotic round spermatids were also seen and no elongating spermatids were observed (Fig 2B). We corroborated that round spermatids were arrested at early stages by immunolabeling for H2AL2. H2AL2 is a transition histone essential for the first replacement of histones by TNP1 and TNP2 before protamine incorporation [26]. H2AL2 was absent from mutant spermatids (S5A Fig). We also used FACs analysis of whole cells from seminiferous tubules to verify this analysis. The results obtained confirmed the presence of a small haploid compartment in *Psm8*^{-/-} testes (Fig 3D and S5B Fig). We conclude from these results that PSMA8 deficiency causes the accumulation of spermatocytes in metaphase I and II which either enter massively into apoptosis or give rise to a low number of aberrant round spermatids that finally apoptose long before histone replacement takes place.

***Psm8*-deficient spermatocytes show normal synapsis/desynapsis and DSB repair but have abnormal metaphases I and II**

Metaphase I accumulation can occur either because of a failure to enter anaphase or because of some event taking place during prophase (SC formation, DSBs repair or chromosome recombination) that aberrantly triggers a checkpoint-mediated delay.

To test this, we first analyzed the assembly/disassembly of the SC by monitoring the distribution of SYCP1, as co-labeling of SYCP3 and SYCP1 highlights regions of synapsis in spermatocytes. We did not observe any differences in this process from zygonema to diakinesis (S6 Fig).

We next studied the kinetics of DSB repair during meiosis. Meiotic DSBs are generated by the nuclease SPO11 and are then resected to form ssDNA ends that invade into the homologous chromosome. DSBs are marked by the presence of phosphorylated H2AX (γ -H2AX) [27]. The distribution of γ -H2AX in mutant spermatocytes was similar to that found in WT cells at prophase I (S7A Fig and S3 Table). We also did not observe any differences in the

Fig 3. Apoptosis, FACs and aberrant metaphase II and spermatid cells in *Psm8*-deficient mice. (A) Double immunolabeling of Caspase3 (green) and (B) TUNEL (green) with SYCP3 (red). Non-apoptotic metaphase I cells from *Psm8*^{+/+} show absence of green staining whereas apoptotic metaphases I from *Psm8*^{-/-} show intense Caspase-3 and TUNEL labeling. Chromatin was counterstained with DAPI. (C) Acrosome positive labeling of round spermatids by PNA staining (green). (D) FACs analysis of cells from whole seminiferous tubules from wild type and *Psm8*^{-/-} showing in both genotypes (N = 2) the presence of 4C, 2C and 1C compartment as a result of the early spermatid arrest. Source data

describing the gating strategy is shown in [S5B Fig](#). (E) Double immunolabeling of metaphase I cells with tubulin (green) and ACA (red) showing normal (*Psm8^{+/+}*) and abnormal spindles (*Psm8^{-/-}*). (F) Double immunolabeling of SYCP3 (green) with ACA (red) in wild-type and *Psm8^{-/-}* spermatocytes at metaphase II which shows aberrant accumulation of SYCP3 at the centromeres. (G) Double immunolabeling of PA200 (green) and SYCP3 (red) in chromosome spreads. PA200 is detected at the chromosome axes of the autosomal and XY bivalents during pachytene in wild type spermatocytes in contrast to the absence of labeling in *Psm8^{-/-}* spermatocytes. Bar in panels (C, E) 5 μ m and 10 μ m (A, B, F and G).

<https://doi.org/10.1371/journal.pgen.1008316.g003>

distribution of RAD51, a recombinase that promotes homologous strand invasion [28], ([S7B Fig](#) and [S3 Table](#)). Because defective DNA repair ultimately abrogates CO formation [29] and because of the involvement of ubiquitylation / sumoylation in CO designation [30], we analyzed the distribution of MLH1 foci [31], a mismatch repair protein (marker of crossover sites) that functions in the resolution of joint molecules at the end of crossover formation [32]. We found a similar value between the KO (24.9 ± 0.9 foci) and the WT (24.3 ± 1.1 foci; [S7C Fig](#) and [S3 Table](#)). These results indicate that the repair of meiotic DSBs and synapsis/desynapsis proceed normally during prophase I in the absence of PSMA8, and is not responsible for the observed metaphase I accumulation.

We also analyzed the morphology of the metaphase I / II cells by staining for tubulin (spindle) and SYCP3. The results showed an aberrant morphology, the presence of multipolar spindles ([Fig 3E](#)), and also a striking aberrant labeling of SYCP3 at the centromeres of metaphase II chromosomes (SYCP3 labeling is barely visible in metaphase II sister kinetochores in WT cells, [Fig 3F](#)). Finally, the arrested round spermatids showed the presence of multiple patches of heterochromatin after DAPI staining ([Fig 3C](#) and [S4 Fig](#), chromocenter fragmentation), suggesting abnormal chromosome segregation or cytokinesis.

PSMA8 deficiency abolishes H4ac turnover from late prophase to round spermatids

During spermiogenesis, most of the histones are replaced by basic transition proteins, and ultimately by protamines, facilitating chromatin compaction. Hyperacetylation of core histones during this process, and especially the acetylation of H4K16, is assumed to play a pivotal role in the initiation of histone displacement and chromatin ultracondensation [33, 34]. The proteasome activator subunit PA200 targets acetylated histones for degradation during histone replacement [5].

The core subunit PSMA8 co-immunoprecipitated PA200 ([S4 Table](#)). Given the stoichiometric relationship between the CP and RP, we analyzed the expression of PA200 by immunofluorescence in the absence of PSMA8. Whilst PA200 decorated the AEs of WT spermatocytes, we failed to observe any signal in the AEs of mutants ([Fig 3G](#) and [S8 Fig](#)). In addition, we were not able to detect PA200 by mass spectrometry analysis of PSMA7/8 immunoprecipitation of *Psm8*-deficient testis extracts (see section Purification of PSMA8-interacting proteins, [S4 Table](#)). These results indicate that PSMA8 is necessary or promotes the assembly of PA200 to the CP. Thus, within the limits of detection, the deficiency of *Psm8* leads to a drastic decrease of PA200.

To understand the acetylated-dependent degradation of histones by the proteasome [5], we measured the acetylation status of three core histones, H2AK5ac, H3ac and H4ac (pan-H4ac and H4K16ac) in chromosome spreads by double immunolabeling for SYCP3 and the corresponding acetylated histone ([Fig 4A–4D](#) and [S9–S12 Figs](#)). This procedure enables a more precise staging of the spermatocytes and is a more efficient mean to quantitate signals than peroxidase immunostaining of testis sections [5]. The loss of PSMA8 led to the accumulation of H2AK5ac, H3ac, H4ac and H4K16ac, albeit to different degrees. Results showed that the levels of H2AK5ac, H3ac, H4ac and H4K16ac were moderately higher in *Psm8^{-/-}* cells, with a

Fig 4. Histone acetylation, nuclei ubiquitylation and proteasome activity in PSMA8-deficient mice. (A-D) Plots represent the quantification of the fluorescence intensity from $Psm8^{+/+}$ and $Psm8^{-/-}$ spermatocytes at early pachytene (EP), mid pachytene (MP), late pachytene (LP), early diplotene (ED), late diplotene (LD), diakinesis (DK), metaphase I (MI) and round spermatid (RS) corresponding to the immunolabeling of (A) H2AK5ac, (B) H3ac, (C) H4ac, and (D) H4K16ac. Representative figures for each immunofluorescence are presented in S9–S12 Figs. (E) Proteasome activity of $Psm8$ -deficient testis. 100 μ g of protein from whole testis extracts of $Psm8^{+/+}$ and $Psm8^{-/-}$ mice were inoculated into 96-well plate and the proteasome peptidases activities were measured. The enzymatic activities relative to WT are shown. (F) Plots represent the quantification of the

fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spread (upper) and squashed (lower) spermatocytes. Welch's *t*-test analysis: * *p* < 0.01; ** *p* < 0.001; *** *p* < 0.0001.

<https://doi.org/10.1371/journal.pgen.1008316.g004>

relative increase at late prophase I (Fig 4A–4D and S9–S12 Figs). We failed to detect staining for H2AK5ac and H3ac in spermatocytes in late diakinesis and round/arrested spermatids. In contrast, pan-H4ac and H4K16ac also labeled metaphase I chromosomes, interkinesis nuclei and round/arrested spermatids, with greater intensity in mutant than in WT cells (Fig 4C and 4D and S11 and S12 Figs). The accumulation of acetylated histones during prophase I and particularly of H4ac and H4K16ac in the arrested round spermatids suggests that the PSMA8--containing proteasomes are involved in the acetylation-dependent degradation of histones.

Proteasomal activity in *Psm8*-deficient mice

We next investigated the biochemical activity of testis extracts lacking PSMA8-containing proteasomes by measuring chymotrypsin-like activity (corresponding to the catalytic subunit β 1), caspase-like activity (corresponding to β 5) and trypsin-like activity (β 3) by a standard fluorogenic assay [35] in the presence and absence of SDS (activated proteasome). Results showed that proteasomal activity in *Psm8*-deficient testis extracts was not noticeably different from that in WT extracts. Indeed, the trypsin-like activity was the only proteolytic function with a modest reduction in the KO (Fig 4E). Overall, these results show that the general proteasome activity of the *Psm8*-deficient testis is not radically changed, which is likely due to the presence of PSMA7-dependent CPs (see dataset 1 in [36]).

To ascertain the degree of activity *in vivo*, we first investigated the steady-state levels of protein ubiquitylation in testis during mouse meiosis. Using immunofluorescence, we analyzed spermatocytes obtained from spreads and squashed preparations with ubiquitin antibodies (Fig 4F and S13 Fig). The results showed a slight decrease of chromatin bound ubiquitylated proteins but an increase in the soluble fraction of ubiquitylated proteins during prophase I (Fig 4F and S13 Fig). These results are partially in agreement with the observed increase in the ubiquitylation state of cultured spermatocytes treated with the proteasome inhibitor MG132 (18), and suggest a specific function of the PSMA8-containing proteasomes in the controlled degradation of ubiquitylated proteins during spermatogenesis.

Purification of PSMA8-interacting proteins

The composition of the CP and its RPs has previously been established by mass-spectrometric analysis of crude preparation of proteasomes from whole testes [37]. To better understand the molecular mechanism underlying the mutant phenotype, we purified PSMA7/8-interacting proteins by single-step affinity chromatography (see [Material and methods](#) for a detailed description). Most of the canonical subunits of the CP and RP were present within the more than 596 proteins of the PSMA8 proteome (S5 Table, using a conservative cut-off, see [methods](#)). In agreement with previous results, among the two activators of the testis-specific proteasome detected (PA200 and Pa28 γ) [5], PA200 was the most abundant. In contrast to previous observations, we were unable to detect Pa28 α and Pa28 β or the inducible catalytic subunits of the immunoproteasome (β 1i, β 2i and β 5i) [5], suggesting a very low abundance or absence. We could not detect PA200 as an interacting protein of PSMA7/8 in testis extracts from *Psm8*-deficient testes (S4 Table).

Among the novel proteasome-interacting proteins (PIPs) detected were chaperones including CCT6b and CCT2, ubiquitin ligases (TRIP12, NEDD4, TRIM36 and RAD18), and novel ubiquitin specific proteases (USPs) such as USP9X, USP34, USP5 and USP47 (S6 Table). We

studied the proteins enriched in the immunoprecipitation through functional (gene ontology, GO) and pathway analysis (KEGG). The top GO and KEGG results were related to the proteasome and to ribonucleoproteins. Pathway analysis showed links to spermatogenesis, cell cycle, and meiosis (see [S1 Text](#)), in accordance with the observed mutant phenotype.

Interestingly, we identified meiotic proteins *a priori* unrelated to the UPS such as DAZL (deleted in azoospermia), SPAG1 (Sperm-associated antigen 1), SPATA5/20 (Spermatogenesis-associated protein 5/20), the tudor domain proteins TDRD1/6/9, MAEL (repressor of transposable elements), and RNF17. These PIPs could represent proteins captured during ubiquitin-dependent targeted degradation [38] and/or proteins interacting *via* ubiquitin-independent proteasomal degradation, as has been shown for the related subunit $\alpha 4$ /PSMA7 [39]. Altogether, the list of novel PIPs included novel potential readers, erasers and writers of the ubiquitin code [40] of the testis-specific proteasome, reflecting its complexity. Among these PIPs, we focused our attention on the following candidates for their role in chromosome segregation and synapsis: SYCP1, TRIP13, TEX30, PIWIL1, PIWIL2 and CDK1 ([S6 Table](#)).

Among the possible interactors, we first evaluated the transverse filament protein SYCP1. Because *Sycp1* mutant mice are infertile but otherwise healthy [41], we analyzed the interaction of SYCP1 with PSMA8 and its localization in mutant meiosis. We co-transfected *Sycp1* with *Psm8* in HEK293T cells and we detected co-immunoprecipitation between SYCP1 and PSMA8 ([Fig 5A](#)). Despite the observation that SYCP1 is properly loaded to the SC and removed from desynapsed regions ([S6 Fig](#)), we observed an abnormal accumulation of SYCP1 in *Psm8*-deficient metaphase I cells, ([Fig 5B](#)). These results suggest defective degradation of SYCP1 with very likely detrimental functional consequences in the exit of meiosis.

We next extended the validation analysis of the remaining candidate interactors by co-immunoprecipitation with PSMA8, making use of the same heterologous system of HEK293T cells. These included TEX30, PIWIL1, PIWIL2, CDK1 and TRIP13. All protein-protein interaction assays carried out were negative ([S14A Fig](#)) with the exceptions of the cyclin dependent kinase CDK1 and the AAA-ATPase TRIP13 (AAA-ATPases associated with diverse cellular activities; see [Figs 6A](#) and [7A](#)). Because of the relevance of CDK1 in metaphase transition, we first determined the expression levels of CDK1 by immunofluorescence. The results showed that more CDK1 but not the related kinase CDK2 [42] could be detected in the centromeres of metaphase I chromosome from mutant cells ([Fig 6B](#) and [S15A Fig](#); KO 0.31 ± 0.2 vs 0.19 ± 0.1 WT; an increase of $\sim 40\%$). To determine whether the increased level of CDK1 corresponded to its active or inactive phosphorylated form, we used an antibody against CDK1-Tyr15-p (inactive form, [Fig 6C](#)). The results showed no differences in the labeling at the centromeres of the metaphase I chromosomes, and therefore a decrease in phospho-CDK1/total CDK1 ratio in mutant cells. Given that CDK1 must be complexed with cyclin B1 to be active, we reasoned that if higher levels of active CDK1 are present, cyclin B1 would be similarly increased. Results showed an increase of cyclin B1 at the centromeres of metaphase I chromosomes ([Fig 6D](#)). This result was congruent with the increased amount of CDK1 and CyclinB1 observed by western blot and in squashed seminiferous tubules ([Fig 6E](#) and [S15B](#) and [S15C Fig](#)). Overall, these findings suggest that loss of PSMA8 causes an increase of CDK1 / CyclinB1 which would cooperate in the accumulation of metaphase I / metaphase II that ultimately results in apoptotic metaphase plates.

We also analyzed the distribution of TRIP13, a pleiotropic ATPase that participates in meiotic DNA repair and chromosome synapsis through HORMAD interaction and somatic spindle assembly checkpoint (SAC) proficiency through MAD2 interaction [43–46]. We first performed immunofluorescence analysis of TRIP13 in *Psm8*-deficient and WT spermatocytes. Results using two independent antibodies showed robust labeling of the telomeres from zygonema (two dots) to pachynema (fused to a single dot) in WT cells, which declined from

Fig 5. SYCP1 interacts with PSMA8 and is accumulated in *Psma8*-deficient metaphase I cells. (A) HEK293T cells were transfected with Flag-PSMA8 and GFP-SYCP1. Protein complexes were immunoprecipitated overnight with either an anti-Flag or anti-EGFP or IgGs (negative control), and were analyzed by immunoblotting with the indicated antibody. PSMA8 co-immunoprecipitates with SYCP1. (B) Double immunolabeling of squashed tubules with SYCP1 (green) and SYCP3 (red) in wild-type and *Psma8*^{-/-} spermatocytes at metaphase I. Chromatin was stained with DAPI (blue). Bar in panel, 10 μ m.

<https://doi.org/10.1371/journal.pgen.1008316.g005>

diplo-nema to diakinesis. The staining pattern was similar but enhanced in mutant spermatocytes (Fig 7B). However, the staining pattern of TRIP13 at metaphase I differed between WT and mutant cells. Specifically, it was detected at the kinetochores of *Psma8*^{-/-} spermatocytes but was absent in WT cells (Fig 7B). This labeling pattern at the metaphase I kinetochores resembles TRIP13 staining in somatic cells [47]. These results thus suggest that TRIP13 accumulates in the absence of a functional PSMA8-containing proteasome.

We next analyzed several downstream effectors of TRIP13, HORMAD1, HORMAD2, and the mitotic checkpoint protein MAD2 [48–50]. No differences were observed in the HORMAD1/2 labeling pattern between WT and mutant cells (S16 Fig). It has been shown in *C. elegans* that in the absence of TRIP13, MAD2 recruitment to kinetochores is delayed and that in addition to its role in checkpoint silencing, TRIP13 also contributes to spindle checkpoint activation [50]. It could thus be argued that an excess of TRIP13 would increase MAD2 loading to kinetochores thereby delaying mitotic exit. We confirmed this prediction and found that MAD2 expression at the kinetochores was enhanced in *Psma8*^{-/-} spermatocytes (Fig 7C), further validating a functional consequence of TRIP13 accumulation at the kinetochores.

In order to validate the substrate specificity of the PSMA8-containing proteasome in protein degradation, we analyzed the expression levels of the separase inhibitor securin (PTTG1),

Fig 6. PSMA8 deficiency causes an accumulation of CDK1 and Cyclin B1 in spermatocytes. (A) HEK293T cells were transfected with Flag-PSMA8 and GFP-CDK1. Protein complexes were immunoprecipitated with either an anti-Flag or anti-EGFP or IgGs (negative control) and were analyzed by immunoblotting with the indicated antibody. PSMA8 co-immunoprecipitates with CDK1 (as well as reciprocally). (B) Double labeling of endogenous CDK1 (green) and SYCP3 (red) in mouse spermatocytes at metaphase I. Chromatin was stained with DAPI (blue). During metaphase I, CDK1 labels in a slight and disperse way the chromosomes and in a more intensely fashion the centromeres of bivalents. This labeling pattern is enhanced in a normal *Psm8*-deficient metaphase I. Plot under the panel represents the quantification of the fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} metaphase I cells. (C) Double labeling of endogenous CDK1-Tyr15phosphorylated (green) and SYCP3 (red) in mouse spermatocytes at

metaphase I showing similar expression levels in *Psm8*^{+/+} and *Psm8*^{-/-}. Chromatin was stained with DAPI (blue). (D) Double labeling of endogenous cyclin B1 (green) and SYCP3 (red) in mouse spermatocytes at metaphase I showing higher expression levels in *Psm8*^{-/-}. Plot under the panel represents the quantification of the fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} metaphase I cells. Welch's *t*-test analysis: * *p*<0.01; ** *p*<0.001; *** *p*<0.0001. (E) CDK1 and CyclinB1 were measured by western blot analysis of protein extracts from whole testis of *Psm8*^{+/+} (WT) and *Psm8*^{-/-} (KO) (*n* = 2 mice). Bar in panels, 10 μm. Welch's *t*-test analysis: * *p*<0.05; ** *p*<0.001; *** *p*<0.0001.

<https://doi.org/10.1371/journal.pgen.1008316.g006>

a well-known substrate of the somatic proteasome. Immunofluorescence analysis showed similar levels of PTTG1 in *Psm8*^{-/-} and WT spermatocytes (S17 Fig). This result suggests that PSMA8-containing proteasomes are not involved in the degradation of classical ubiquitylated substrates degraded by the somatic proteasome.

PSMA8 interacts with proteins of the synaptonemal complex

To investigate the molecular basis of PSMA8 localization in the SC, and considering the alteration of SYCP3 and SYCP1 in *Psm8*^{-/-} spermatocytes (Fig 3F and Fig 5B), we used a candidate gene approach to identify additional putative interactors of PSMA8. We co-transfected *Psm8* with cDNAs encoding each of the known central element proteins (SIX6OS1, SYCE1, SYCE2, SYCE3, and TEX12), and the AE protein SYCP3. As positive controls, we exploited the well-known interaction between SYCE2 and TEX12 [51] (S14C Fig). Surprisingly, we detected specific co-immunoprecipitation of PSMA8 with SIX6OS1 and SYCE3 (Fig 8A and S14B Fig). We were unable to immunoprecipitate transfected SYCP3 (using several tags or antibodies against SYCP3), likely due to the highly complex structures of transfected SYCP3, which prevented to perform co-immunoprecipitation experiments. Because SYCP3 forms filamentous structures in the cytoplasm of transfected cells, termed polycomplexes [52], co-expression of an interacting protein with SYCP3 may lead to its recruitment to polycomplexes [24], an indication of protein interaction. Indeed, we obtained self assembled higher structures when *Psm8* was co-transfected with *Sycp3* (Fig 8B). This SYCP3-dependent cytological interaction was not observed when *Psm8* was co-transfected (Fig 8B), further validating the specificity of the interaction given the extensive protein similarity between both PSMA8 and PSMA7 (92%). To validate this interaction *in vivo*, we performed a detailed analysis of SYCP3 in mouse mutant squashed spermatocytes, a procedure in which no solubilization or protein extraction is performed. We observed SYCP3 aggregates/polycomplexes in the *Psm8*-deficient spermatocytes during prophase I and metaphase I / II (Fig 8C and 8D and S7 Table). SYCP3 accumulated in metaphase II chromosomes as abnormal SYCP3 labeling at the centromeres between sister kinetochores and as aggregates in the cytosol (Fig 3F and Fig 8D). Global accumulation of SYCP3 was also observed by western blot of whole testis under high denaturing conditions (Fig 8E) [53]. Interestingly, it has been previously shown that cultured spermatocytes chemically treated with the proteasome inhibitor MG132 form SYCP3 aggregates [17]. Overall, our results suggest that SYCP3 is targeted for degradation by the PSMA8-containing proteasome and that in the absence of PSMA8 its accumulation could mediate, at least in part, the arrest and apoptosis of spermatocytes.

Discussion

The testis-specific proteasome is one of the three tissue-specific proteasomes identified in mammals (together with the immunoproteasome and the thymoproteasome); however, little is known about its biochemical and physiological function. The groundbreaking work of Xiao-Bo Qiu and colleagues showing the acetyl-histone preference of the PA200 subunit of the proteasome [5] has provided novel insights into the proteasome-dependent degradation of non-ubiquitylated proteins and led to the designation of spermatoproteasome to the

Fig 7. TRIP13 and MAD2 levels are increased in *Pisma8*-deficient spermatocytes. (A) HEK293T cells were transfected with a plasmid encoding GFP-TRIP13 and Flag-PSMA8. Protein complexes were immunoprecipitated with either an anti-Flag or anti-EGFP or IgGs (negative control), and

immunoblotted with the indicated antibody. (B) Double immunolabeling of TRIP13 (green) and SYCP3 (red). TRIP13 labels the telomeres at pachytene and the intensity of the labeling decreases through desynapsis at diplotene and diakinesis. This labeling is enhanced during prophase I in the *Psm8* mutants but its main pattern is not altered. At metaphase I, a faint labeling of sister kinetochores is observed in the *Psm8*^{-/-} spermatocytes that is absent in the wild type. Plot over the panel represents the quantification of the fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes at pachytene and late diplotene. (C) MAD2 (green) labels with enhanced intensity the centromeres of the chromosomes from *Psm8*^{-/-} metaphase I cells in comparison with the WT controls. Plot right to the panel represents the quantification of the fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes at metaphase I spermatocytes. Bar in panels, 10 μ m. Welch's *t*-test analysis: * $p < 0.01$; ** $p < 0.001$; *** $p < 0.0001$.

<https://doi.org/10.1371/journal.pgen.1008316.g007>

PA200-containing proteasome. However, following the criteria employed for the designation of the thymoproteasome, which were devised based on the restricted expression of its $\beta 5t$ subunit in the thymus [9] (GTEx portal), we suggest that the term spermatoproteasome be restricted exclusively to the PSMA8-containing proteasome instead of the widely expressed PA200 subunit [5].

We have shown that genetic depletion of *Psm8* causes the delocalization and the drastic decrease (loss of detection) of the proteasome activator PA200 in spermatocytes. Accordingly, *Psm8*-deficient spermatocytes accumulate acetylated histones. PSMA8 deficiency is comparatively more severe than that of the PA200 single mutant (subfertile) and of the PA200 and PA28 γ double mutant, which do not show an arrest in spermatogenesis despite being infertile *in vivo* but not *in vitro* (spermatozoa are not motile but can fertilize *in vitro* [54]). From a genetic analysis perspective, this result would suggest that PSMA8 has additional functions that are independent of the activators PA200 and PA28 γ . Our proteomic analysis, together with other data [10], supports this notion and indicates that PSMA8-containing proteasomes can be associated with other regulators such as the 19S subunit, expanding its targets.

Beyond its role in initiation of histone replacement [34], H4K16ac is involved in the three waves of H2AX phosphorylation during prophase I [55]. We have shown that *Psm8* deficiency causes the accumulation of H4ac and H4K16ac during prophase I. However, we did not observe defects in this process in the form of a different staining pattern for γ -H2AX (leptonema and zygonema), including the expansion of γ -H2AX staining to the chromatin of the sex body (in pachynema). However, the observed premature accumulation of H4K16ac at early round spermatid might cause a defect in histone removal later on in spermiogenesis if the *Psm8*^{-/-} mutants spermatids would not have entered apoptosis before this event.

We have shown that spermatoproteasome deficiency causes severe defects in protein turnover of key meiotic players that affect metaphase I/II exit, but not the complex process of meiotic recombination that occurs during prophase I (CO). By using a candidate approach of PIPs, we have identified CDK1 and TRIP13 as likely crucial proteins that have an abnormal expression pattern during meiotic metaphase in mutant mice. Given the key roles of these proteins in all aspects of mitotic/meiotic division (including SAC activation), the accumulation of aberrant metaphase I/II spermatocytes in *Psm8*-deficient mice is to be expected.

The role of CDK1 in the metaphase-anaphase transition is complex and is multifaceted. CDK1 inhibits and activates APC/C by promoting the SAC and also by a SAC-independent mechanism [56]. The balance between these opposing functions determines cyclin B1 destruction and separase activation, giving rise to cohesin cleavage and anaphase onset [57]. Based on the normal expression levels of PTTG1 in *Psm8*^{-/-} metaphase I cells, it can be argued that there is no precocious APC activation in *Psm8*-deficient cells (S17 Fig). Given that CDK1 activation of the SAC is dominant over the activation of APC^{Cdc20} [58] in oocytes, we suggest that the former effect is acting on *Psm8*-deficient spermatocytes. The question how CDK1 promotes the SAC is still unresolved in oocytes and even less is known about this in spermatocytes.

Fig 8. PSMA8 interacts with proteins of the SC. (A) PSMA8 co-immunoprecipitates with SIX6OS1 and SYCE3. HEK293T cells were transfected with plasmids encoding Flag-PSMA8 and GFP-SIX6OS1 or GFP-SYCE3. Protein complexes were immunoprecipitated overnight with either an anti-Flag or anti-EGFP or IgGs (negative control), and were analyzed by immunoblotting with the indicated antibody. (B) Double immunofluorescence of transfected HEK293T cells with plasmids encoding Flag-PSMA8 and Flag-PSMA7 alone or together with plasmid encoding SYCP3-HA and immuno-detected with antibodies against Flag (green) or HA (red). Transfected PSMA8 alone is delocalized and occupies the whole cell whereas when co-transfected with SYCP3-HA is recruited to form polycomplexes. PSMA7 do not form polycomplexes when co-transfected with SYCP3-HA. (C-D) SYCP3 is accumulated *in vivo* in *Pisma8^{-/-}* spermatocytes. (C) Double immunolabeling of squashed tubules with SYCP3 (red) and SYCP1 (green) in wild-type and *Pisma8^{-/-}* spermatocytes at prophase I showing large SYCP3 aggregates surrounding the nuclei (arrows). (D) Double immunolabeling of squashed tubules with SYCP3 (green) and ACA (red) in wild-type and *Pisma8^{-/-}* spermatocytes at metaphase I and II. *Pisma8^{-/-}* metaphases I show labeling of SYCP3 in aggregates (arrows, absent in the WT) in addition to its typical labeling at the centromeres. Metaphases II from *Pisma8^{-/-}* show labeling for SYCP3 at the centromeres between the sister kinetochores and as aggregates in the cytosol (arrows) whereas wild type metaphases II show barely visible SYCP3 labeling. (E) SYCP3 was measured by western blot analysis of protein extracts from whole testis of *Pisma8^{+/+}* (WT) and *Pisma8^{-/-}* (KO) (n = 2 mice). Bar in panels, 10 μm. Welch's *t*-test analysis: * *p* < 0.05; ** *p* < 0.001; *** *p* < 0.0001.

<https://doi.org/10.1371/journal.pgen.1008316.g008>

Another group of proteins found to be deregulated in spermatoproteasome-deficient mice are the SC structural proteins SYCP1 and SYCP3. The precise effect of the accumulated SYCP1 in the cytoplasm of *Pisma8^{-/-}* spermatocytes cannot be experimentally analyzed. However, the coiled-coil structure and self-assembly abilities of SYCP1 strongly suggest a functionally detrimental consequence. Similarly, the presence of SYCP3 aggregates during pachynema and metaphase I mutant spermatocytes and its persistence at metaphase II

centromeres, where SYCP3 is barely visible in WT cells, also suggest a detrimental effect on these cells causing their entrance into apoptosis.

We have also shown that PSMA8 is delocalized in the severe synapsis *Six6os1* mutant, which is consistent with the observed co-immunoprecipitation of PSMA8 with SYCP1, SIX6OS1 and SYCE3. All the synapsis-less mutants of CE proteins failed to load properly or lacked SYCP1 and the remaining CE proteins [24, 59–61]. Thus, we would predict delocalization of the spermatoproteasome from the SC in the remaining mouse mutants of the CE proteins. Overall, our results support the idea of a physical anchorage or recruitment of the spermatoproteasome to the SC especially through SYCP3, possibly facilitated or mediated by SYCP1, SIX6OS1 and SYCE3 as their most relevant structural partners. Supporting this notion, the Zip1 transverse filament protein of the yeast SC participates in the recruitment of the proteasome to the SC [22], suggesting an evolutionary conservation of the mechanism.

Yeast mutated for a nonessential subunit of the proteasome (pre9) showed abnormal meiotic recombination, pairing and synapsis [22]. Similar but milder defects were also observed in spermatocytes cultured with a proteasome inhibitor [17]. It has been proposed that the UPS regulates the proteostatic turnover of the ZMM which is required for efficient synapsis and CO [17], through the RNF212 (E3 sumo ligase)-Hei10 (E3 ubiquitin ligase) pathway [31]. Given this, the lack of a meiotic recombination phenotype (DSBs are generated and repaired and COs are generated normally) in our *Psma8*-deficient mouse is surprising. It can be argued that PSMA7-containing proteasomes are still present and at the early stages of meiosis are compensating for the loss of function of *Psma8*. Another possible but not mutually exclusive explanation is that the main targets of the PSMA8-containing proteasome are proteins from mid-prophase I onwards.

The spermatoproteasome through its complex interactome would serve as a hub for the fine tuning of several fundamental key molecules of the spermatogenic process such as those analyzed during the present work (SYCP1, SYCP3, TRIP13, CDK1 and acetyl-histones). Our data suggest that deregulation of proteostasis of key meiotic proteins promoting cell division leads to the presence of multipolar spindles and aberrant meiotic exit. Thus, we favor an explanation in which the joint contribution of several pathways is responsible for the observed infertility.

In relation to human disease, protein degradation was one of the top cellular functions found in an unbiased differential proteomic profiling of spermatozoa proteins from infertile men with a varicocele [62]. More specifically, PSMA8 is among the top 7 in this list of proteins that are differentially expressed, suggesting a causal role in the severity of the disease. From an organismal perspective, *Psma8* transcription is mainly restricted to the human testis and to some tumors like Burkitt lymphoma and melanoma (TCGC database). Altogether, and considering the PSMA8 dependency of the mouse male germline, we suggest that the spermatoproteasome may be an effective target for male contraception and for the treatment of some human malignancies.

Material and methods

In vivo electroporation of testes

Testes were freed from the abdominal cavity and 10 μ l of DNA solution (50 μ g) mixed with 1 μ l of 10 $\times$ FastGreen (Sigma Aldrich F7258) was injected into the rete testis with a DNA embryo microinjection tip. After a period of 1 h following the injection, testes were held between electrodes and four electric pulses were applied (35 V for 50 ms each pulse) using a CUY21 BEX electroporator.

Production of CRISPR/Cas9-Edited mice

Psmas8-sgRNAs G71 5'-GGGCATACT CCACTTGGAAA -3' G84 5'-ACCGCGGTAAGCTG CTCCCC-3' targeting exon 1 and intron 1 were predicted at crispr.mit.edu. *Psmas8*-sgRNAs were produced by cloning annealed complementary oligos at the BbsI site of pX330 (#42230, Addgene), generating PCR products containing a T7 promoter sequence that were purified (NZYtech), and then *in vitro* transcribed with the MEGAshortscrip T7 Transcription Kit (Life Technologies). The plasmid pST1374-NLS-flag-linker-Cas9 (#44758; Addgene) was used for generating Cas9 mRNA. After linearization with AgeI, it was transcribed and capped with the mMESSAGE mMACHINE T7 Transcription Kit (AM1345; Life Technologies). RNAs were purified using the RNeasy Mini Kit (Qiagen). RNAs (100 ng/μl Cas9 and 50ng/μl each guide RNA) were microinjected into B6/CBA F2 zygotes (hybrids between strains C57BL/6J and CBA/J) [63] at the Transgenic Facility of the University of Salamanca. Edited founders were identified by PCR amplification (Taq polymerase, NZYtech) with primers flanking exons 1 and intron 1 (Primer F 5'-CTTCTCGGTATGACAGGGCAATC-3' and R 5'-ACTCTACCTC CACTGCCAAC CTG-3') and either direct sequenced or subcloned into pBlueScript (Stratagene) followed by Sanger sequencing. The predicted best null mutation was selected by PCR sequencing of the targeted region of *Psmas8* (S3B Fig). The selected mutant allele was 166 bp long versus 222bp of the wild-type. The founder was crossed with wild-type C57BL/6J to eliminate possible unwanted off-targets. *Psmas8*^{+/-} heterozygous mice were re-sequenced and crossed to give rise to *Psmas8*^{-/-} homozygous. Genotyping was performed by analysis of the PCR products of genomic DNA with primers F and R. Mouse mutants for Rec8 and Six6os1 have been previously developed [24, 25].

Histology

For histological analysis of adult testes, mice were perfused and their testes were processed into serial paraffin sections and stained with hematoxylin-eosin or were fixed in Bouin's fixative and stained with Periodic acid-Schiff (PAS) and hematoxylin.

Microscopy

Slides were visualized at room temperature using a microscope (Axioplan 2; Carl Zeiss, Inc.) with 63 × objectives with an aperture of 1.4 (Carl Zeiss, Inc.). Images were taken with a digital camera (ORCA-ER; Hamamatsu) and processed with OPENLAB 4.0.3 and Photoshop (Adobe). Quantification of fluorescence signals was performed using Image J software. Squashed preparations were visualized with a Delta vision microscopy station. Stimulated emission depletion (STED) microscopy (SP8, Leica) was used to generate the super-resolution images. Secondary antibodies for STED imaging were conjugated to Alexa 555 and 488 (Invitrogen). Slides were mounted in Prolong Antifade Gold without DAPI.

Immunocytology

Testes were detunicated and processed for spreading using a conventional "dry-down" technique or squashing [64]. Antibody against the C-term of PSMA8 was a gift from Dr. Murata (Univ of Tokyo, Japan) and has been previously described [10]. Rabbit polyclonal antibodies against PSMA8 were developed by Proteintech (R1 and R2) against a fusion protein of poly-His with full length PSMA8 (pET vector) of mouse origin (see S1 Fig for validation) and was used to validate the immunofluorescence and western results. The primary antibodies used for immunofluorescence were rabbit αSYCP1 IgG ab15090 (1:200) (Abcam), rabbit anti-γH2AX (ser139) IgG #07-164 (1:200) (Millipore), ACA or purified human α-centromere proteins IgG

15–235 (1:5, Antibodies Incorporated), mouse α MLH1 51-1327GR (1:5, BD Biosciences), mouse α SYCP3 IgG sc-74569 (1:100), rabbit α RAD51 PC130 (1:50, Calbiochem), Mouse α CDK1 sc-54 (1:20 IF; 1:1000 wb, Santa Cruz), rabbit α CDK1 Tyr15p #4539 (1:10, Cell Signaling), rabbit α CDK2 sc-6248 (1:20, Santa Cruz), rabbit α PTTG1 serum K783 (1:20 IF, 1:1000 wb), rabbit α TRIP13 19602-1-AP (1:20, Proteintech), rabbit α H2AL2 (1:100, from Dr. Saadi Khochbin), rabbit α PA200 (1:20, Bethyl A303-880A), rabbit α -Caspase3 #9661 (1:30, Cell Signaling), rabbit α H2AK5ac ab45152 (1:20, Abcam), Rabbit α H4K16ac #07–329 (1:50 Millipore), Rabbit α H3ac (K9 and K14) #06–599 (1:20, Millipore), Rabbit α H4ac (K5, K8, K12 and K16) #06–598 (1:20, Millipore), Mouse α Ubiquitin 11023 (1:20 IF, 1:1000 wb, QED Bioscience), Rabbit α HORMAD1 and α HORMAD2 and chicken anti SYCP1 (1:50, from Dr. Attila Toth; [65]), Rabbit anti p-ser10-H3 06–570 (1:100, Millipore), Mouse anti α -tubulin T9026 (1:100, Sigma), Rabbit α Cyclin B1 ab72 (1:20, Abcam), Rabbit α MAD2 (1:30 provided by Dr. Stemmann), Peanut agglutinin lectin L7381 (15 μ g/ml, Sigma), SMC6 ab18039 (1:50, Abcam), Human α VASA 560189 (1:100, BD), Rabbit α INCENP 1186 (1:50, provided by Dr. Earnshaw). TUNEL staining of chromosome spreads was performed with the *in situ* cell death detection kit (Roche).

FACs analysis

Psm8^{+/+} and *Psm8*^{-/-} testicular cells preparation and measurement of their DNA content were performed by a standard procedure [66]. Briefly, the testes were detunicated and the seminiferous tubules were kept in 5 ml of ice-cold separation medium (DMEM supplemented with 10% FCS, 0.1 mM NEAA, 1.5 mM sodium pyruvate, 4 mM L-glutamine and 75 μ g/ml ampicillin). They were treated with 0.1 mg/ml collagenase at 37°C for 10 min under mild shaking. The sedimented seminiferous tubules were washed twice with separation medium and treated for 2 min at 37°C with 2.5 μ g/ml trypsin and 1 U/ml DNase I in separation medium and transferred to ice. Afterwards, single cells were extracted from the seminiferous cords with a Pasteur pipette and filtered through a 40 μ m nylon mesh. The cell suspension (2 $\times$ 10⁶ cells/ml) was diluted 1:1 with a solution containing 0.05 mg/ml propidium iodide and 0.1 mg/ml RNase for 15 min. Finally, the cells were analyzed through flow cytometry in a cytometer FACSCalibur and the BD Cell-Quest software. The cell cycle distribution was analyzed with the Kaluza Analysis software (Beckman Coulter).

Proteasome assay

The 26S proteasome assay was carried out in a total volume of 250 μ l in 96 well plates with 2 mM ATP in 26S buffer using 100 μ g of protein supernatants from whole extracts of mouse testis. Fluorescently labeled substrates employed were: succinyl-Leu-Leu-Val-Tyr-7-amino-4-methylcoumarin (Suc-LLVY-AMC), Z-Ala-Arg-Arg-AMC (Z-ARR-AMC, Bachem), and Z-Leu-Leu-Glu-AMC (Z-LLE-AMC) for the detection of the chymotrypsin- (β 5 catalytic subunit), trypsin- (β 2 catalytic subunit) and caspase- (β 1 catalytic) like activity measurements respectively. The final substrate concentration in each assay was 100 μ M.

Cell lines

The HEK293T, GC1-spg, Leydig TM3, and Sertoli TM4 cell lines were directly purchased at the ATCC and cultured in standard cell media. HEK293T cell line was transfected with Lipofectamine (Invitrogen) or Jetpei (PolyPlus). Cell lines were tested for mycoplasma contamination (Mycoplasma PCR ELISA, Sigma).

Generation of plasmids

Full-length cDNAs encoding PSMA8, PSMA7, CDK1, SYCP1 and SIX6OS1, SYCP3, SYCE2, TEX12, TEX30, PIWIL1 and PIWIL2 were RT-PCR amplified from murine testis RNA. Full-length cDNAs were cloned into the EcoRV pcDNA3-2XFlag or SmaI pEGFP-C1 expression vectors under the CMV promoter. In frame cloning was verified by Sanger sequencing.

Immunoprecipitation and western blotting

200 µg of antibody R1 and R2 were bound to 100 µl of sepharose beads slurry (GE Healthcare). Testis extracts were prepared in 50mM Tris HCl (pH8), 500mM NaCl, 1mM EDTA 1% TritonX-100. 20 mg of proteins extracts were incubated o/n with the Sepharose beads. Protein-bound beads were packed into columns and washed in extracting buffer for three times. Protein were eluted in 100 mM glycine pH3. The whole immunoprecipitation of PSMA8 was performed in a buffer lacking ATP and glycerol to increase the stringency of the interactors and regulators/activators subunits. HEK293T cells were transiently transfected and whole cell extracts were prepared and cleared with protein G Sepharose beads (GE Healthcare) for 1 h. The antibody was added for 2 h and immunocomplexes were isolated by adsorption to protein G-Sepharose beads o/n. After washing, the proteins were eluted from the beads with 2xSDS gel-loading buffer 100mM Tris-HCl (pH 7), 4% SDS, 0.2% bromophenol blue, 200mM β-mercaptoethanol and 20% glycerol, and loaded onto reducing polyacrylamide SDS gels. The proteins were detected by western blotting with the indicated antibodies. Immunoprecipitations were performed using mouse αFlag IgG (5µg; F1804, Sigma-Aldrich), mouse αGFP IgG (4 µg; CSB-MA000051M0m, Cusabio), rabbit αMyc Tag IgG (4µg; #06-549, Millipore), mouse αHA.11 IgG MMS- (5µL, aprox. 10µg/1mg prot; 101R, Covance), ChomPure mouse IgG (5µg/1mg prot; 015-000-003), ChomPure rabbit IgG (5µg/1mg prot.; 011-000-003, Jackson ImmunoResearch), ChomPure goat IgG (5µg/1mg prot.; 005-000-003, Jackson ImmunoResearch). Primary antibodies used for western blotting were rabbit αFlag IgG (1:2000; F7425 Sigma-Aldrich), goat αGFP IgG (sc-5385, Santa Cruz) (1:3000), rabbit αHA IgG (H6908, Sigma-Aldrich) (1:1.000), mouse αMyc obtained from hybridoma cell myc-1-9E10.2 ATCC (1:5). Secondary horseradish peroxidase-conjugated α-mouse (715-035-150, Jackson ImmunoResearch), α-rabbit (711-035-152, Jackson ImmunoResearch), or α-goat (705-035-147, Jackson ImmunoResearch) antibodies were used at 1:5000 dilution. Antibodies were detected by using Immobilon Western Chemiluminescent HRP Substrate from Millipore. Protein extracts for the analysis of SYCP3, CDK1 and CyclinB1 were extracted in Tris-HCl 250mM, SDS10%, Glycerol 50% (denaturing buffer).

MS/MS data analysis

Raw MS data were analyzed using MaxQuant (v. 1.5.7.4) and Perseus (v. 1.5.6.0) programmes 71. Searches were generated versus the *Mus musculus* proteome (UP000000589, May 2017 release) and Maxquant contaminants. All FDRs were of 1%. Variable modifications taken into account were oxidation of M, acetylation of the N-term and ubiquitylation remnants di-Gly and LRGG, while fixed modifications included considered only carbamidomethylation of C. The maximum number of modifications allowed per peptide was 5. For the case of the protein group of CDK1 to 3, experimental results showed that the protein detected was CDK1. For the PSMA8 antibodies R1 and R2, ratios of their respective iBAQ intensity versus the correspondent iBAQ intensity in the control sample were calculated. Proteins with ratio higher or equal to 5 and two or more unique peptides for at least one RP antibody were selected for ulterior analysis. Additionally, in order to avoid filtering rare proteins, those with at least one unique

peptide and one peptide for both Rabbit antibodies (R1 and R2) and none for anti-IgG were also selected for further analysis.

Functional and pathway analysis

GO and KEGG over-representation tests were performed using the R package *clusterProfiler* [67] using standard parameters except for a FDR cutoff of 0.01. KEGG pathways where some key genes (TRIP13, CDK1, SYCP1, DDX4, SYCP3, SYCE3, SIX6OS1) operate and the role of the co-immunoprecipitated proteins were studied using the R package *pathview* [68].

Statistics

In order to compare counts between genotypes at different stages, we used the Welch's t-test (unequal variances t-test), which was appropriate as the count data were not highly skewed (i.e., were reasonably approximated by a normal distribution) and in most cases showed unequal variance. We applied a two-sided test in all the cases. Asterisks denote statistical significance: *p-value <0.01, **p-value <0.001 and ***p-value <0.0001.

Ethics statement

Mice were housed in a temperature-controlled facility (specific pathogen free, spf) using individually ventilated cages, standard diet and a 12 h light/dark cycle, according to EU laws at the "Servicio de Experimentación Animal, SEA". Mouse protocols were approved by the Ethics Committee for Animal Experimentation of the University of Salamanca (USAL). We made every effort to minimize suffering and to improve animal welfare. Blinded experiments were not possible since the phenotype was obvious between wild type and *Psma8*-deficient mouse for all of the experimental procedures used. No randomization methods were applied since the animals were not divided in groups/treatments. The minimum size used for each analysis was two animals/genotype.

Supporting information

S1 Fig. Validation of the antibodies raised against PSMA8. (A) HEK293T cells were transfected with a plasmid encoding PSMA8-GFP, PSMA7-GFP or GFP and the whole extracts were analyzed by western blot using rabbit α -PSMA8 C-terminal (left panel, α 4S), rabbit α -PSMA8 (central panel, R2) and α -GFP (right panel, GFP). Immunodetection of β -actin was used as loading control. The rabbit α - α 4S antibody detected exclusively the 60 kDa band representing PSMA8-GFP. The rabbit α -PSMA8 R2 antibody detected both bands representing PSMA8-GFP and PSMA7-GFP. The bands of 60 kDa (PSMA7 and PSMA8) and 30 kDa (GFP) were all detected with the goat α -GFP validating the experiments. (B) Immunofluorescence of HEK293T cells transfected with plasmids encoding PSMA8-GFP, PSMA7-GFP or GFP. Both PSMA8 and PSMA7 were detected with rabbit α -PSMA8-R2 (red) and GFP by direct fluorescence signal (green). Green and red signals co-localize in the cytoplasm of the transfected HEK293T cells. The experiments were reproduced three times. Bar represents 10 μ m. (TIF)

S2 Fig. Localization of PSMA8 in mouse spermatocytes. (A) Double immunolabeling of endogenous PSMA8 (R2 antibody, green) and SYCP3 (red) in mouse spermatocytes. From the leptotene to zygotene stage, PSMA8 is detected at the synapsed autosomal LEs. At pachytene, PSMA8 is located at the totally synapsed axes and at the PAR of the sex XY bivalent. In diplotene, PSMA8 localizes at the still synapsed AEs and disappears at diakinesis. (B) Double

immunolabeling of spermatocytes spread preparations with PSMA8 (green) and SYCP1 (red), showing that PSMA8 localizes to the synapsed LEs but do not perfectly co-localize with SYCP1 (upper panel). Magnification of the XY bivalent (lower panel) showing the PAR (arrow). Bars represent 10 μ m (A and B, upper panel) and 1.5 μ m (B, lower panel).

(TIF)

S3 Fig. Generation and genetic characterization of *Pisma8*-deficient mice. (A) Diagrammatic representation of the mouse *Pisma8* locus (WT) and the genome editing strategy showing the sgRNAs located on exon 1 and intron 1 (see [methods](#)), the corresponding coding exons (light grey) and non-coding exons (open boxes). Thin (non-coding) and thick (coding sequences) lines under exons represent the expected transcript derived from wild-type (black) and *Pisma8* edited allele (blue). ATG, initiation codon; TGA and *, stop codon. The nucleotide sequence of the 56 base pair deletion derived from PCR amplification of DNA from the *Pisma8*^{edited/edited} is indicated (Δ). Primers (F and R) are represented by arrows. (B) PCR analysis of genomic DNA from three littermate progeny of *Pisma8*^{+/-} heterozygote crosses. The PCR amplification with primers F and R revealed 222 and 166 bp fragments for wild-type and disrupted alleles respectively. Wild-type (WT, +/+), heterozygous (Het, +/-), and homozygous knock-out (KO, -/-) animals. (C) Western blot analysis of protein extracts from wild type testis (P22 and adult), KO testis (P16, P22 and adult) with a specific antibody against the C-terminal (α 4S) and whole recombinant PSMA8 protein (PSMA8-R2). β -actin was used as loading control. The corresponding bands to PSMA8 and PSMA7 are indicated in the right of the panel. Note that at the P22 and in adult stages the intensity of both bands abolishes its independent observation. (D) Double immunofluorescence of spermatocytes at pachytene stage obtained from *Pisma8*^{+/+} and *Pisma8*^{-/-} mice using SYCP3 (red) and PSMA8 (R2 antibody, green). Green labeling in *Pisma8*^{-/-} spermatocytes (49% of the wild type) represents cross-reactivity of the antiserum with PSMA7. Plot under the image panel represents the quantification of intensity from *Pisma8*^{+/+} and *Pisma8*^{-/-} spermatocytes. Welch's *t*-test analysis: * $p < 0.01$. Bar in panel, 10 μ m.

(TIF)

S4 Fig. Validation of the identity of round spermatids with molecular markers. (A) PNA staining (green) of acrosome in spread preparations from wild type and *Pisma8*^{-/-} cells. Double labeling of squash tubules of VASA (chromatoid body), INCENP [1], SMC6 [2] (green) with SYCP3 (red) from wild type and *Pisma8*^{-/-} mice. The combined labeling of INCENP (labels both interkinesis and round spermatids, [1]) and SYCP3 (mainly labels interkinesis with a typical barr patterning at the chromocenters, see below [S4B Fig](#)) is compatible with round spermatids. The combined double immunolabeling of SMC6 (labels both interkinesis and round spermatids, [2]) and SYCP3 (mainly labels interkinesis with a typical barr patterns at the chromocenters, see below [S4B Fig](#)) is also compatible being round spermatids. (B) Double labeling of SYCP3 (green) and ACA (red) showing the different pattern of secondary spermatocytes at interkinesis and round spermatids. Bars in panels represent 10 μ m (A, PNA panel) and 5 μ m (rest of panels).

(TIF)

S5 Fig. Early arrest of *Pisma8*^{-/-} spermatids and gating strategy of the FACs analysis. (A) Immunolabeling of H2AL2 (green) show positive staining in elongating spermatids from wild type mice but lack of staining in *Pisma8*^{-/-} mice. Chromatin was stained with DAPI. Bar represents 10 μ m. (B) Gating strategy employed in the FACs analysis of [Fig 3D](#). Grey dots represent cells that were excluded from the analysis whilst dots included in the polygon represent cells that were employed for the analysis. Red dots enclose 1C cells, blue dots represent 2C cells and

green dots enclose 4C cells.
(TIF)

S6 Fig. Normal synopsis and desynopsis in spermatocytes lacking PSMA8. Double immunolabeling of SYCP3 (red) and SYCP1 (green) showing normal synopsis and desynopsis from early zygotene to diakinesis in *Psm8*^{-/-} in comparison with *Psm8*^{+/+}. Bar represents 10 μm.
(TIF)

S7 Fig. DSBs are generated and repaired as COs in spermatocytes lacking PSMA8. (A) Double immunolabeling of γ-H2AX (green) with SYCP3 (red) in wild-type and *Psm8*^{-/-} spermatocytes from leptotene to diplotene (upper panel). In WT and KO leptoneas, γ-H2AX labels intensely the chromatin. After repair, γ-H2AX labeling remains only in the chromatin of the sex body of the pachynemas. Plot right to the panel represent the quantification of the fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes at leptotene and pachytene. Late round spermatids (LR) but not early round spermatids (ER) from wild type mice show positive staining for γ-H2AX but these highly differentiated cells are lacking in the *Psm8*^{-/-} tubules which are arrested at early round spermatids without γ-H2AX staining (bottom panel). (B) Double immunolabeling of SYCP3 (red) and RAD51 (green). RAD51 foci associates to the AEs in leptoneas spermatocytes of both genotypes (similar number of foci) and dissociate towards pachytene with a similar kinetics. Plot right to the image panel represents the quantification of the number of foci from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes. (C) Double immunolabeling of SYCP3 (red) with MLH1 (green). MLH1 foci are present along each autosomal SC in wild-type and *Psm8*^{-/-} pachynema meocytes in a similar way. Plot right to the panel represents the quantification of the values of the MLH1 foci from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes. Bars represent 10 μm. Welch's *t*-test analysis: * *p*<0.01; ** *p*<0.001; *** *p*<0.0001. Quantification data is indicated in [S3 Table](#).
(TIF)

S8 Fig. PA200 localization in prophase I from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes. Double immunolabeling of PA200 (green) and SYCP3 (red) in chromosome spreads from zygotene to diakinesis. PA200 is detected at the chromosome axes in wild type spermatocytes in contrast to the absence of labeling in *Psm8*^{-/-} spermatocytes. Bar in panels, 10 μm.
(TIF)

S9 Fig. PSMA8 deficiency provokes an slight increase of H2AK5ac at prophase I. Double immunolabeling of H2AK5ac (green) with SYCP3 (red) in wild-type (left panel) and *Psm8*^{-/-} spermatocytes (right panel). In WT and KO spermatocytes chromatin start to be labelled at early pachytene around chromosomes axes. Plots from each panel representing the quantification of fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes are depicted in [Fig 4A](#). Bar represents 10 μm.
(TIF)

S10 Fig. PSMA8 deficiency provokes an slight increase of H3ac at prophase I. Double immunolabeling of H3ac (green) with SYCP3 (red) in wild-type (left panel) and *Psm8*^{-/-} spermatocytes (right panel). Spermatocytes from *Psm8*^{+/+} and *Psm8*^{-/-} show labeling for H3ac at early pachytene in a very diffuse manner surrounding chromosomes axes. Plots from each panel representing the quantification of fluorescence intensity from *Psm8*^{+/+} and *Psm8*^{-/-} spermatocytes are in [Fig 4B](#). Bar represents 10 μm.
(TIF)

S11 Fig. PSMA8 deficiency provokes an slight increase of H4ac at prophase I and in round spermatids. Double immunolabeling of H4ac (green) with SYCP3 (red) in wild-type and

Psma8^{-/-} spermatocytes. Spermatocytes from *Psma8*^{+/+} and *Psma8*^{-/-} show labeling for H4ac in a very diffuse manner surrounding chromosomes from pachytene to metaphase I (right panel). In wild type metaphase I, H4ac labeling appears weakly painting the chromosomes and on some of the centromeres. However, *Psma8*-deficient cells show a more intense labeling specially at the centromeres (lower panel). Round spermatid from *Psma8*^{-/-} accumulates H4ac labeling at the chromatin in comparison with the WT. Plots from each panel representing the quantification of fluorescence intensity from *Psma8*^{+/+} and *Psma8*^{-/-} spermatocytes are in Fig 4C. Bars represent 10 μm.

S12 Fig. PSMA8 deficiency provokes an increase of H4K16ac at prophase I and in metaphase I / round spermatids. Double immunolabeling of H4K16ac (green) with SYCP3 (red) in wild-type and *Psma8*^{-/-} spermatocytes. Spermatocytes from *Psma8*^{+/+} and *Psma8*^{-/-} show labeling for H4K16ac in a very diffuse manner surrounding chromosomes from pachytene to metaphase I (right panel). In wild type metaphase I, H4K16ac labeling appears weakly painting the chromosomes. However, *Psma8*-deficient cells show enhance labeling in the chromosomes of metaphase I cells (lower panel). Round spermatid from *Psma8*^{-/-} accumulates H4K16ac labeling at the chromatin in comparison with the WT. Plots from each panel representing the quantification of fluorescence intensity from *Psma8*^{+/+} and *Psma8*^{-/-} spermatocytes are in Fig 4D. Bars represent 10 μm.

S13 Fig. PSMA8 deficiency alters Ubiquitylation of mouse spermatocytes. (A) Double immunolabeling of Ubiquitin (green) and SYCP3 (red) in mouse chromosome spreads at pachytene stage from *Psma8*^{+/+} and *Psma8*^{-/-} mice. (B) Double immunolabeling of Ubiquitin (green) and SYCP3 (red) in mouse squashed tubules from *Psma8*^{+/+} and *Psma8*^{-/-} mice. Chromatin was stained with DAPI. Bars represent 10 μm (A) and 5 μm (B).

S14 Fig. Lack of co-immunoprecipitation of PSMA8 with candidate interactors. (A-B) HEK293T cells were co-transfected with GFP-TEX30, GFP-PIWIL1, GFP-PIWIL2, GFP-SYCE1, GFP-SYCE2, and GFP-TEX12, and with Flag-PSMA8. PSMA8 does not co-immunoprecipitates (co-IP) with any of them. (C) Positive control was generated by transfecting HEK293T cells with Flag-SYCE2 and GFP-TEX12. Protein complexes were immunoprecipitated overnight with either an anti-Flag or anti-EGFP or IgGs (negative control) and were analyzed by immunoblotting with the indicated antibody.

S15 Fig. CDK1 / Cyclin B1, but not CDK2, are accumulated in *Psma8* mutant spermatocytes. (A) Double immunolabeling of endogenous CDK2 (green) and SYCP3 (red) in WT and KO mouse chromosome spreads at pachytene and metaphase I showing similar labeling at the telomeres and centromeres, respectively. (B) Double immunolabeling of CDK1 (green) and SYCP3 (red) in mouse squashed metaphases I from *Psma8*^{+/+} and *Psma8*^{-/-} mice showing CDK1 accumulation. Plot right to the panel represents the quantification of total CDK1 fluorescence intensity from *Psma8*^{+/+} and *Psma8*^{-/-} metaphase I cells. (C) Double immunolabeling of Cyclin B1 (green) and SYCP3 (red) in mouse squashed tubules from *Psma8*^{+/+} and *Psma8*^{-/-} mice showing CyclinB1 accumulation. Plot right to the panel represents the quantification of total CyclinB1 fluorescence intensity in metaphase I cells. Bars represent 10 μm (A), and 5 μm (B,C). Welch's *t*-test analysis: * *p*<0.01; ** *p*<0.001; *** *p*<0.0001.

S16 Fig. HORMADs are not affected by the increased expression of TRIP13 in the *Psma8*^{-/-} spermatocytes. (A-B) Double immunolabeling of HORMAD1 (A) and HORMAD2 (B) (green) with SYCP3 (red) in *Psma8*^{+/+} and *Psma8*^{-/-} spermatocytes at zygotene and pachytene stages. As synapsis progresses HORMAD1 and HORMAD2 are released from the AEs and maintained at the AE of the sex body similarly in the wild type and in the mutant spermatocytes. Bars represent 10 μ m.
(TIF)

S17 Fig. PTTG1 expression is not altered in the absence of PSMA8. Double immunofluorescence of PTTG1 (green) and SYCP3 (red) in metaphase I cells showing similar expression levels of PTTG1. Plot under the panel represents the quantification of the fluorescence intensity from *Psma8*^{+/+} and *Psma8*^{-/-} metaphase I cells. Bar in panels, 10 μ m. Welch's *t*-test analysis: * $p < 0.01$; ** $p < 0.001$; *** $p < 0.0001$.
(TIF)

S1 Table. Fertility assessment of *Psma8*^{+/+}, *Psma8*^{+/-} and *Psma8*^{-/-} mice.
(PDF)

S2 Table. Quantification of metaphases I/II in *Psma8*^{-/-} testis. (A) Quantification of the proportion of tubules with metaphase I/II in PAS stained tubule sections from the histology example shown in Fig 2B. (B) Quantification of the number of metaphase I and II cells present in p-Ser10-H3 stained tubules that show meiotic divisions (Fig 2C). (C) Quantification of the percentage of metaphases-anaphases I and metaphases-anaphases II in squash preparations (double immunolabeled with ACA and SYCP3) measured as the N° of Metaphase-Anaphase I/II divided by the N° of cells (prophase I + Metaphase-Anaphase I + Interkinesis + Metaphase-Anaphase II) (Fig 2D). Apoptotic Metaphase-Anaphase I and Metaphase-Anaphase II within each genotype are indicated.
(PDF)

S3 Table. Quantification of γ H2AX levels, RAD51 foci, and MLH1 foci (S7 Fig).
(PDF)

S4 Table. Proteasome subunits and proteasome regulators co-immunoprecipitated with PSMA8 from *Psma8*^{+/+} and *Psma8*^{-/-} testis protein extracts using anti-PSMA8 R2 antibody.
(PDF)

S5 Table. Proteasome subunits and proteasome regulators co-immunoprecipitated with PSMA8 selected after analysis and filtering of the data.
(PDF)

S6 Table. Selection of some of the proteasome-related proteins co-immunoprecipitated with PSMA8 selected after analysis and filtering of the data.
(PDF)

S7 Table. Quantification of the percentage of spermatocytes showing SYCP3 aggregates during prophase I stages in squash of seminiferous tubules of *Psma8*^{+/+} and *Psma8*^{-/-} testis. They have been classified in cells with small or large aggregates (n = 2 mice).
(PDF)

S1 Text. Exploratory representation of representative KEGG pathways. (A) Cell cycle (mmu04110). (B) Progesterone-mediated oocyte maturation (mmu04914). (C) Oocyte meiosis (mmu04114). In red, proteins detected in the co-IP experiment over the established cut-off.
(HTM)

S2 Text. Supporting information references.
(DOCX)

Acknowledgments

We wish to express our sincere thanks to Drs. Liu (Univ. of Toledo, USA), Dr Schimenti (Cornell Univ, USA), A. Toth (Dresden Univ. Germany), S. Khochbin (Univ. of Grenoble, France), Dr. Stemmann (Bayreuth Univ., Germany), Dr. Earnshaw, Dr. Murata (Univ of Tokyo, Japan) and Dr M. Sacristan for providing antibodies (TRIP13, Hormad1, Hormad2, SYCP1, H2AL2, MAD2, INCENP, PSMA8, and CDK1) and reagents (plasmid and REC8 mutant mice).

Author Contributions

Conceptualization: Elena Llano, Alberto M. Pendas.

Data curation: Laura Gómez-H.

Formal analysis: Laura Gómez-H, Dirk G. de Rooij, Alberto M. Pendas.

Funding acquisition: Alberto M. Pendas.

Investigation: Laura Gómez-H, Alberto M. Pendas.

Methodology: Laura Gómez-H, Natalia Felipe-Medina, Yazmine B. Condezo, Isabel Ramos, Manuel Sánchez-Martín, Dirk G. de Rooij.

Project administration: Alberto M. Pendas.

Resources: Natalia Felipe-Medina, José Luis Barbero, Ignasi Roig, Manuel Sánchez-Martín, Alberto M. Pendas.

Software: Rodrigo Garcia-Valiente.

Supervision: Alberto M. Pendas.

Validation: Laura Gómez-H, José Angel Suja.

Visualization: Laura Gómez-H, José Angel Suja.

Writing – original draft: Laura Gómez-H, Alberto M. Pendas.

Writing – review & editing: Laura Gómez-H, Dirk G. de Rooij, Elena Llano, Alberto M. Pendas.

References

1. Collins GA, Goldberg AL. The Logic of the 26S Proteasome. *Cell*. 2017; 169(5):792–806. <https://doi.org/10.1016/j.cell.2017.04.023> PMID: 28525752
2. Schmidt M, Haas W, Crosas B, Santamaria PG, Gygi SP, Walz T, et al. The HEAT repeat protein Bim10 regulates the yeast proteasome by capping the core particle. *Nature structural & molecular biology*. 2005; 12(4):294–303.
3. Murata S, Yashiroda H, Tanaka K. Molecular mechanisms of proteasome assembly. *Nature reviews Molecular cell biology*. 2009; 10(2):104–15. <https://doi.org/10.1038/nrm2630> PMID: 19165213
4. Finley D. Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annual review of biochemistry*. 2009; 78:477–513. <https://doi.org/10.1146/annurev.biochem.78.081507.101607> PMID: 19489727
5. Qian MX, Pang Y, Liu CH, Haratake K, Du BY, Ji DY, et al. Acetylation-mediated proteasomal degradation of core histones during DNA repair and spermatogenesis. *Cell*. 2013; 153(5):1012–24. <https://doi.org/10.1016/j.cell.2013.04.032> PMID: 23706739

6. Khor B, Bredemeyer AL, Huang CY, Turnbull IR, Evans R, Maggi LB Jr., et al. Proteasome activator PA200 is required for normal spermatogenesis. *Molecular and cellular biology*. 2006; 26(8):2999–3007. <https://doi.org/10.1128/MCB.26.8.2999-3007.2006> PMID: 16581775
7. Cascio P, Hilton C, Kisselev AF, Rock KL, Goldberg AL. 26S proteasomes and immunoproteasomes produce mainly N-extended versions of an antigenic peptide. *The EMBO journal*. 2001; 20(10):2357–66. <https://doi.org/10.1093/emboj/20.10.2357> PMID: 11350924
8. Griffin TA, Nandi D, Cruz M, Fehling HJ, Kaer LV, Monaco JJ, et al. Immunoproteasome assembly: cooperative incorporation of interferon gamma (IFN-gamma)-inducible subunits. *The Journal of experimental medicine*. 1998; 187(1):97–104. <https://doi.org/10.1084/jem.187.1.97> PMID: 9419215
9. Murata S, Sasaki K, Kishimoto T, Niwa S, Hayashi H, Takahama Y, et al. Regulation of CD8+ T cell development by thymus-specific proteasomes. *Science*. 2007; 316(5829):1349–53. <https://doi.org/10.1126/science.1141915> PMID: 17540904
10. Uechi H, Hamazaki J, Murata S. Characterization of the testis-specific proteasome subunit alpha4s in mammals. *The Journal of biological chemistry*. 2014; 289(18):12365–74. <https://doi.org/10.1074/jbc.M114.558866> PMID: 24668818
11. Kisselev AF, Akopian TN, Castillo V, Goldberg AL. Proteasome active sites allosterically regulate each other, suggesting a cyclical bite-chew mechanism for protein breakdown. *Molecular cell*. 1999; 4(3):395–402. PMID: 10518220
12. Inobe T, Matouschek A. Paradigms of protein degradation by the proteasome. *Current opinion in structural biology*. 2014; 24:156–64. <https://doi.org/10.1016/j.sbi.2014.02.002> PMID: 24632559
13. Guo X, Wang X, Wang Z, Banerjee S, Yang J, Huang L, et al. Site-specific proteasome phosphorylation controls cell proliferation and tumorigenesis. *Nature cell biology*. 2016; 18(2):202–12. <https://doi.org/10.1038/ncb3289> PMID: 26655835
14. Belle A, Tanay A, Bitincka L, Shamir R, O'Shea EK. Quantification of protein half-lives in the budding yeast proteome. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103(35):13004–9. <https://doi.org/10.1073/pnas.0605420103> PMID: 16916930
15. Glotzer M, Murray AW, Kirschner MW. Cyclin is degraded by the ubiquitin pathway. *Nature*. 1991; 349(6305):132–8. <https://doi.org/10.1038/349132a0> PMID: 1846030
16. Meyer HJ, Rape M. Enhanced protein degradation by branched ubiquitin chains. *Cell*. 2014; 157(4):910–21. <https://doi.org/10.1016/j.cell.2014.03.037> PMID: 24813613
17. Rao HB, Qiao H, Bhatt SK, Bailey LR, Tran HD, Bourne SL, et al. A SUMO-ubiquitin relay recruits proteasomes to chromosome axes to regulate meiotic recombination. *Science*. 2017; 355(6323):403–7. <https://doi.org/10.1126/science.aaf6407> PMID: 28059716
18. Lake CM, Hawley RS. Becoming a crossover-competent DSB. *Seminars in cell & developmental biology*. 2016; 54:117–25.
19. Zickler D, Kleckner N. Recombination, Pairing, and Synapsis of Homologs during Meiosis. *Cold Spring Harbor perspectives in biology*. 2015; 7(6).
20. Kim J, Ishiguro K, Nambu A, Akiyoshi B, Yokobayashi S, Kagami A, et al. Meikin is a conserved regulator of meiosis-I-specific kinetochore function. *Nature*. 2015; 517(7535):466–71. <https://doi.org/10.1038/nature14097> PMID: 25533956
21. Baudat F, Imai Y, de Massy B. Meiotic recombination in mammals: localization and regulation. *Nature reviews Genetics*. 2013; 14(11):794–806. <https://doi.org/10.1038/nrg3573> PMID: 24136506
22. Ahuja JS, Sandhu R, Mainpal R, Lawson C, Henley H, Hunt PA, et al. Control of meiotic pairing and recombination by chromosomally tethered 26S proteasome. *Science*. 2017; 355(6323):408–11. <https://doi.org/10.1126/science.aaf4778> PMID: 28059715
23. Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science*. 2015; 348(6235):648–60. <https://doi.org/10.1126/science.1262110> PMID: 25954001
24. Gomez HL, Felipe-Medina N, Sanchez-Martin M, Davies OR, Ramos I, Garcia-Tunon I, et al. C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nature communications*. 2016; 7:13298. <https://doi.org/10.1038/ncomms13298> PMID: 27796301
25. Bannister LA, Reinholdt LG, Munroe RJ, Schimenti JC. Positional cloning and characterization of mouse mei8, a disrupted allele of the meiotic cohesin Rec8. *Genesis*. 2004; 40(3):184–94. <https://doi.org/10.1002/gene.20085> PMID: 15515002
26. Barral S, Morozumi Y, Tanaka H, Montellier E, Govin J, de Dieuleveuil M, et al. Histone Variant H2A.L.2 Guides Transition Protein-Dependent Protamine Assembly in Male Germ Cells. *Molecular cell*. 2017; 66(1):89–101 e8. <https://doi.org/10.1016/j.molcel.2017.02.025> PMID: 28366643

27. Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *The Journal of biological chemistry*. 1998; 273(10):5858–68. <https://doi.org/10.1074/jbc.273.10.5858> PMID: 9488723
28. Mimitou EP, Symington LS. Nucleases and helicases take center stage in homologous recombination. *Trends in biochemical sciences*. 2009; 34(5):264–72. <https://doi.org/10.1016/j.tibs.2009.01.010> PMID: 19375328
29. Dai J, Voloshin O, Potapova S, Camerini-Otero RD. Meiotic Knockdown and Complementation Reveals Essential Role of RAD51 in Mouse Spermatogenesis. *Cell reports*. 2017; 18(6):1383–94. <https://doi.org/10.1016/j.celrep.2017.01.024> PMID: 28178517
30. Qiao H, Prasada Rao HB, Yang Y, Fong JH, Cloutier JM, Deacon DC, et al. Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nature genetics*. 2014; 46(2):194–9. <https://doi.org/10.1038/ng.2858> PMID: 24390283
31. Shinohara M, Oh SD, Hunter N, Shinohara A. Crossover assurance and crossover interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nature genetics*. 2008; 40(3):299–309. <https://doi.org/10.1038/ng.83> PMID: 18297071
32. Moens PB, Marcon E, Shore JS, Kochakpour N, Spyropoulos B. Initiation and resolution of interhomolog connections: crossover and non-crossover sites along mouse synaptonemal complexes. *Journal of cell science*. 2007; 120(Pt 6):1017–27. <https://doi.org/10.1242/jcs.03394> PMID: 17344431
33. Gaucher J, Boussouar F, Montellier E, Curtet S, Buchou T, Bertrand S, et al. Bromodomain-dependent stage-specific male genome programming by Brdt. *The EMBO journal*. 2012; 31(19):3809–20. <https://doi.org/10.1038/emboj.2012.233> PMID: 22922464
34. Lu LY, Wu J, Ye L, Gavrilina GB, Saunders TL, Yu X. RNF8-dependent histone modifications regulate nucleosome removal during spermatogenesis. *Developmental cell*. 2010; 18(3):371–84. <https://doi.org/10.1016/j.devcel.2010.01.010> PMID: 20153262
35. Gomes AV, Young GW, Wang Y, Zong C, Eghbali M, Drews O, et al. Contrasting proteome biology and functional heterogeneity of the 20 S proteasome complexes in mammalian tissues. *Molecular & cellular proteomics: MCP*. 2009; 8(2):302–15.
36. da Cruz I, Rodriguez-Casuriaga R, Santinaque FF, Farias J, Curti G, Capoano CA, et al. Transcriptome analysis of highly purified mouse spermatogenic cell populations: gene expression signatures switch from meiotic-to postmeiotic-related processes at pachytene stage. *BMC genomics*. 2016; 17:294. <https://doi.org/10.1186/s12864-016-2618-1> PMID: 27094866
37. Bousquet-Dubouch MP, Baudelet E, Guerin F, Matondo M, Uttenweiler-Joseph S, Burlet-Schiltz O, et al. Affinity purification strategy to capture human endogenous proteasome complexes diversity and to identify proteasome-interacting proteins. *Molecular & cellular proteomics: MCP*. 2009; 8(5):1150–64.
38. Verma R, Chen S, Feldman R, Schieltz D, Yates J, Dohmen J, et al. Proteasomal proteomics: identification of nucleotide-sensitive proteasome-interacting proteins by mass spectrometric analysis of affinity-purified proteasomes. *Molecular biology of the cell*. 2000; 11(10):3425–39. <https://doi.org/10.1091/mbc.11.10.3425> PMID: 11029046
39. Sanchez-Lanzas R, Castano JG. Proteins directly interacting with mammalian 20S proteasomal subunits and ubiquitin-independent proteasomal degradation. *Biomolecules*. 2014; 4(4):1140–54. <https://doi.org/10.3390/biom4041140> PMID: 25534281
40. Dittmar G, Selbach M. Deciphering the Ubiquitin Code. *Molecular cell*. 2017; 65(5):779–80. <https://doi.org/10.1016/j.molcel.2017.02.011> PMID: 28257698
41. de Vries FA, de Boer E, van den Bosch M, Baarends WM, Ooms M, Yuan L, et al. Mouse Sycp1 functions in synaptonemal complex assembly, meiotic recombination, and XY body formation. *Genes & development*. 2005; 19(11):1376–89.
42. Mikolcevic P, Isoda M, Shibuya H, del Barco Barrantes I, Igea A, Suja JA, et al. Essential role of the Cdk2 activator RingoA in meiotic telomere tethering to the nuclear envelope. *Nature communications*. 2016; 7:11084. <https://doi.org/10.1038/ncomms11084> PMID: 27025256
43. Roig I, Dowdle JA, Toth A, de Rooij DG, Jasin M, Keeney S. Mouse TRIP13/PCH2 is required for recombination and normal higher-order chromosome structure during meiosis. *PLoS genetics*. 2010; 6(8).
44. Yost S, de Wolf B, Hanks S, Zachariou A, Marozzi C, Clarke M, et al. Biallelic TRIP13 mutations predispose to Wilms tumor and chromosome missegregation. *Nature genetics*. 2017; 49(7):1148–51. <https://doi.org/10.1038/ng.3883> PMID: 28553959
45. Bolcun-Filas E, Rinaldi VD, White ME, Schimenti JC. Reversal of female infertility by Chk2 ablation reveals the oocyte DNA damage checkpoint pathway. *Science*. 2014; 343(6170):533–6. <https://doi.org/10.1126/science.1247671> PMID: 24482479

46. Wojtasz L, Daniel K, Roig I, Bolcun-Filas E, Xu H, Boonsanay V, et al. Mouse HORMAD1 and HORMAD2, two conserved meiotic chromosomal proteins, are depleted from synapsed chromosome axes with the help of TRIP13 AAA-ATPase. *PLoS genetics*. 2009; 5(10):e1000702. <https://doi.org/10.1371/journal.pgen.1000702> PMID: 19851446
47. Wang K, Sturt-Gillespie B, Hittle JC, Macdonald D, Chan GK, Yen TJ, et al. Thyroid hormone receptor interacting protein 13 (TRIP13) AAA-ATPase is a novel mitotic checkpoint-silencing protein. *The Journal of biological chemistry*. 2014; 289(34):23928–37. <https://doi.org/10.1074/jbc.M114.585315> PMID: 25012665
48. Wojtasz L, Cloutier JM, Baumann M, Daniel K, Varga J, Fu J, et al. Meiotic DNA double-strand breaks and chromosome asynapsis in mice are monitored by distinct HORMAD2-independent and -dependent mechanisms. *Genes & development*. 2012; 26(9):958–73.
49. Daniel K, Lange J, Hached K, Fu J, Anastasiadis K, Roig I, et al. Meiotic homologue alignment and its quality surveillance are controlled by mouse HORMAD1. *Nature cell biology*. 2011; 13(5):599–610. <https://doi.org/10.1038/ncb2213> PMID: 21478856
50. Nelson CR, Hwang T, Chen PH, Bhalla N. TRIP13PCH-2 promotes Mad2 localization to unattached kinetochores in the spindle checkpoint response. *The Journal of cell biology*. 2015; 211(3):503–16. <https://doi.org/10.1083/jcb.201505114> PMID: 26527744
51. Davies OR, Maman JD, Pellegrini L. Structural analysis of the human SYCE2-TEX12 complex provides molecular insights into synaptonemal complex assembly. *Open biology*. 2012; 2(7):120099. <https://doi.org/10.1098/rsob.120099> PMID: 22870393
52. Winkel K, Alsheimer M, Ollinger R, Benavente R. Protein SYCP2 provides a link between transverse filaments and lateral elements of mammalian synaptonemal complexes. *Chromosoma*. 2009; 118(2):259–67. <https://doi.org/10.1007/s00412-008-0194-0> PMID: 19034475
53. Miake H, Mizusawa H, Iwatsubo T, Hasegawa M. Biochemical characterization of the core structure of alpha-synuclein filaments. *The Journal of biological chemistry*. 2002; 277(21):19213–9. <https://doi.org/10.1074/jbc.M110551200> PMID: 11893734
54. Huang L, Haratake K, Miyahara H, Chiba T. Proteasome activators, PA28gamma and PA200, play indispensable roles in male fertility. *Scientific reports*. 2016; 6:23171. <https://doi.org/10.1038/srep23171> PMID: 27003159
55. Jiang H, Gao Q, Zheng W, Yin S, Wang L, Zhong L, et al. MOF influences meiotic expansion of H2AX phosphorylation and spermatogenesis in mice. *PLoS genetics*. 2018; 14(5):e1007300. <https://doi.org/10.1371/journal.pgen.1007300> PMID: 29795555
56. Yang Q, Ferrell JE Jr. The Cdk1-APC/C cell cycle oscillator circuit functions as a time-delayed, ultrasensitive switch. *Nature cell biology*. 2013; 15(5):519–25. <https://doi.org/10.1038/ncb2737> PMID: 23624406
57. Hellmuth S, Pohlmann C, Brown A, Bottger F, Sprinzl M, Stemmann O. Positive and negative regulation of vertebrate separase by Cdk1-cyclin B1 may explain why securin is dispensable. *The Journal of biological chemistry*. 2015; 290(12):8002–10. <https://doi.org/10.1074/jbc.M114.615310> PMID: 25659430
58. Rattani A, Vinod PK, Godwin J, Tachibana-Konwalski K, Wolna M, Malumbres M, et al. Dependency of the spindle assembly checkpoint on Cdk1 renders the anaphase transition irreversible. *Current biology*. 2014; 24(6):630–7. <https://doi.org/10.1016/j.cub.2014.01.033> PMID: 24583015
59. Bolcun-Filas E, Hall E, Speed R, Taggart M, Grey C, de Massy B, et al. Mutation of the mouse Syce1 gene disrupts synapsis and suggests a link between synaptonemal complex structural components and DNA repair. *PLoS genetics*. 2009; 5(2):e1000393. <https://doi.org/10.1371/journal.pgen.1000393> PMID: 19247432
60. Bolcun-Filas E, Costa Y, Speed R, Taggart M, Benavente R, De Rooij DG, et al. SYCE2 is required for synaptonemal complex assembly, double strand break repair, and homologous recombination. *The Journal of cell biology*. 2007; 176(6):741–7. <https://doi.org/10.1083/jcb.200610027> PMID: 17339376
61. Schramm S, Fraune J, Naumann R, Hernandez-Hernandez A, Hoog C, Cooke HJ, et al. A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility. *PLoS genetics*. 2011; 7(5):e1002088. <https://doi.org/10.1371/journal.pgen.1002088> PMID: 21637789
62. Agarwal A, Sharma R, Durairajanayagam D, Cui Z, Ayaz A, Gupta S, et al. Differential proteomic profiling of spermatozoal proteins of infertile men with unilateral or bilateral varicocele. *Urology*. 2015; 85(3):580–8. <https://doi.org/10.1016/j.urology.2014.11.030> PMID: 25733269
63. Singh P, Schimenti JC, Bolcun-Filas E. A mouse geneticist's practical guide to CRISPR applications. *Genetics*. 2015; 199(1):1–15. <https://doi.org/10.1534/genetics.114.169771> PMID: 25271304
64. Page J, Suja JA, Santos JL, Rufas JS. Squash procedure for protein immunolocalization in meiotic cells. *Chromosome research: an international journal on the molecular, supramolecular and evolutionary aspects of chromosome biology*. 1998; 6(8):639–42.

65. Papanikos F, Clement JAJ, Testa E, Ravindranathan R, Grey C, Dereli I, et al. Mouse ANKRD31 Regulates Spatiotemporal Patterning of Meiotic Recombination Initiation and Ensures Recombination between X and Y Sex Chromosomes. *Molecular cell*. 2019.
66. Malkov M, Fisher Y, Don J. Developmental schedule of the postnatal rat testis determined by flow cytometry. *Biol Reprod*. 1998; 59(1):84–92. <https://doi.org/10.1095/biolreprod59.1.84> PMID: 9674997
67. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology*. 2012; 16(5):284–7. <https://doi.org/10.1089/omi.2011.0118> PMID: 22455463
68. Brouwer CA, Postma A, Hooimeijer HL, Smit AJ, Vonk JM, van Roon AM, et al. Endothelial damage in long-term survivors of childhood cancer. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2013; 31(31):3906–13.

BIOCHEMISTRY

Meiotic chromosome synapsis depends on multivalent SYCE1-SIX6OS1 interactions that are disrupted in cases of human infertility

Fernando Sánchez-Sáez^{1*}, Laura Gómez-H^{1*}, Orla M. Dunne², Cristina Gallego-Páramo², Natalia Felipe-Medina¹, Manuel Sánchez-Martín³, Elena Llano⁴, Alberto M. Pendas^{1†}, Owen R. Davies^{2†}

Copyright © 2020
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim to
original U.S. Government
Works. Distributed
under a Creative
Commons Attribution
License 4.0 (CC BY).

Meiotic reductional division depends on the synaptonemal complex (SC), a supramolecular protein assembly that mediates homologous chromosomes synapsis and promotes crossover formation. The mammalian SC has eight structural components, including SYCE1, the only central element protein with known causative mutations in human infertility. We combine mouse genetics, cellular, and biochemical studies to reveal that SYCE1 undergoes multivalent interactions with SC component SIX6OS1. The N terminus of SIX6OS1 binds and disrupts SYCE1's core dimeric structure to form a 1:1 complex, while their downstream sequences provide a distinct second interface. These interfaces are separately disrupted by SYCE1 mutations associated with nonobstructive azoospermia and premature ovarian failure (POF), respectively. Mice harboring SYCE1's POF mutation and a targeted deletion within SIX6OS1's N terminus are infertile with failure of chromosome synapsis. We conclude that both SYCE1-SIX6OS1 binding interfaces are essential for SC assembly, thus explaining how SYCE1's reported clinical mutations give rise to human infertility.

INTRODUCTION

Meiotic cell division is defined by a unique and highly dynamic program of events that result in homologous chromosome synapsis, crossover (CO) formation, and subsequent homolog segregation into haploid germ cells (1–3). Homologous chromosome pairs are established through interhomolog recombination searches from up to 400 induced double-strand breaks (DSBs) per cell (4). Once established, local recombination-mediated alignments are converted into the single continuous synapsis of aligned homologous chromosomes through the zipper-like assembly of the synaptonemal complex (SC) (5). The SC's supramolecular protein structure mediates continuous 100-nm tethering between homologous chromosome axes and provides the necessary three-dimensional framework for crossover formation (2). Following SC disassembly, crossovers provide the sole physical links between homologs at metaphase I, so are essential for ensuring correct homolog segregation in addition to providing genetic diversity (2).

The SC has an iconic and highly conserved tripartite structure that has been observed across meiotically reproducing eukaryotes (6). This consists of lateral elements (LEs) that coat the two homologous chromosome axes and a midline central element (CE), with a series of transverse filaments that bind together these longitudinal electron-dense structures (Fig. 1A) (7). The protein components of the mammalian SC have been identified as transverse filaments protein SYCP1 (Synaptonemal complex protein 1) (8), CE proteins

SYCE1, SYCE2, and SYCE3 (Synaptonemal complex central element proteins 1 to 3), SIX6OS1, and TEX12 (Testis-expressed protein 12) (9–12), and LE proteins SYCP2 and SYCP3 (13, 14). All transverse filament and CE components are essential for SC assembly, and their individual disruption leads to infertility owing to meiotic arrest with failure of DSB repair (10, 11, 15–18). In contrast, disruption of LE components produces a sexual dimorphism of male infertility and female subfertility (19, 20), with SYCP3 deficiency in females promoting germ cell aneuploidy and embryonic death (21).

In recent years, a variety of cellular imaging, biochemical and structural biology approaches have begun to uncover the molecular structures, interactions, and mechanisms responsible for mammalian SC assembly. SYCP1 self-assembles into a supramolecular lattice that provides the underlying 100-nm synapsis between chromosome axes (22, 23), while SYCP3 assembles into regularly repeating filaments that support chromosomal looping (24, 25). The five CE proteins provide essential structural supports for the SYCP1 lattice that enable its continuous and cooperative extension along the entire chromosome length. In this capacity, CE proteins have been categorized as synaptic initiation factors (SYCE3, SYCE1, and SIX6OS1) and elongation factors (SYCE2 and TEX12), of which their disruption leads to complete loss of tripartite SC structure and failure of extension of short SC-like stretches, respectively (10, 11, 16–18). Of synaptic initiation factors, SYCE3 forms dimers that undergo potentially limitless self-assembly (26, 27), SYCE1 forms antiparallel dimeric assemblies (28), and SIX6OS1 is an SYCE1-interacting protein of unknown structure (11). These likely act as short-range structural supports between SYCP1 molecules, possibly in transverse, longitudinal, and vertical orientations to stabilize a local three-dimensional SYCP1 lattice (22). In contrast, SYCE2 and TEX12 exist as a seemingly constitutive complex that undergoes self-assembly into fibers of many micrometers in length (29), which likely provide the long-range structural supports that stabilize continuous growth of the SYCP1 lattice along the entire chromosome axis (22).

¹Molecular Mechanisms Program, Centro de Investigación del Cáncer and Instituto de Biología Molecular y Celular del Cáncer (CSIC-Universidad de Salamanca), Salamanca, Spain. ²Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, UK. ³Departamento de Medicina, Universidad de Salamanca, Salamanca, Spain. ⁴Departamento de Fisiología y Farmacología, Universidad de Salamanca, Salamanca, Spain.

*These authors contributed equally to this work.

†Corresponding author. Email: owen.davies@newcastle.ac.uk (O.R.D.); amp@usal.es (A.M.P.)

Fig. 1. SYCE1_{POF} retains its core dimeric structure. (A) Schematic of the SC demonstrating its tripartite structure of two chromosome-bound LEs and a midline CE. Synapsis is achieved through N-terminal head-to-head assembly of SYCP1 molecules, which are bound via their C termini to meiotic chromosomes. SYCP1 head-to-head assembly is structurally supported within the CE by SYCE3 (red), an SYCE1-SIX6OS1 complex (yellow), and SYCE2-TEX12 fibrous assemblies (green). (B) Human SYCE1 (top) and SIX6OS1 (bottom) sequence schematics indicating the location and consequence of infertility-associated mutations of SYCE1 and Δ10–21 internal deletion of SIX6OS1, alongside the principal constructs used in this study. (C) SDS–polyacrylamide gel electrophoresis (SDS–PAGE) analysis of the purified recombinant proteins used in this study. The dominant degradation product of SYCE1_{POF} is indicated by an asterisk; its identity was confirmed by the observed cleavage of degraded MBP- and His-SYCE1_{POF} fusion proteins upon treatment with TEV protease (fig. S1, A and B), consistent with it representing C-terminal degradation down to SYCE1's structural core. *M_w*, weight-average molecular weight. (D) SEC–MALS analysis. SYCE1_{core} (yellow), SYCE1_{POF} (green), and full-length SYCE1 (violet) are dimeric species of 36, 48 (39 kDa for the degradation product), and 86 kDa, respectively (theoretical dimers: 37, 55, and 80 kDa). dRI, differential refractive index. Data for SYCE1_{core} and full-length SYCE1 are reproduced from (28).

Owing to the essential roles of meiotic recombination, synapsis, and chromosome dynamics in mammalian meiosis (15, 30–34), their defects are associated with human infertility, recurrent miscarriage, and aneuploidies (35, 36). As genetic causes of infertility, they typically fall within the category of idiopathic cases, having no readily diagnosable and clinically resolvable cause. Within the 10 to 15% of couples who suffer from infertility, approximately 25% are idiopathic and of likely genetic origin, comprising 50 to 80% of cases of nonobstructive azoospermia (NOA) and premature ovarian failure (POF) (36, 37). While individual infertility mutations are inherently unlikely to become widespread in a population, they can be found within families, especially when consanguineous (38), and provide crucial insights into their common targets and the molecular mechanisms that they disrupt.

Within the SC, familial infertility mutations have been identified for SYCP3 and SYCE1 (36). All identified SYCP3 mutations are autosomal dominant and alter or delete its structural core's C terminus that mediates filamentous assembly, so likely sequester wild-type (WT) molecules into inactive complexes (24, 36). In contrast, the three identified SYCE1 mutations are autosomal recessive and

were found in two familial cases of NOA and one of POF (36). The two NOA cases are splice-site mutations, c.197-2A>G and c.375-2A>G, which are predicted to result in a truncated product of amino acids 1 to 65 and an internal deletion of amino acids 126 to 155, respectively (39, 40). These remove or delete part of human SYCE1's structural core that is encoded by amino acids 25 to 179, so can be explained by disruption of its dimeric structure (Fig. 1B) (28, 36). The POF mutation c.613C>T generates a premature stop codon (p.Gln241*) to give a truncated product of amino acids 1 to 240, relative to the canonical 351–amino acid isoform (Fig. 1B) (41). However, as this truncation lies outside SYCE1's structural core, the molecular mechanism that is disrupted, and thereby responsible for infertility, remains unknown.

Here, we combine mouse genetics and cellular and biochemical studies to reveal a multivalent interaction mode between SYCE1 and SIX6OS1 that is disrupted by infertility-associated mutations of SYCE1. We find that the SIX6OS1 N terminus binds and disrupts the core dimeric structure of SYCE1 (amino acids 25 to 179) to form a 1:1 complex as the first interface, and its downstream sequence binds to SYCE1 amino acids 177 to 305 as the second interface.

SYCE1's infertility-associated mutations c.375-2A>G (NOA) and c.613C>T (POF) specifically disrupt the first and second interfaces, respectively. Mice harboring the SYCE1 POF mutation and a targeted deletion within SIX6OS1 (which disrupts the first interface) are infertile, with failure of SC assembly. We conclude that both SYCE1-SIX6OS1 binding interfaces are essential for SC assembly and meiotic division, thus explaining how human infertility results from the differential targeting of binding interfaces by SYCE1's reported clinical mutations.

RESULTS

SYCE1 POF mutation c.613C>T retains its core dimeric structure

The SYCE1 POF mutation c.613C>T encodes a premature stop codon (p.Gln241*) that is predicted to generate a truncated protein product of amino acids 1 to 240, relative to SYCE1's canonical 351-amino acid isoform (Fig. 1B) (41). We previously demonstrated that an N-terminal structural core encoded by amino acids 25 to 179 (SYCE1core) forms an α -helical antiparallel coiled-coil structure that mediates head-of-head dimerization of SYCE1 (28). As this core region is retained (Fig. 1B), we predicted that SYCE1's antiparallel dimeric structure would be maintained within the 1- to 240-amino acid truncated product of the POF mutation (SYCE1_{POF}). To test this, we purified recombinant SYCE1_{POF}, generating purified material that contained approximately equal quantities of the full protein and a degradation product of apparent size consistent with degradation to the C-terminal boundary of its structural core (Fig. 1C and fig. S1, A and B). Circular dichroism (CD) spectroscopy confirmed that SYCE1_{POF} contains a proportion of α -helical structure consistent with retention of the 25–179 core structure (fig. S1C), and SYCE1_{POF} and SYCE1core demonstrated identical melting temperatures (T_m) of 39°C (fig. S1D). Furthermore, analysis by size exclusion chromatography multiangle light scattering (SEC-MALS) confirmed that the full and degraded proteins are homodimers of 48 and 39 kDa, respectively (Fig. 1D). We conclude that SYCE1_{POF} retains the dimeric structure imposed by its core 25–179 region, so its SC and meiotic defects must result from additional structural or functional roles of its deleted C terminus.

The SYCE1 POF mutation leads to failure of SC assembly and infertility in mice

Having established its retention of core dimeric structure, we next sought to determine the structural and functional consequence of the SYCE1 POF mutation on the SC and meiotic division *in vivo*. We thus generated mice harboring mutations of *Syce1* alleles to introduce stop codons at amino acid position 243, equivalent to the human p.Gln241* mutation (figs. S2 and S3). While heterozygotes (designated *Syce1*^{POF/WT}) were fertile, both male and female homozygotes (designated *Syce1*^{POF/POF}) were infertile, replicating the autosomal recessive pattern of the POF mutation in humans (41). In male mutant mice, we observed reduced testis size (63% smaller, $n = 3$ mice at 2 months of age; fig. S4A) and a zygotene-like arrest similar to that observed in the SYCE1 knockout (16). There was defective SC assembly, with reduced staining for SYCP1 (Fig. 2A) and SYCE3 (Fig. 2B) and no staining for SYCE1 (Fig. 2C), SIX6OS1 (Fig. 2D), and SYCE2-TEX12 (fig. S4, B and C). Analysis of SYCE1 expression in the testis of *Syce1*^{POF/POF} mice confirmed the presence of *Syce1* transcript and a protein product of the correct molecular

weight, albeit at reduced levels in comparison with WT (fig. S4, D and E, and table S1A). The *Syce1*^{POF} open reading frame achieved WT levels of protein expression in a heterologous 293T cellular system (fig. S4F). We next studied the kinetics of DSB repair. Meiotic DSBs are generated by the nuclease SPO11 and are then resected to form single-stranded DNA ends that invade into the homologous chromosome by the recombinases RAD51 (DNA repair protein RAD51 homolog 1) and DMC1 (Meiotic recombination protein DMC1/LIM15 homolog) (42). DSBs are labeled by the presence of phosphorylated H2AX (γ -H2AX) (43). The distribution of γ -H2AX in mutant spermatocytes was similar to that found in WT cells at early prophase I but show increased staining at zygotene-like arrest (Fig. 2E). The distributions of RAD51 and DMC1 were detected on aligned LEs (Fig. 2, F and G) but in absence of mismatch repair protein MLH1 (DNA mismatch repair protein Mlh1) (marker of crossing-overs) (Fig. 2H). Together, these data indicate generation of DSBs but with failure of their repair and CO formation in *Syce1*^{POF/POF}. In female mutant mice, we observed no follicles in adult ovaries (fig. S5A), and embryonic oocytes demonstrated zygotene arrest with mostly unaligned chromosome axes, recapitulating the human POF syndrome. Analysis of the SC revealed similar defects, with reduction in SYCP1 and SYCE3 (Fig. 3, A and B) staining (though to a lesser extent than males), and absence of SYCE1, SIX6OS1 (Fig. 3, C and D), and SYCE2-TEX12 (fig. S5, B and C). The distribution of γ -H2AX, RAD51, and DMC1 labeling in zygotene-like mutant oocytes was also increased and lacked MLH1 foci (Fig. 3, E to H). Thus, the SYCE1 POF mutation leads to male and female infertility with phenotypes of failed DSB repair, synapsis, and lastly SC assembly, similar to those previously observed upon disruption of structural components of the SC CE (10, 11, 16–18).

SYCE1_{POF} retains SIX6OS1 binding but lacks SYCE3 binding in heterologous systems

As the *Syce1*^{POF/POF} mouse strain indicated a clear structural defect in the SC, we wondered whether the POF mutation may disrupt the known interaction between SYCE1 and fellow SC CE components SIX6OS1 and SYCE3 (11). The expression of SYCE1 and SIX6OS1 in COS7 cells produced cytoplasmic signals that became colocalized in foci upon coexpression (95% cells; Fig. 4A and fig. S6), in keeping with our previous findings (11). SYCE1_{POF} formed similar or slightly reduced numbers of foci that equally colocalized with SIX6OS1, indicating a retention of SIX6OS1 binding (89% cells; Fig. 4A). We further demonstrated a similar coimmunoprecipitation of SIX6OS1 by WT SYCE1 and SYCE1_{POF} upon coexpression in human embryonic kidney (HEK) 293 cells (Fig. 4B). Thus, the SYCE1-SIX6OS1 interaction is retained in the SYCE1 POF mutation. Could other disrupted functions contribute to the effect of the POF mutation? The only other known SYCE1 interactor is SYCE3, which undergoes low-affinity binding, as determined by its dissociation during purification (fig. S7, A and B). In contrast with the WT protein, the expression of SYCE1_{POF} (cytoplasmic foci) in COS7 cells failed to recruit SYCE3 (preferentially nuclear) to their cytoplasmic foci (colocalization between SYCE3 and SYCE1 was observed for 95% of cells expressing WT SYCE1 and 21% of cells expressing SYCE1_{POF}; Fig. 4C and fig. S6). Similarly, SYCE1_{POF} failed to coimmunoprecipitate SYCE3 upon coexpression in HEK293 cells (Fig. 4D). Thus, while the SYCE1-SIX6OS1 complex is retained, the low-affinity SYCE1-SYCE3 complex is largely abolished in the SYCE1 POF mutation.

Fig. 2. *Syce1*^{POF/POF} spermatocytes are not able to synapse and DSBs are deficiently repaired. (A) Double immunolabeling of WT pachytene and *Syce1*^{POF/POF} zygotene-like spermatocytes with SYCP3 (red) and SYCP1 (green). In *Syce1*^{POF/POF} spermatocytes, AEs fail to synapse and show a weak staining of SYCP1 along the axial elements (AEs). a.u., arbitrary units. (B to D) Double immunolabeling of spermatocyte spreads with SYCP3 (red) and the CE proteins (green). *Syce1*^{POF/POF} zygotene-like spermatocytes showed a highly reduced signal of SYCE3 (B) and the absence of (C) SYCE1 and (D) SIX6OS1 from the AEs. (E) Double immunolabeling of γ-H2AX (green) and SYCP3 (red) in spermatocyte spreads from WT and *Syce1*^{POF/POF} mice. γ-H2AX staining was persistent in *Syce1*^{POF/POF} zygotene-like spermatocytes, but was restricted to the sex body in WT pachytene cells. (F and G) Double immunofluorescence of (F) RAD51 or (G) DMC1 (green) and SYCP3 (red). *Syce1*^{POF/POF} zygotene-like spermatocytes showed increased numbers of foci of RAD51 and DMC1 along the AEs in comparison with WT, indicating unrepaired DSBs. (H) Double immunolabeling of MLH1 (green) and SYCP3 (red) showing the absence of COs (MLH1) in arrested *Syce1*^{POF/POF} spermatocytes. Fluorescence intensity levels (A, B, and E) and number of foci (F and G) from WT and zygotene-like arrested spermatocytes are quantified in the right-hand plots. Welch's *t* test analysis: ****P* < 0.0001. Scale bars, 10 μm.

Fig. 3. *Syce1*^{POF/POF} oocytes fail to synapse and do not properly repair DSBs. (A) Double immunolabeling of oocyte spreads from WT and *Syce1*^{POF/POF} mice with SYCP3 (red) and SYCP1 (green). *Syce1*^{POF/POF} oocytes became arrested in a zygotene-like stage where AEs remain unsynapsed and unaligned, with reduced levels of SYCP1. (B to D) Double immunolabeling of oocyte spreads with SYCP3 (red) and the CE proteins (green). *Syce1*^{POF/POF} zygotene-like oocytes showed reduced SYCE3 signal (B) and a complete absence of (C) SYCE1 and (D) SIX6OS1 from the AEs. IP, immunoprecipitation. (E) Double immunostaining of spread preparations of WT pachytene and *Syce1*^{POF/POF} zygotene-like oocytes with γ-H2AX (green) and SYCP3 (red). In *Syce1*^{POF/POF} oocytes, the levels of γ-H2AX increased and were more restricted to AEs in comparison with WT pachytene cells. (F to G) Double immunolabeling of (F) RAD51 or (G) DMC1 (green) and SYCP3 (red), showing higher numbers of foci in AEs from mutant oocytes. (H) Labeling of MLH1 (green) and SYCP3 (red). MLH1 foci are absent from the AEs of *Syce1*^{POF/POF} oocytes. Fluorescence intensity levels (A, B, and E) and number of foci (F and G) from WT and *Syce1*^{POF/POF} zygotene-like oocytes are quantified in the right-hand plots. Welch's t test analysis: ****P* < 0.0001. Scale bars, 10 μm.

Fig. 4. SYCE1_{POF} retains SIX6OS1 binding but fails to retain the SYCE3-interaction in heterologous systems. (A) Mouse SIX6OS1 colocalized with mouse SYCE1 and SYCE1_{POF} in a cytoplasmic punctate pattern upon coexpression in COS7 cells; the percentage of cells exhibiting colocalization is shown in the right-hand plot ($n = 100$ cells). DAPI, 4',6-diamidino-2-phenylindole. (B) HEK293T cells were cotransfected with the indicated expression vectors. Protein complexes were immunoprecipitated with anti-Flag or anti-enhanced green fluorescent protein (EGFP) antibodies, or mouse immunoglobulin G (IgG) as a negative control, and were analyzed by immunoblotting with the indicated antibody. GFP-mSIX6OS1 coimmunoprecipitated with Flag-mSYCE1 and Flag-mSYCE1_{POF}, suggesting that the POF mutation of SYCE1 alone is insufficient to block the interaction. (C) COS7 cells were transfected with mouse *Syce1* or *Syce1*_{POF} as indicated. SYCE1 colocalized with SYCE3 in its own cytoplasmic punctate pattern, and colocalization was substantially diminished for SYCE1_{POF} ($n = 100$ cells). (D) Immunoprecipitation of protein complexes from HEK293T-cotransfected cells with an anti-Myc or anti-EGFP antibody or mouse IgG. SYCE1 coimmunoprecipitated with SYCE3, and the interaction was disrupted for SYCE1 POF, suggesting that the C-terminal region of SYCE1 is required for its interaction with SYCE3. The untransfected lanes in (B) and (D) show the absence of all the proteins in total protein extracts from untransfected 293T cells. Scale bars, 20 μ m.

SYCE1 core undergoes conformational change to form a 1:1 complex with SIX6OS1

What is the molecular basis of SIX6OS1 binding by SYCE1? As this is retained in SYCE1^{POF}, we reasoned that SIX6OS1 binding must be mediated by SYCE1's structural core. We screened SYCE1 core against a library of SIX6OS1 constructs through bacterial coexpression and identified a robust interaction with amino acids 1 to 67 of SIX6OS1, herein referred to as SIX6OS1_N (Figs. 1B and 5A). We were able to purify the SYCE1 core-SIX6OS1_N complex by reciprocal affinity chromatography, ion exchange, and size exclusion chromatography (Fig. 5B) and found it to be stable under all experimental conditions tested. We were further able to purify similar complexes for SYCE1^{POF} (with the same degradation product as upon isolated expression) and full-length SYCE1 (Fig. 1C and fig. S1B), confirming that SIX6OS1 binding is retained by all constructs containing the 25–179 core. CD analysis revealed similar α -helical content for SYCE1-SIX6OS1_N complexes as for their isolated SYCE1 proteins (fig. S1C). CD thermal denaturation revealed slightly increased cooperativity of unfolding and melting temperatures for SYCE1-SIX6OS1_N complexes relative to their isolated SYCE1 proteins (increasing from 39° to 43°C, 39° to 41°C, and 38° to 40°C for SYCE1 core, SYCE1^{POF}, and full length, respectively; Fig. 5C and fig. S1D). SEC-MALS analysis revealed that all three SYCE1-SIX6OS1_N complexes are 1:1, with molecular weights of 27, 37, and 46 kDa, respectively (Fig. 5D and fig. S7C). Thus, the SYCE1 core undergoes conformational change from an antiparallel homodimer to a 1:1 complex upon binding to SIX6OS1_N (Fig. 5E).

We analyzed the conformation of the SYCE1 core-SIX6OS1_N complex by size exclusion chromatography small-angle x-ray scattering (SEC-SAXS; fig. S7, D and E). The SAXS real-space pair-distance $P(r)$ distribution (the distribution of interatomic distances within a protein structure) demonstrates positive skew, indicating that SYCE1 core-SIX6OS1_N retains the rod-like structure of SYCE1 core, but with a reduction in its molecular length from 186 to 138 Å (Fig. 5F). Furthermore, its cross-sectional radius is slightly increased from 9 to 11 Å (fig. S7F), suggesting an increase from a two- to four-helical coiled coil. These geometric changes are consistent with the SYCE1 core-SIX6OS1_N 1:1 complex forming a shorter but wider coiled coil than the isolated SYCE1 core dimer, as indicated by their SAXS *ab initio* models (Fig. 5G). Furthermore, the SAXS $P(r)$ distribution of SYCE1^{POF} indicates a similar elongated structure but with an increased tail to a maximum dimension of 180 Å (Fig. 5F), consistent with it containing the same SYCE1 core-SIX6OS1_N structure with an extended and potentially unstructured C terminus to amino acid 240. We conclude that SYCE1 core mediates a direct interaction with SIX6OS1_N that imposes a conformational change to a 1:1 complex that adopts a shorter and wider coiled-coil conformation than the isolated SYCE1 core antiparallel homodimer.

SYCE1^{POF} disrupts a second SYCE1-SIX6OS1 binding interface

Does the SYCE1 core-SIX6OS1_N complex represent the sole means by which SYCE1 interacts with SIX6OS1? We were unable to obtain soluble biochemical complexes containing SIX6OS1 sequences beyond its N terminus and so used yeast two-hybrid (Y2H) to test SYCE1 binding by full-length SIX6OS1. Having confirmed direct binding of SYCE1 core to full-length SIX6OS1, we used C-terminal truncation to dissect its minimal binding site to amino acids 1 to 75, in keeping with our biochemical findings, and identified an addi-

tional interaction between SYCE1 177–305 and full-length SIX6OS1 (Fig. 6A).

To establish whether SYCE1 core and 177–305 bind to the same or distinct sites within SIX6OS1, we established an internal deletion of SIX6OS1 amino acids 10 to 21 (Δ 10–21) that blocks formation of the SYCE1 core-SIX6OS1_N biochemical complex (Fig. 5A). SIX6OS1 1–22 did not interact with any SYCE1 construct (Fig. 6A), indicating that amino acids 10 to 21 are necessary but not sufficient for SYCE1 core binding. While Δ 10–21 completely abrogated the Y2H interaction of full-length SIX6OS1 with SYCE1 core (25–179), it retained a robust interaction with SYCE1 177–305, suggesting distinct SIX6OS1-binding sites (Fig. 6A). Furthermore, Δ 10–21 blocked the ability of SIX6OS1 1–262 to interact with SYCE1 core and SYCE1^{POF} (amino acids 25 to 240) while retaining its binding to full-length and 25–315 SYCE1 (Fig. 6A). Thus, SYCE1 undergoes multivalent interactions with SIX6OS1, with the first binding interface mediated by SYCE1 core and SIX6OS1_N (1–67), and the second interface mediated by SYCE1 177–305 and downstream sequence within SIX6OS1 1–262. Furthermore, the first and second binding interfaces are specifically disrupted by SIX6OS1 deletion Δ 10–21 and the SYCE1 POF mutation, respectively, and in both cases, an SYCE1-SIX6OS1 complex is retained through the unaffected alternative site (Fig. 6B).

SIX6OS1 Δ 10–21 retains SYCE1 binding in heterologous systems

Our biochemical and Y2H analyses concluded that SIX6OS1 Δ 10–21 would disrupt the first SYCE1-SIX6OS1 binding interface while retaining complex formation through the second interface. In support of this, we found that SIX6OS1 Δ 10–21 retained its ability to form intense colocalized foci with SYCE1 upon coexpression in COS7 cells (98% of the cells; Fig. 6C), similar to our previous observations for the SYCE1 POF mutation (Fig. 4A). Similarly, SIX6OS1 Δ 10–21 retained its ability to coimmunoprecipitate SYCE1 upon coexpression in HEK293 cells (Fig. 6D). Thus, localization and coimmunoprecipitation data from heterologous systems support our Y2H findings that the second SYCE1-SIX6OS1 binding interface is retained in SIX6OS1 Δ 10–21, mirroring the retention of only the second binding interface that is predicted for the 126–155 deletion of the SYCE1 c.375-2A>G NOA mutation (40).

SIX6OS1 Δ 10–21 leads to failure of SC assembly and murine infertility

Having established that the severe phenotype of the SYCE1 POF mutation likely results from the disruption of the second SYCE1-SIX6OS1 binding interface and its interaction with SYCE3, we wondered whether a similar phenotype would result from the sole disruption of the first SYCE1-SIX6OS1 binding interface. To test this, we generated mice harboring mutations of *Six6os1* alleles encoding internal in-frame deletions of amino acids 10 to 21 (equivalent numbering to the human protein) (fig. S8, A and B). While heterozygotes (designated *Six6os1* ^{Δ 10–21/*WT*}) were fertile, both male and female homozygotes (designated *Six6os1* ^{Δ 10–21/ Δ 10–21}) were infertile, similar to the SYCE1 POF mutation. In males, we observed reduced testis size (Fig. 7A) and a zygotene-like arrest similar to that observed in the *Six6os1* and *Syce1* knockouts (11, 16). The mutant spermatocytes were defective in synapsis and SC assembly, with reduced staining for SC proteins SYCP1 (Fig. 7B) and SYCE3 (Fig. 7C) and no staining for SYCE2-TEX12 (Fig. 7, F and G). In

Fig. 5. SYCE1core undergoes conformational change to form a 1:1 complex with SIX6OS1N. (A) Amylose pull-down following coexpression of MBP-SIX6OS1 1-75, 1-67, 1-75 Δ10-21, and free MBP with His-SYCE1core. (B) SDS-PAGE of the copurification of the SYCE1core-SIX6OS1N complex. Ni-NTA, Ni-nitrilotriacetic acid. (C) CD thermal denaturation recording the CD helical signature at 222 nm between 5° and 95°C, as % unfolded; estimated melting temperatures (T_m) are indicated. (D) SEC-MALS analysis. SYCE1core-SIX6OS1N (blue), SYCE1POF-SIX6OS1N (red) and full-length SYCE1-SIX6OS1N (black) are 1:1 complexes of 27, 37 (29 kDa for the degradation product complex), and 46 kDa, respectively (theoretical 1:1 to 27, 36, and 48 kDa), while MBP-SIX6OS1N (gray) is a 57-kDa monomer (theoretical, 53 kDa). SDS-PAGE of the SYCE1POF-SIX6OS1N sample is shown in Fig. 1C. (E) Schematic of the conformational change of the SYCE1core antiparallel dimer (yellow) into a 1:1 SYCE1core-SIX6OS1N complex (yellow-blue). (F and G) SEC-SAXS analysis. (F) SEC-SAXS P(r) interatomic distance distributions of SYCE1core-SIX6OS1N (blue), SYCE1POF-SIX6OS1N (red), and SYCE1core (yellow), revealing maximum dimensions (D_{max}) of 138, 180, and 186 Å, respectively. Their cross-sectional radii (R_c) are indicated (fig. S7D). (G) SAXS ab initio models of SYCE1core-SIX6OS1N (blue) and SYCE1core (yellow); averaged models were generated from 20 independent DAMMIF runs. Data for SYCE1core and full-length SYCE1 are reproduced from (28).

Fig. 6. SYCE1 undergoes multivalent interaction with SIX6OS1 in yeast, but SIX6OS1 $\Delta 10-21$ retains SYCE1 binding in heterologous systems. (A) Y2H analysis of interactions between SYCE1 and SIX6OS1 in which positive reactions are indicated by the growth of blue colonies. These data are representative of three repeats. (B) Schematic of the SYCE1-SIX6OS1 interaction based on the Y2H data in (A), with the two binding sites highlighted in red and green. The SYCE1 POF mutation blocks the second binding interface between SYCE1 177–305 and SIX6OS1 downstream sequence within region 1–262, whereas the SIX6OS1 $\Delta 10-21$ deletion blocks the first binding interface between SYCE1 core (25–179) and SIX6OS1N (1–67). (C) COS7 cells were transfected with mouse *Six6os1* $\Delta 10-21$ alone or in combination with mouse *Syce1*. SIX6OS1 $\Delta 10-21$ showed nuclear localization with some cytoplasmic signal and colocalized in cytoplasmic foci with SYCE1; the percentage of cells exhibiting colocalization is shown. Scale bars, 20 μ m. (D) Coimmunoprecipitation of SIX6OS1 $\Delta 10-21$ and Flag-SYCE1 from cotransfected HEK293T cells using anti-Myc or anti-EGFP antibodies, or mouse IgG as a negative control. SIX6OS1 $\Delta 10-21$ coimmunoprecipitated SYCE1, indicating that the second SYCE1 binding interface is retained. The untransfected lanes confirm the absence of SIX6OS1 $\Delta 10-21$ and SYCE1 in total protein extracts of untransfected 293T cells.

contrast with their complete absence in the SYCE1 POF mutation, we observed some residual staining for SYCE1 (Fig. 7D) and SIX6OS1 (Fig. 7E) even though the levels of transcription of *Six6os1* $\Delta 10-21$ appeared to be increased in the mutant testis (fig. S9 and table S1B). We detected γ -H2AX (fig. S10A) and DMC1/RAD51 foci (fig. S10, B and C) on aligned axial elements but no MLH1 foci (fig. S10D), indicating the proper induction of DSBs with their failed repair and absence of COs. Thus, SIX6OS1 $\Delta 10-21$ leads to infertility with a phenotype of failed DSB repair and SC assembly, similar to the SYCE1 POF mutation and those reported for disruption of structural components of the CE (10, 11, 16–18).

Thus, we conclude that both first and second SYCE1-SIX6OS1 binding interfaces are essential for SC assembly and meiotic progression. Furthermore, these findings explain how the sole disruption of individual SYCE1-SIX6OS1 binding interfaces by SYCE1 NOA (c.375-2A>G) and POF (c.613C>T) mutations result in the reported familial cases of human infertility.

DISCUSSION

The structural and functional integrity of the SC is contingent on the structure and assembly of its constituent protein components.

Here, we report that SC assembly depends on multivalent interactions between CE components SYCE1 and SIX6OS1 that are disrupted by infertility-associated mutations of SYCE1. The first binding interface is formed by the structural core of SYCE1 (SYCE1core; amino acids 25 to 179), which undergoes conformational change from an antiparallel homodimer to a 1:1 complex upon interaction with SIX6OS1's N terminus (SIX6OS1N; amino acids 1 to 67). The second binding interface is formed by downstream sequence within SIX6OS1 1–262 interacting directly with SYCE1 177–305. Through the generation of mice harboring an internal deletion of SIX6OS1's N terminus ($\Delta 10-21$) and the SYCE1 POF mutation (murine p. Gln243*), which specifically block the first and second binding interfaces, respectively, we find that integrity of both SYCE1-SIX6OS1 binding interfaces is essential for SC assembly and meiotic progression in vivo.

What is the structure of the SYCE1-SIX6OS1 complex? SEC-SAXS analysis revealed that the SYCE1core-SIX6OS1N 1:1 complex formed by the first binding interface has a length and cross-sectional radius of 138 and 11 Å, in comparison with 186 and 9 Å for the SYCE1core dimer. We previously reported a model for SYCE1core in which amino acids 52 to 179 form an antiparallel dimeric coiled coil containing a midline “kink”, with α helices of amino acids 25 to 50 packing

Fig. 7. Synapsis between homologs is disrupted in *Six6os1*^{Δ10-21/Δ10-21} spermatocytes. (A) Genetic deletion of amino acids 10 to 21 of SIX6OS1 led to a reduction of the testis size compared to the WT (mice of 3 months of age). (B) Double immunolabeling of WT pachytene and *Six6os1*^{Δ10-21/Δ10-21} zygotene-like spermatocytes with SYCP3 (red) and SYCP1 (green). AEs failed to synapse in *Six6os1*^{Δ10-21/Δ10-21} spermatocytes despite partial alignment, with reduced loading of SYCP1 along the AEs. (C to G) Double immunolabeling of spermatocyte spreads with SYCP3 (red) and all CE components (green). *Six6os1*^{Δ10-21/Δ10-21} zygotene-like spermatocytes showed reduced signals of (C) SYCE3, (D) SYCE1, and (E) SIX6OS1, and the absence of (F) SYCE2 and (G) TEX12 from the AEs. Scale bars, 10 μ m. Plots represent the quantification of fluorescence intensity levels in *Six6os1*^{Δ10-21/Δ10-21} zygotene-like and WT pachytene spermatocytes (B to E). Welch's *t* test analysis: ****P* < 0.0001. (H) Schematic of how the SYCE1 antiparallel dimer (yellow) undergoes conformational change upon interaction with SIX6OS1 (blue) to form a possible 1:1 complex through consecutive binding interfaces mediated by SYCE1core-SIX6OS1_N (site 1) and SYCE1 177–305 and downstream sequence within SIX6OS1 1–262 (site 2). The consequence of SYCE1 mutations associated with POF (c.613C>T) and NOA (c.375-2A>G) and SIX6OS1 Δ 10–21 on the integrity, predicted stoichiometry, and conformation of resultant SYCE1-SIX6OS1 complexes is illustrated. Photo credit (A): Laura Gómez-H, Instituto de Biología Celular y Molecular del Cáncer.

against this structural core (fig. S11A) (28). A maximum dimension of 138 Å for SYCE1core-SIX6OS1_N suggests a coiled-coil length of approximately 92 amino acids, given a helical rise of 1.5 Å per amino acid (44). This could be explained by the 52–179 region forming a helix-turn-helix structure through exaggeration of the kink to a full turn, which may combine with the α helix formed by amino acids 25 to 50 and an α helix from SIX6OS1_N to form a four-helical coiled coil, consistent with its 11-Å cross-sectional radius (fig. S11B). The second binding interface between SYCE1 177–305 and downstream sequence within SIX6OS1 1–262 suggests that SYCE1core-SIX6OS1_N likely adopts a parallel configuration to form a single SYCE1-

SIX6OS1 1:1 complex of consecutive first and second binding interfaces (Fig. 7H).

Our analysis of the SYCE1-SIX6OS1 complex reveals how the three reported clinical mutations of SYCE1 differentially affect its interaction with SIX6OS1. The SYCE1 NOA mutation c.197-2A>G is predicted to result in a truncated product of amino acids 1 to 65 (39), which would disrupt both binding sites and so likely abrogates SYCE1-SIX6OS1 complex formation and thus works as a null mutation. The SYCE1 NOA mutation c.375-2A>G is predicted to result in internal deletion of amino acids 126 to 155 (40), which would disrupt the first binding interface while retaining the second binding

interface, and so is likely to result in a conformationally altered 1:1 complex (Fig. 7H). In contrast, while $\Delta 10-21$ SIX6OS1 similarly disrupts the first binding interface and retains the second binding interface, the SYCE1 core remains unaffected and so is predicted to enable formation of a head-to-head 2:2 complex (Fig. 7H). The SYCE1 POF mutation c.613C>T generates a premature stop codon (p.Gln241*) that gives a truncated product of amino acids 1 to 240 (41), which we have demonstrated disrupts the second binding interface while retaining the first binding interface (Fig. 7H). Thus, the latter two infertility-associated mutations of SYCE1 specifically disrupt one SYCE1-SIX6OS1 interface while retaining the other, which combine with our mouse genetic studies to confirm that both interfaces are essential for the structural assembly of the SC and its function in meiosis.

What are the structural roles of SYCE1 and SYCE1-SIX6OS1 within the SC? Our analyses of *Syce1*^{POF/POF} and *Six6os1* ^{$\Delta 10-21/\Delta 10-21$} mouse strains revealed similar phenotypes with retention of some SYCP1 and SYCE3 recruitment to chromosome axes, with absence or substantial reduction of SYCE1 and SIX6OS1, and lack of recruitment of SYCE2-TEX12. This pattern suggests a hierarchical model of SC assembly in which SYCE1 and SYCE1-SIX6OS1 lie downstream of SYCP1 and SYCE3, and upstream of SYCE2-TEX12 (Fig. 1A), which is consistent with existing knockout data (10, 11, 15–18). The disruption of SYCE3 binding by the POF mutation suggests that its SYCE1-SIX6OS1 complex would be defective for SC recruitment, whereas the SYCE1-SYCE3 interaction, and hence SC recruitment, should be retained for the SYCE1-SIX6OS1 complex of the SIX6OS1 $\Delta 10-21$ internal deletion. This explains the greater severity of the CE loading defect in *Syce1*^{POF/POF} than *Six6os1* ^{$\Delta 10-21/\Delta 10-21$} , in which SYCE1 and SIX6OS1 staining was substantially reduced in the latter (83.77% of SYCE1 reduction, 0.12 ± 0.02 in the *Six6os1* ^{$\Delta 10-21/\Delta 10-21$} versus 0.73 ± 0.21 in the WT; 68.27% of SIX6OS1 reduction, 0.24 ± 0.02 in the *Six6os1* ^{$\Delta 10-21/\Delta 10-21$} versus 0.76 ± 0.15 in the WT) but completely absent in the former. Thus, we conclude that the first and second SYCE1-SIX6OS1 interfaces are essential for initiation of SC CE formation and likely function by stabilizing a local three-dimensional SC structure that mediates recruitment and self-assembly of SYCE2-TEX12 into fibers that mediate SC elongation along the chromosome axis. Furthermore, the SYCE1 POF mutation is likely worsened by its additional disruption of SYCE3 binding that removes the residual SYCE1-SIX6OS1 SC recruitment observed for the SIX6OS1 $\Delta 10-21$ internal deletion.

The existence of SYCE1 core as an isolated antiparallel homodimer and in a 1:1 complex with SIX6OS1_N raises the question of which is the biologically relevant conformation. It is important to highlight that the CD melting temperatures of SYCE1-SIX6OS1_N complexes and isolated SYCE1 dimers are very similar, ranging between 38° and 41°C. In contrast, highly stable SC components SYCE2-TEX12 and SYCP3 have melting temperatures of approximately 65°C (24, 29). Thus, the relatively low melting temperatures of SYCE1-SIX6OS1_N complexes and SYCE1 suggest that they may undergo conformational change in vivo, with each conformation functioning at different stages of meiosis and/or at different locations within the SC. Furthermore, our analysis of SYCE1 infertility-associated mutations and a targeted internal deletion of SIX6OS1 revealed at least four possible conformations of SYCE1 and SYCE1-SIX6OS1 complexes (Fig. 7H). Owing to the direct competition between SIX6OS1_N binding and SYCE1 core dimerization, these conformations could be achieved in the absence of mutations,

through alterations of protein levels, local concentrations, allosteric changes, and posttranslational modifications. Hence, alternative conformations of SYCE1 and SYCE1-SIX6OS1 are intriguing candidates for local structural heterogeneity and the propagation of signals along the length of the SC, which could function in roles such as crossover enforcement and interference. Thus, as we progress toward a full molecular understanding of the mammalian SC, the multivalent SYCE1-SIX6OS1 interactions described herein provide tantalizing possibilities for a dynamic role of SC structure in its enigmatic functions in the mechanics of meiosis.

MATERIALS AND METHODS

Recombinant protein expression and purification

Human SYCE1 sequences were cloned into pHAT4 and pMAT11 vectors (45) for bacterial expression as His- and His-MBP (Maltose-Binding Protein) fusions with TEV (Tobacco Etch Virus) cleavage sites for fusion protein removal. Human SIX6OS1 was cloned into pRSF-Duet1 vectors with a TEV-cleavable N-terminal MBP fusion for coexpression with SYCE1. Proteins were expressed in BL21(DE3) *Escherichia coli* cells (Novagen), in 2xYT (Yeast Extract Tryptone) media. Expression was induced with addition of 0.5 mM isopropyl- β -D-thiogalactopyranoside with the cells incubated at 25°C for 16 hours. Cells were lysed via sonication in 20 mM Tris (pH 8.0) and 500 mM KCl, followed by centrifugation. Supernatant was applied to an amylose (New England Biolabs) affinity chromatography column, followed by HiTrap Q HP (GE Healthcare) anion exchange chromatography. His- and His-MBP/MBP tags were removed by incubation with TEV protease at 4°C for 16 hours. The cleaved proteins were further purified by HiTrap Q HP (GE Healthcare) anion exchange chromatography followed by size exclusion chromatography (HiLoad 16/600 Superdex 200, GE Healthcare). The purified proteins/complexes were concentrated using Microsep Advance 3 kDa (PALL) centrifugal filter units and stored at –80°C. Protein samples were analyzed for purity using Coomassie-stained SDS-polyacrylamide gel electrophoresis. Protein molecular weights and extinction coefficients were calculated using ExPASy ProtParam (<http://web.expasy.org/protparam/>) with protein concentrations determined using a Cary 60 ultraviolet (UV) spectrophotometer (Agilent).

Circular dichroism

Far-UV CD spectra were collected using a Jasco J-810 spectropolarimeter (Institute for Cell and Molecular Biosciences, Newcastle University). Wavelength scans were recorded at 4°C from 260 to 185 nm at 0.2-nm intervals using a 0.2-mm path length quartz cuvette (Hellma). Protein samples were measured at 0.2 to 0.4 mg/ml in 10 mM Na₂HPO₄ (pH 7.5) and 150 mM NaF. Nine measurements were taken for each sample, averaged, buffer-corrected and converted to mean residue ellipticity (MRE) ($[\theta]$) ($\times 1000 \text{ deg} \cdot \text{cm}^2 \cdot \text{dmol}^{-1}$ per residue). Spectral deconvolutions were carried out using the Dichroweb CDSSTR algorithm (<http://dichroweb.cryst.bbk.ac.uk>). CD thermal melts were recorded at 222 nm between 5° and 95°C, at intervals of 0.5°C with a 1°C/min ramping rate. Protein samples were measured at 0.1 mg/ml in 20 mM Tris (pH 8.0), 150 mM KCl, and 2 mM dithiothreitol (DTT), using a 1-mm path length quartz cuvette (Hellma). The data were plotted as % unfolded after conversion to MRE ($[\theta]_{222,x} - [\theta]_{222,5}) / ([\theta]_{222,95} - [\theta]_{222,5})$. The melting temperature was determined as the temperature at which the proteins are 50% unfolded.

Size exclusion chromatography multiangle light scattering

SEC-MALS analysis of protein samples was carried out at concentrations of 5 to 20 mg/ml in 20 mM Tris (pH 8.0), 150 mM KCl, and 2 mM DTT. Samples were loaded onto a Superdex 200 Increase 10/300 GL (GE Healthcare) column at 0.5 ml/min using an ÄKTA Pure (GE Healthcare) system. The eluate was fed into a DAWN HELEOS II MALS detector (Wyatt Technology), followed by an Optilab T-rEX differential refractometer (Wyatt Technology). SEC-MALS data were collected and analyzed using ASTRA 6 software (Wyatt Technology), using Zimm plot extrapolation with a 0.185 ml/g dn/dc value to determine absolute protein molecular weights.

Size exclusion chromatography small-angle x-ray scattering

SEC-SAXS experiments were carried out on beamline B21 at the Diamond Light Source synchrotron facility (Oxfordshire, UK). Protein samples at concentrations 6 to 20 mg/ml were loaded onto a Superdex 200 Increase 10/300 GL size exclusion chromatography column (GE Healthcare) in 20 mM Tris (pH 8.0) and 150 mM KCl at 0.5 ml/min using an Agilent 1200 high-performance liquid chromatography system. The eluate was fed through the experimental cell, with SAXS data recorded at 12.4 keV, in 3.0-s frames with a detector distance of 4.014 m. ScÅtter 3.0 (www.bioisis.net) was used to subtract and average the frames and carry out the Guinier analysis for the R_g and cross-sectional R_g (R_c). $P(r)$ distributions were fitted using PRIMUS. Ab initio modeling was performed using DAMMIF (46) imposing P1 symmetry. Twenty independent runs were averaged. The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC was used to generate images of the SAXS ab initio models.

Yeast two-hybrid

Constructs of human SYCE1 and SIX6OS1 were cloned into pGBKT7 and pGADT7 vectors (Clontech). Y2H experiments were carried out using the Matchmaker Gold system (Clontech) according to the manufacturer's guidelines. Y187 yeast strain was transformed with pGBKT7 vectors, while the Y2H gold strain was transformed with pGADT7 vectors. Yeast transformations were carried out using standard lithium acetate methods. Mating of the two strains was carried out in 0.5 ml 2× YPDA (Yeast Peptone Dextrose Adenine) at 30°C, 40 rpm, by mixing respective colonies. After 24 hours, the cultures were centrifuged and pellets were resuspended in 0.5×YPDA. These were then plated onto SD/-Trp/-Leu to select for mated colonies and onto SD/-Trp/-Leu/-Ade/-His with X-α-gal to detect mated colonies through ADE1, HIS3, and MEL1 reporter gene activation. Plates were then incubated for 5 days at 30°C.

Production of CRISPR-Cas9-edited mice

For developing the *Syce1*^{POF/POF} model, *Syce1*-single-guide RNA (sgRNA) 5'-TGACTTCTTCCACACTATC-3' targeting the intron 10 was predicted at https://eu.idtdna.com/site/order/designtool/index/CRISPR_SEQUENCE. This crRNA (CRISPR RNA), the tracrRNA (trans-activating CRISPR RNA), and the ssODN (single-stranded donor oligonucleotides) (5'-GGGACTCTTCTCCG-AAGCCATGAGGAGCTGCAGCAATGTAAGATGCAGGGT-GGGGCAGGAGGAGGAAATGTCTAGCACTGACTTCTTTC-CACACCCCGAGGTAGATCTTCAAGGATGAGAA-CAAGAAAGCTGAGG

AGTTCCTAGAGGCTGCAGCTCAGCAGCAGCAGCAGCT-GCAGCAGAGGTGCCACCACTACAG-3') were produced by chemical synthesis at IDT. The ssODN contains the mutated base

(C>T, p.Gln241*) and the peptidyl-glycine α-amidating monooxygenase (PAM) was mutated by substituting it by the human intron sequence (ACTATCAG > CCCCCAG). The crRNA and tracrRNA were annealed to obtain the mature sgRNA. A mixture containing the sgRNAs, recombinant Cas9 protein (IDT), and the ssODN [Cas9 (30 ng/μl), annealed sgRNA (20 ng/μl each), and ssODN (10 ng/μl)] were microinjected into B6/CBA F2 zygotes (hybrids between strains C57BL/6 J and CBA/J) (47) at the Transgenic Facility of the University of Salamanca. Edited founders were identified by polymerase chain reaction (PCR) amplification (Taq polymerase, NZYTech) with primers flanking the exon 11 (primer F 5'-CTGTAGAGAACT-GATGAAAGT-3' and R 5'-CAAGAAAATATGAAGAGACATAC-3') producing an amplicon of 398 base pairs (bp) for both edited and WT alleles, and either direct sequenced or subcloned into pBlueScript (Stratagene) followed by Sanger sequencing, selecting the point mutation in the targeted region of *Syce1* (fig. S2). For generating the *Six6os1*^{Δ10-21/Δ10-21} (named as *Six6os1*^{Δ/Δ}), *Six6os1*-crRNA G68 5'-ATCTGTTTGTCTGAGTTTGGAC-3' and *Six6os1*-crRNA G75 5'-TACTTATGTCTTGCTCATAC-3' targeting exons 2 and 3 and the ssODN (5'-GTTCTTACTTTATGTATGCTCTTTTATATAT-GGCTTCTGAAAGTTTTATTTATTTTATTTTACA-CAGTGTCCCAAGATGAATGATAATCTGTTTGT-CAGTTTGTCAAGACATAAGTATTTAAAGAAGATACGAT-TCAAAGAATTAATAGTAAGTAGTTTGCATGAAATA-AATATTTTAGTCTTTTGGTTTATCTTATATAGCA-3') were predicted, produced, and microinjected, as previously described. Edited founders with the predicted deletion were identified through PCR using primers flanking this region (primer F 5'-CACTTACATTTTCTTTTAAGAATGC-3' and R 5'-CCCCTCTCATACATACAAGTTGC-3'). The Δ10-21 allele was 285 bp long versus 413 bp of the WT allele (fig. S8, A and B). The founders were crossed with WT C57BL/6 J to eliminate possible unwanted off-targets. Heterozygous mice were resequenced and crossed to give rise to edited homozygous. Genotyping was performed by analysis of the PCR products of genomic DNA with primers F and R.

Histology

For histological analysis of ovaries, after the necropsy of the mice, their ovaries were removed and fixed in formol 10%. They were processed into serial paraffin sections and stained with hematoxylin and eosin. The samples were analyzed using a microscope OLYMPUS BX51, and images were taken with a digital camera OLYMPUS DP70.

Immunocytology

Testes were detunicated and processed for spreading using a conventional "dry-down" technique. Oocytes from fetal ovaries (E17.5 embryos) were digested with collagenase, incubated in hypotonic buffer, disaggregated, and fixed in paraformaldehyde. Both meiotic preparations were incubated with the following primary antibodies for immunofluorescence (IF): rabbit αSIX6OS1 R1 and R2 [1:100, Proteogenix (11)], rabbit αSYCE1 17406-1-AP (1:50, Proteintech), guinea pig αSYCE1 (1:100, provided by C. Höög), mouse αSYCP3 immunoglobulin G (IgG) sc-74569 (1:1000, Santa Cruz Biotechnology), rabbit serum αSYCP3 K921 (1:500), rabbit αSYCP1 IgG ab15090 (1:200), guinea pig αSYCE3 (1:20, provided by R. Benavente), guinea pig αSYCE2 (1:100, provided by C. Höög), rabbit αTEX12 IgG (1:100, provided by R. Benavente), rabbit anti-γ-H2AX (ser139) IgG #07-164 (1:200) (Millipore), mouse αMLH1 51-1327GR (1:5, BD Biosciences), rabbit αRAD51 PC130 (1:50, Calbiochem), and rabbit

α DMC1 R1 and R2 (1:500, Proteogenix). The secondary antibodies used were goat Alexa 555 α -mouse A-32727, goat Alexa 488 α -mouse A-11001, donkey Alexa 555 α -rabbit A-31572 (1:200, Thermo Fisher Scientific), goat Alexa 488-Fab α -rabbit 111-547-003, and donkey fluorescein isothiocyanate α -guinea pig 706-095-148 (1:100, Jackson ImmunoResearch). Slides were visualized at room temperature using a microscope (Axioplan 2; Carl Zeiss Inc.) with 63 $\times$ objectives with an aperture of 1.4 (Carl Zeiss Inc.). Images were taken with a digital camera (ORCA-ER; Hamamatsu) and processed with OPENLAB 4.0.3 and Photoshop (Adobe). Quantification of fluorescence signals was performed using ImageJ software.

Cell lines and transfections

HEK293T and COS7 cell lines were obtained from the American Type Culture Collection (ATCC). Cell lines were tested for mycoplasma contamination (Mycoplasma PCR ELISA, Sigma-Aldrich). They were transfected with Jetpei (PolyPlus) according to the manufacturer's protocol.

Immunoprecipitation and Western blotting

HEK293T cells were transiently transfected, and whole-cell extracts were prepared and cleared with protein G Sepharose beads (GE Healthcare) for 1 hour. The antibody was added for 2 hours, and immunocomplexes were isolated by adsorption to protein G Sepharose beads overnight. After washing, the proteins were eluted from the beads with 2 $\times$ SDS gel-loading buffer 100 mM Tris-HCl (pH 7), 4% SDS, 0.2% bromophenol blue, 200 mM β -mercaptoethanol, and 20% glycerol and loaded onto reducing polyacrylamide SDS gels. The proteins were detected by Western blotting with the indicated antibodies. Immunoprecipitations were performed using mouse α -Flag IgG (5 μ g; F1804, Sigma-Aldrich), mouse α -green fluorescent protein (α -GFP) IgG (4 μ g; CSB-MA000051M0m, Cusabio), mouse α -Myc obtained from hybridoma cell myc-1-9E10.2 ATCC (4 μ g), and ChromPure mouse IgG (5 μ g/1 mg protein; 015-000-003). Primary antibodies used for Western blotting were rabbit α -Flag IgG (1:2000; F7425 Sigma-Aldrich), goat α -GFP IgG (sc-5385, Santa Cruz Biotechnology) (1:3000), and rabbit α -Myc Tag IgG (1:3000; #06-549, Millipore). Secondary horseradish peroxidase-conjugated α -mouse (715-035-150, Jackson ImmunoResearch), α -rabbit (711-035-152, Jackson ImmunoResearch), or α -goat (705-035-147, Jackson ImmunoResearch) antibodies were used at 1:5000 dilution. Antibodies were detected by using Immobilon Western Chemiluminescent HRP Substrate from Millipore. Both *Syce1*^{POF} and *Six6os1* Δ 10-21 complementary DNAs (cDNAs) used for IF and coimmunoprecipitation experiments were reverse transcription PCR-amplified (the primers used for it were *Syce1* 5'-GAGCAGTATGGC-CACCAGACC-3' and *Syce* AS 5'-GAGGAGGGTATTAGGTCCTGC-3'; *Six6os1* 5'-AGTGTCGAAGATGAATGATAATCTG-3' and *Six6os1* AS 5'-GTTCAAAAATAATAACTCAAAAAAAC-3') from total RNA extracted from *Syce1*^{POF/POF} and *Six6os1* ^{Δ 10-21/ Δ 10-21} mice, respectively. PCR-amplified fragments were cloned in pcDNA3-based mammalian expression vectors with different tags (enhanced GFP or Flag) and verified by Sanger sequencing.

Quantitative PCR

Total RNA was isolated from testis of WT and mutant mice. To analyze the expression of *Syce1* and *Six6os1* mRNAs, equal amounts of cDNA were synthesized using SuperScript II Reverse Transcriptase (Invitrogen, Life Technologies) and Oligo (dT). Quantitative

PCR (qPCR) was performed using FastStart Universal SYBR Green Master Mix (ROX) (Roche) and specific forward and reverse primers: qSYCE1_F 5'-GGACATGGTAAAAAGTTGCAG-3' and qSYCE1_R 5'-CAGTTCCTTCTGCAGGTTGTC-3' for *Syce1*, and qSIX6OS1_F 5'-GCTGAATGTGGAGATAAAGAG-3' and qSIX6OS1_R 5'-AGGAGTTTCAGGAGTTTGAGG-3' for *Six6os1*. All qPCR reactions were performed at 95°C for 10 min and then 40 cycles of 95°C for 15 s and 62°C for 1 min on the iQ5 Thermal Cycler (Bio-Rad). β -Actin was amplified as a housekeeping gene with the primers q β -actin_F 5'-GGCACCACACCTTCTACAATG-3' and q β -actin_R 5'-GTG-GTGGTGAAGCTGTAGCC-3'.

Statistics

To compare counts between genotypes, we used the Welch's *t* test (unequal variances *t* test), which was appropriate as the count data were not highly skewed (i.e., were reasonably approximated by a normal distribution) and, in most cases, showed unequal variance. We applied a two-sided test in all the cases. Asterisks denote statistical significance: **P* < 0.01, ***P* < 0.001, and ****P* < 0.0001.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/6/36/eabb1660/DC1>

[View/request a protocol for this paper from Bio-protocol.](#)

REFERENCES AND NOTES

1. N. Hunter, Meiotic recombination: The essence of heredity. *Cold Spring Harb. Perspect. Biol.* **7**, a016618 (2015).
2. D. Zickler, N. Kleckner, Recombination, pairing, and synapsis of homologs during meiosis. *Cold Spring Harb. Perspect. Biol.* **7**, (2015).
3. J. Loidl, Conservation and variability of meiosis across the eukaryotes. *Annu. Rev. Genet.* **50**, 293–316 (2016).
4. F. Baudat, Y. Imai, B. de Massy, Meiotic recombination in mammals: Localization and regulation. *Nat. Rev. Genet.* **14**, 794–806 (2013).
5. C. K. Cahoon, R. S. Hawley, Regulating the construction and demolition of the synaptonemal complex. *Nat. Struct. Mol. Biol.* **23**, 369–377 (2016).
6. M. Westergaard, D. von Wettstein, The synaptonemal complex. *Annu. Rev. Genet.* **6**, 71–110 (1972).
7. K. Schmekel, B. Daneholt, The central region of the synaptonemal complex revealed in three dimensions. *Trends Cell Biol.* **5**, 239–242 (1995).
8. R. L. Meuwissen, H. H. Offenberg, A. J. Dietrich, A. Riesewijk, M. van Iersel, C. Heyting, A coiled-coil related protein specific for synapsed regions of meiotic prophase chromosomes. *EMBO J.* **11**, 5091–5100 (1992).
9. Y. Costa, R. Speed, R. Öllinger, M. Alsheimer, C. A. Semple, P. Gautier, K. Maratou, I. Novak, C. Höög, R. Benavente, H. J. Cooke, Two novel proteins recruited by synaptonemal complex protein 1 (SYCP1) are at the centre of meiosis. *J. Cell Sci.* **118**, 2755–2762 (2005).
10. S. Schramm, J. Fraune, R. Naumann, A. Hernandez-Hernandez, C. Höög, H. J. Cooke, M. Alsheimer, R. Benavente, A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility. *PLoS Genet.* **7**, e1002088 (2011).
11. L. Gomez-H, N. Felipe-Medina, M. Sánchez-Martín, O. R. Davies, I. Ramos, I. García-Turiñón, D. G. de Rooij, I. Dereli, A. Tóth, J. L. Barbero, R. Benavente, E. Llano, A. M. Pendas, C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nat. Commun.* **7**, 13298 (2016).
12. G. Hamer, K. Gell, A. Kouznetsova, I. Novak, R. Benavente, C. Höög, Characterization of a novel meiosis-specific protein within the central element of the synaptonemal complex. *J. Cell Sci.* **119**, 4025–4032 (2006).
13. H. H. Offenberg, J. A. Schalk, R. L. Meuwissen, M. van Aalderen, H. A. Kester, A. J. Dietrich, C. Heyting, SCP2: A major protein component of the axial elements of synaptonemal complexes of the rat. *Nucleic Acids Res.* **26**, 2572–2579 (1998).
14. J. H. Lammers, H. H. Offenberg, M. van Aalderen, A. C. Vink, A. J. Dietrich, C. Heyting, The gene encoding a major component of the lateral elements of synaptonemal complexes of the rat is related to X-linked lymphocyte-regulated genes. *Mol. Cell. Biol.* **14**, 1137–1146 (1994).

15. F. A. T. de Vries, E. de Boer, M. van den Bosch, W. M. Baarends, M. Ooms, L. Yuan, J.-G. Liu, A. A. van Zeeland, C. Heyting, A. Pastink, Mouse *Sycp1* functions in synaptonemal complex assembly, meiotic recombination, and XY body formation. *Genes Dev.* **19**, 1376–1389 (2005).
16. E. Bolcun-Filas, E. Hall, R. Speed, M. Taggart, C. Grey, B. de Massy, R. Benavente, H. J. Cooke, Mutation of the mouse *Syce1* gene disrupts synapsis and suggests a link between synaptonemal complex structural components and DNA repair. *PLoS Genet.* **5**, e1000393 (2009).
17. E. Bolcun-Filas, Y. Costa, R. Speed, M. Taggart, R. Benavente, D. G. de Rooij, H. J. Cooke, SYCE2 is required for synaptonemal complex assembly, double strand break repair, and homologous recombination. *J. Cell Biol.* **176**, 741–747 (2007).
18. G. Hamer, H. Wang, E. Bolcun-Filas, H. J. Cooke, R. Benavente, C. Hoog, Progression of meiotic recombination requires structural maturation of the central element of the synaptonemal complex. *J. Cell Sci.* **121**, 2445–2451 (2008).
19. L. Yuan, J. G. Liu, J. Zhao, E. Brundell, B. Daneholt, C. Höög, The murine SCP3 gene is required for synaptonemal complex assembly, chromosome synapsis, and male fertility. *Mol. Cell* **5**, 73–83 (2000).
20. F. Yang, R. D. L. Fuente, N. A. Leu, C. Baumann, K. J. McLaughlin, P. J. Wang, Mouse SYCP2 is required for synaptonemal complex assembly and chromosomal synapsis during male meiosis. *J. Cell Biol.* **173**, 497–507 (2006).
21. L. Yuan, J. G. Liu, M. R. Hoja, J. Wilbertz, K. Nordqvist, C. Höög, Female germ cell aneuploidy and embryo death in mice lacking the meiosis-specific protein SCP3. *Science* **296**, 1115–1118 (2002).
22. J. M. Dunce, O. M. Dunne, M. Ratcliff, C. Millán, S. Madgwick, I. Usón, O. R. Davies, Structural basis of meiotic chromosome synapsis through SYCP1 self-assembly. *Nat. Struct. Mol. Biol.* **25**, 557–569 (2018).
23. K. Schucker, T. Holm, C. Franke, M. Sauer, R. Benavente, Elucidation of synaptonemal complex organization by super-resolution imaging with isotropic resolution. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 2029–2033 (2015).
24. J. L. Syrjänen, L. Pellegrini, O. R. Davies, A molecular model for the role of SYCP3 in meiotic chromosome organisation. *eLife* **3**, (2014).
25. J. L. Syrjänen, I. Heller, A. Candelli, O. R. Davies, E. J. G. Peterman, G. J. L. Wuite, L. Pellegrini, Single-molecule observation of DNA compaction by meiotic protein SYCP3. *eLife* **6**, e22582 (2017).
26. J. Lu, Y. Gu, J. Feng, W. Zhou, X. Yang, Y. Shen, Structural insight into the central element assembly of the synaptonemal complex. *Sci. Rep.* **4**, 7059 (2014).
27. O. M. Dunne, O. R. Davies, A molecular model for self-assembly of the synaptonemal complex protein SYCE3. *J. Biol. Chem.* **294**, 9260–9275 (2019).
28. O. M. Dunne, O. R. Davies, Molecular structure of human synaptonemal complex protein SYCE1. *Chromosoma* **128**, 223–236 (2019).
29. O. R. Davies, J. D. Maman, L. Pellegrini, Structural analysis of the human SYCE2–TEX12 complex provides molecular insights into synaptonemal complex assembly. *Open Biol.* **2**, 120099 (2012).
30. A. Kouznetsova, R. Benavente, A. Pastink, C. Hoog, Meiosis in mice without a synaptonemal complex. *PLoS ONE* **6**, e28255 (2011).
31. H. F. Horn, D. I. Kim, G. D. Wright, E. S. M. Wong, C. L. Stewart, B. Burke, K. J. Roux, A mammalian KASH domain protein coupling meiotic chromosomes to the cytoskeleton. *J. Cell Biol.* **202**, 1023–1039 (2013).
32. X. Ding, R. Xu, J. Yu, T. Xu, Y. Zhuang, M. Han, SUN1 is required for telomere attachment to nuclear envelope and gametogenesis in mice. *Dev. Cell* **12**, 863–872 (2007).
33. P. J. Romanienko, R. D. Camerini-Otero, The mouse Spo11 gene is required for meiotic chromosome synapsis. *Mol. Cell* **6**, 975–987 (2000).
34. K. Yoshida, G. Kondoh, Y. Matsuda, T. Habu, Y. Nishimune, T. Morita, The mouse RecA-like gene Dmc1 is required for homologous chromosome synapsis during meiosis. *Mol. Cell* **1**, 707–718 (1998).
35. M. A. Handel, J. C. Schimenti, Genetics of mammalian meiosis: Regulation, dynamics and impact on fertility. *Nat. Rev. Genet.* **11**, 124–136 (2010).
36. A. Geisinger, R. Benavente, Mutations in genes coding for synaptonemal complex proteins and their impact on human fertility. *Cytogenet. Genome Res.* **150**, 77–85 (2016).
37. P. Laisue, Aetiological coding sequence variants in non-syndromic premature ovarian failure: From genetic linkage analysis to next generation sequencing. *Mol. Cell. Endocrinol.* **411**, 243–257 (2015).
38. S. Caburet, V. A. Arboleda, E. Llano, P. A. Overbeek, J. L. Barbero, K. Oka, W. Harrison, D. Vaiman, Z. Ben-Neriah, I. García-Tuñón, M. Fellous, A. M. Pendas, R. A. Veitia, E. Vilain, Mutant cohesin in premature ovarian failure. *N. Engl. J. Med.* **370**, 943–949 (2014).
39. E. Maor-Sagie, Y. Cinnamon, B. Yaacov, A. Shaag, H. Goldsmidt, S. Zenvirt, N. Laufer, C. Richler, A. Frumkin, Deleterious mutation in SYCE1 is associated with non-obstructive azoospermia. *J. Assist. Reprod. Genet.* **32**, 887–891 (2015).
40. M. Pashaei, M. M. Rahimi Bidgoli, D. Zare-Abdollahi, H. Najmabadi, R. Haji-Seyed-Javadi, F. Fatehi, A. Alavi, The second mutation of SYCE1 gene associated with autosomal recessive nonobstructive azoospermia. *J. Assist. Reprod. Genet.* **37**, 451–458 (2020).
41. L. de Vries, D. M. Behar, P. Smirni-Yosef, I. Lagovsky, S. Tzur, L. Basel-Vanagaite, Exome sequencing reveals SYCE1 mutation associated with autosomal recessive primary ovarian insufficiency. *J. Clin. Endocrinol. Metab.* **99**, E2129–E2132 (2014).
42. E. P. Mimitou, L. S. Symington, Nucleases and helicases take center stage in homologous recombination. *Trends Biochem. Sci.* **34**, 264–272 (2009).
43. S. K. Mahadevaiah, J. M. A. Turner, F. Baudat, E. P. Rogakou, P. de Boer, J. Blanco-Rodriguez, M. Jasin, S. Keeney, W. M. Bonner, P. S. Burgoyne, Recombinational DNA double-strand breaks in mice precede synapsis. *Nat. Genet.* **27**, 271–276 (2001).
44. A. N. Lupaş, M. Gruber, The structure of a-Helical coiled coils. *Adv. Protein Chem.* **70**, 37–78 (2005).
45. J. Peranen, M. Rikonen, M. Hyvönen, L. Kaariainen, T7 vectors with modified T7lac promoter for expression of proteins in *Escherichia coli*. *Anal. Biochem.* **236**, 371–373 (1996).
46. D. Franke, D. I. Svergun, DAMMIF, a program for rapid ab-initio shape determination in small-angle scattering. *J. Appl. Cryst.* **42**, 342–346 (2009).
47. P. Singh, J. C. Schimenti, E. Bolcun-Filas, A mouse geneticist's practical guide to CRISPR applications. *Genetics* **199**, 1–15 (2015).

Acknowledgments: We thank Diamond Light Source and the staff of beamline B21 (proposals sm15836, sm21777, and sm23510). We thank H. Waller for assistance with CD data collection.

Funding: O.R.D. is a Sir Henry Dale Fellow jointly funded by the Wellcome Trust and Royal Society (grant number 104158/Z/14/Z). This work was supported by MINECO (BFU2017-89408-R) and by Junta de Castilla y León (CS1239P18). F.S.-S., L.G.-H., and N.F.-M. are supported by European Social Fund/JCyL grants (EDU/556/2019, EDU/1083/2013, and EDU/310/2015). CIC-IBMC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia cofunded by the Castilla-León autonomous government and the European Regional Development Fund (CLC-2017-01). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. **Ethics statement:** Mice were housed in a temperature-controlled facility (specific pathogen free) using individually ventilated cages, standard diet, and a 12-hour light/dark cycle, according to European Union laws at the "Servicio de Experimentación Animal, SEA." Mouse protocols were approved by the Ethics Committee for Animal Experimentation of the University of Salamanca (USAL). We made every effort to minimize suffering and to improve animal welfare. Blinded experiments were not possible since the phenotype was obvious between WT and mutant mice for all of the experimental procedures used. No randomization methods were applied since the animals were not divided in groups or treatments. The minimum size used for each analysis was two animals per genotype.

Author contributions: F.S.-S., L.G.-H., O.M.D., N.F.-M., C.G.-P., M.S.-M., and O.R.D. performed experiments. O.R.D. and A.M.P. designed experiments, analyzed data, and wrote the manuscript. A.M.P., E.L., and O.R.D. supervised and designed the work. **Competing financial interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 4 February 2020

Accepted 17 July 2020

Published 2 September 2020

10.1126/sciadv.abb1660

Citation: F. Sánchez-Sáez, L. Gómez-H, O. M. Dunne, C. Gallego-Páramo, N. Felipe-Medina, M. Sánchez-Martín, E. Llano, A. M. Pendas, O. R. Davies, Meiotic chromosome synapsis depends on multivalent SYCE1-SIX6OS1 interactions that are disrupted in cases of human infertility. *Sci. Adv.* **6**, eabb1660 (2020).