



Phycomyces MADB interacts with MADA to form the primary photoreceptor complex for fungal phototropism

Catalina Sanz^{a,1}, Julio Rodríguez-Romero^{b,1}, Alexander Idnurm^{c,d}, John M. Christie^e, Joseph Heitman^c, Luis M. Corrochano^{b,2}, and Arturo P. Eslava^{a,2}

^aArea de Genética, Departamento de Microbiología y Genética and Centro Hispano Luso de Investigaciones Agrarias, Universidad de Salamanca, E-37007 Salamanca, Spain; ^bDepartamento de Genética, Facultad de Biología, Universidad de Sevilla, Apartado 1095, E-41080 Sevilla, Spain; ^cDepartment of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC 27710; ^dDivision of Cell Biology and Biophysics, School of Biological Sciences, University of Missouri-Kansas City, Kansas City, MO 64110; and ^ePlant Science Group, Division of Molecular and Cellular Biology, Faculty of Biomedical and Life Sciences, University of Glasgow, Glasgow G12 8QQ, United Kingdom

Edited by Winslow R. Briggs, Carnegie Institution of Washington, Stanford, CA, and approved March 13, 2009 (received for review January 27, 2009)

The fungus *Phycomyces blakesleeanus* reacts to environmental signals, including light, gravity, touch, and the presence of nearby objects, by changing the speed and direction of growth of its fruiting body (sporangiophore). Phototropism, growth toward light, shares many features in fungi and plants but the molecular mechanisms remain to be fully elucidated. *Phycomyces* mutants with altered phototropism were isolated ≈ 40 years ago and found to have mutations in the *mad* genes. All of the responses to light in *Phycomyces* require the products of the *madA* and *madB* genes. We showed that *madA* encodes a protein similar to the *Neurospora* blue-light photoreceptor, zinc-finger protein WC-1. We show here that *madB* encodes a protein similar to the *Neurospora* zinc-finger protein WC-2. MADA and MADB interact to form a complex in yeast 2-hybrid assays and when coexpressed in *E. coli*, providing evidence that phototropism and other responses to light are mediated by a photoresponsive transcription factor complex. The *Phycomyces* genome contains 3 genes similar to *wc-1*, and 4 genes similar to *wc-2*, many of which are regulated by light in a *madA* or *madB* dependent manner. We did not detect any interactions between additional WC proteins in yeast 2-hybrid assays, which suggest that MADA and MADB form the major photoreceptor complex in *Phycomyces*. However, the presence of multiple *wc* genes in *Phycomyces* may enable perception across a broad range of light intensities, and may provide specialized photoreceptors for distinct photoresponses.

blue light | LOV domain | White Collar protein | zinc finger | gene duplication

Organisms sense and interact with the surrounding environment to increase their probability of survival. Fungi respond to many environmental signals to modify their patterns of growth and behavior (1). Light, particularly blue light, serves as a signal to regulate fungal development and behavior, presumably for the optimization of spore production and dispersal (2). In addition, blue light activates metabolic pathways and directs the growth of fungal structures (3, 4).

The zygomycete fungus *Phycomyces blakesleeanus* has served as a model organism to investigate the responses of fungi to light (3, 5). Use of *Phycomyces* in sensory transduction research was promoted by the Nobel laureate Max Delbrück in the 1950s (6). Blue light regulates several aspects of *Phycomyces* biology: it regulates the development of fruiting bodies (sporangiophores), stimulates the biosynthesis of beta-carotene, and modifies the direction (phototropism) and speed of growth of the sporangiophores (3). In addition, the *Phycomyces* sporangiophore can change the direction of growth after sensing other environmental signals, like gravity, wind, touch, and the presence of nearby objects, making this unicellular structure a unique experimental object (3). Much of the attention in *Phycomyces* research has

focused on its responses to light. *Phycomyces* responds to a wide interval of light intensities extending 10 orders of magnitude. This remarkable sensory dexterity approximates that of the human eye and is achieved through the action of 2 photosystems optimized to operate at different light intensities (7).

A genetic screen for phototropic mutants, conducted in Delbrück's lab, allowed the isolation and characterization of *mad* mutants, and the first outline of the sensory transduction pathway for *Phycomyces* (8). The discovery of additional *mad* mutants and detailed genetic characterization led to the identification of 10 unlinked *mad* genes, *madA* through *madJ* (9, 10). Mutants of the *madA* and *madB* genes are defective in phototropism and other light responses suggesting that the corresponding gene products play key roles in *Phycomyces* photobiology (3).

Most of our understanding of fungal photobiology comes from studies with the ascomycete fungus *Neurospora crassa*. Mutations in the *wc-1* or *wc-2* genes disrupt all of the responses of *Neurospora* to blue light (4, 11). The WC-1 protein contains a zinc-finger, a chromophore-binding domain (named LOV), and PAS domains for protein-protein interactions (12). The LOV domain binds the flavin FAD, allowing WC-1 to act as a photoreceptor (13, 14). LOV was initially identified in phototropins, plant blue light photoreceptors for phototropism (15), and the structure of the LOV domain in a small *Neurospora* photoreceptor, VVD, has been determined (16). The WC-2 protein contains a zinc-finger and 1 PAS domain (17), and interacts with WC-1 to form a complex that binds to the promoters of light-inducible genes, presumably to activate their transcription (13, 18, 19). WC proteins are required for the responses to blue light in the basidiomycete fungi *Cryptococcus neoformans* (20, 21) and *Coprinus cinereus* (22), and 3 *wc-1* genes have been described in the zygomycetes *Rhizopus oryzae* and *Mucor circinelloides* (23, 24). A *Mucor* WC-1 protein is modified by ubiquitylation, presumably to regulate its activity (25). Red- and blue-light photoreceptors regulate development and secondary metabolism in the ascomycete fungus *Aspergillus nidulans* (26–28). Protein complexes containing photoreceptors or

Author contributions: C.S., J.R.-R., A.I., J.M.C., J.H., L.M.C., and A.P.E. designed research; C.S., J.R.-R., and A.I. performed research; C.S., J.R.-R., A.I., J.M.C., J.H., L.M.C., and A.P.E. analyzed data; and C.S., J.R.-R., A.I., J.M.C., J.H., L.M.C., and A.P.E. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The sequence reported in this paper has been deposited in the EMBL Nucleotide Sequence Database (FM178798 *wcoB*, FM178799 *madB*, FM178800 *wctB*, FM179475 *wctC*, FM178801 *wctD*).

¹C.S. and J.R.-R. contributed equally to this work.

²To whom correspondence may be addressed. E-mail: eslava@usal.es or corrochano@us.es.

This article contains supporting information online at www.pnas.org/cgi/content/full/0900879106/DCSupplemental.




Fig. 1. *Phycomyces* *wc* genes and proteins. (A) Genomic structure of *Phycomyces* *wc* genes including flanking genes with their putative identities. Exons, introns, and direction of transcription are indicated. (B) WC proteins in the *Phycomyces* genome and their domains. (C) Phylogenies of the WC proteins. Phylogenetic trees of the LOV domains from fungal WC-1 proteins and photoreceptor proteins from plants and bacteria, and from WC-2 proteins (for accession numbers see [SI Text](#)). Bars represent the number of substitutions per site. The *wctA* gene and the WCTA protein are indicated as *madB* and MADB, respectively.

transcriptional regulators participate in *Aspergillus* photobiology (27, 29). The existence of proteins similar to WC-1 and WC-2 in ascomycete, basidiomycete, and zygomycete fungi led to the proposal that the White Collar Complex arose early in fungal evolution as a photoreceptive transcription factor (1, 30, 31).

In *Phycomyces*, however, the molecular nature of the photoreceptor or the proteins involved in light-signal transduction remained unknown for many decades. In addition, none of the *mad* genes had been identified despite their initial isolation in the mid 1960s. Our recent discovery that the *Phycomyces* *madA* gene is homologous to *Neurospora* *wc-1* suggested that the MADA protein should act as a light-dependent transcription factor and opened the way to the molecular characterization of *Phycomyces* photoreception (23).

We report here the identification and characterization of the *Phycomyces* *madB* gene, encoding a member of the WC-2 family of zinc-finger proteins. In addition, we describe the complete set of *wc* genes in the *Phycomyces* genome and characterize their expression after light exposure. We observe the physical interaction between MADA and MADB, suggesting the presence of a MAD complex that regulates gene transcription by light in *Phycomyces*. Identification of the *madB* gene closes a relevant chapter in *Phycomyces* photobiology by uncovering the molecular identity of a key component of the photoreceptor complex.

Results

Multiple *wc-1* and *wc-2* Genes in the *Phycomyces* Genome. Our discovery that MADA, a protein similar to the *Neurospora*

photoreceptor WC-1, is required for light sensing (23) suggested that *Phycomyces* might use a protein similar to *Neurospora* WC-2 to interact with MADA to mediate blue light responses.

We amplified and cloned a *Phycomyces* gene similar to *wc-2* by PCR. In addition, when the *Phycomyces* genome sequence became available we analyzed the gene set to identify additional *wc* genes. The *Phycomyces* genome contains 3 genes similar to *wc-1* and 4 genes similar to *wc-2*. Two of the *wc-1* genes had been identified, *madA* and *wcoA* (23), and we named the third gene *wcoB* (for white collar one gene B). The 4 *wc-2* genes were named *wctA* to *wctD* (for white collar two gene). The *wctA* gene was identified initially by PCR, and was later renamed *madB* based on our genetic and sequencing results. The predicted gene structure for each *wc* gene was established by PCR amplifying and sequencing the corresponding cDNAs. The only exception was the *wctC* gene, because we were unable to amplify the cDNA from RNA samples obtained from cultures grown in the dark or after exposure to light. All of the *wc-1* genes have 6 exons and 5 introns, and all of the *wc-2* genes have 3 exons and 2 introns (Fig. 1A).

The *Phycomyces* WC proteins are shorter than their *Neurospora* counterparts and contain a PAS domain for protein interactions, and a zinc finger located at the carboxyl terminus, except for WCOB where we could not identify a standard zinc finger although the presence of conserved cysteine residues suggests WCOB may contain a derivative of the GATA-factor class of

zinc fingers (Fig. 1B and Fig. S1). In addition, MADA, WCOA, and WCOB contain an LOV domain that should allow each of these proteins to act as a photoreceptor (Fig. S2). The *Phycomyces* LOV domains contain a conserved cysteine that would support light-dependent formation of a cysteinyl adduct with the flavin chromophore, as observed in plant LOV domains (15) (Fig. S2). The domains in the *Phycomyces* WC-1 proteins suggest that these proteins may function as light-regulated transcription factors.

The *wc* genes were examined by phylogenetic analyses and DNA sequence comparisons to ascertain if these may have arisen through gene duplication, and to assess the extent of duplication in the genome (Fig. 1). The duplication that gave rise to the *madA* and *wcoA* gene pair incorporates an upstream gene encoding a MAP kinase. There is no evidence of a MAP kinase upstream of the third homolog, *wcoB*. The intron positions of the *wcoB* gene are also conserved, with 4 of the 5 introns sharing the same splice sites with *madA* and *wcoA*, which suggests that this gene is related to the other two. Comparison of the zygomycete LOV domains suggests that the event that gave rise to *wc-1* gene triplication occurred before the last common ancestor diverged, and that each of the 3 *Phycomyces* WC-1 homologs is most closely related to its ortholog from the divergent species *Mucor circinelloides* than to the paralogs within species (Fig. 1C).

Examination of the phylogeny and local gene synteny for the 4 *wc-2* homologs reveals that these genes arose through 3 sequential duplication events. All 4 copies of *wc-2* contain 2 introns that are conserved in position. The ancestral *wc-2* gene duplicated, giving rise to 2 paralogs that both subsequently duplicated, giving rise to 2 pairs of genes, corresponding to *wctA/wctB* and *wctC/wctD*. The *wctC* and *wctD* genes are each flanked by a cyclin gene that duplicated along with the *wct* gene. We conclude that the *Phycomyces* *wc* genes were derived from limited local duplications within the genome yet all are related to common ancestral *wc-1* and *wc-2* genes.

***wctA* Gene Has a Splicing Mutation in the Phototropic Mutant *madB*.** We sequenced all of the 7 *wc* genes in 50 representative strains carrying all of the *mad* mutations to identify nucleotide changes that might have been responsible for the deficient phototropic phenotype. We discovered a mutation in the *wctA* gene in *Phycomyces* *madB* mutants. No other mutations were detected in any of the other *wc* genes in any of the other *mad* mutants, with the exception of the characterized *madA* alleles in *madA* mutants (23). The mutation in *madB* strains is a G to A transition at nucleotide 907 of the *wctA* gene (from the initiation ATG) that alters the 5' splicing site of the first intron. This mutation prevents correct mRNA splicing, resulting in longer mRNA that would yield a truncated protein of 327 aa without the zinc finger. The splicing mutation was confirmed after amplification and sequencing of the corresponding cDNAs in the wild type and several *madB* mutants (Fig. 2A).

Mutation in *wctA* Cosegregates with Impaired Phototropism in *madB* Mutants in a Genetic Analysis. To test the hypothesis that *wctA* corresponds to the *madB* locus, genetic crosses between *madB* strains (C109 or C111) and an isogenic strain (A56) were performed. The *wctA* gene was amplified by PCR from each progeny DNA, and cleaved with the restriction enzyme *RsaI*. The G907A mutation identified in *madB* strains mutates an *RsaI* recognition site, resulting in 2 fragments for this *wctA* allele after enzymatic digestion, compared with 3 fragments for the wild type allele. From the C109 x A56 cross (24 progeny from 7 zygospores) the G907A mutation cosegregated with the 16 progeny exhibiting reduced phototropism whereas the remaining 8 progeny with wild type phototropism contained the wild type *wctA* allele. After the C111 x A56 cross (38 progeny from 14 independent zygospores) 15 progeny with reduced light sensi-




Fig. 2. The gene *madB* is similar to *Neurospora* *wc-2*. (A) Aberrant cDNA splicing of the *wctA* gene in the *madB* mutants. The cDNAs for the *wctA* gene from the wild-type strain and 3 *madB* mutants were amplified by PCR and resolved by gel electrophoresis (Upper). Nucleotide sequence of a single transcript cloned from the wild type and *madB* strains C109, C111, or C112. The G907A mutation is in bold. Coding nucleotides are shown in uppercase, and intron nucleotides are shown in lowercase font. For the mutant transcripts the zinc-finger domain will be deleted by introduction of a premature stop codon (underlined) (Lower). (B) Genetic evidence that the G907A mutation in *wctA* is linked to reduced phototropism in *madB* strains. Progeny (15 derived from 15 independent germinated zygospores) from crosses of strains C109 x A56 or C111 x A56 were examined for phototropism, and 50% of the Petri dishes were photographed. The *wctA/madB* gene was amplified by PCR and cleaved with *RsaI* to produce 3 (wild type) or 2 (mutant) fragments. The parents *madB* (phototropism mutant, sex - , 2 fragments) and A56 (phototropism wild type, sex + , 3 fragments) are not shown. The sex locus shows meiotic recombination in the progeny.

tivity had the mutated *wctA* allele with the 23 other wild type progeny containing the wild type allele. A subset of this analysis is illustrated in Fig. 2B. Sex segregated independently of the *madB* mutation (Fig. 2B). In addition, we confirmed the presence of the G907A mutation in the *madB* progeny (A820 and A821) but not in the wild type progeny (A818 and A819) obtained after a cross between a *madB* strain (A520) and a phototropic wild type (C169).

As all of the *madB* strains have a mutation in the *wctA* gene, and the mutation in *wctA* cosegregates with the *madB* phenotype in genetic crosses, we conclude that the *madB* phenotype is caused by a mutation in the *wctA* gene that we will henceforth call *madB*.

Transcriptional Regulation of the *wc* Genes by Blue Light. Blue light induced the expression of several *Phycomyces* *wc* genes to different levels. After exposing *Phycomyces* mycelia to 30 min of blue light we observed a 5-fold increase in *wcoB* mRNA, and a thirtyfold increase in *wcoA* mRNA. The induction by blue light was more pronounced for the *wctB* and *wctD* genes, 180- to 250-fold (Fig. 3A). In comparison, we observed a 10-fold photoactivation for the heat-shock gene *hspA* (32). On the contrary, expression of *madA*, *madB*, and *wctC* genes was not induced by light, and a slight photorepression was detected for *madA* and *madB* (Fig. 3A).

Blue-light exposures of different duration showed that maximum mRNA accumulation for *wcoA*, *wcoB*, *wctB*, and *wctD* genes occurred after 15–30 min of light, but photoactivation was observed after 5-min exposures (Fig. S3). Longer light exposures reduced light-dependent mRNA accumulation, but gene photoactivation was still observed in mycelia exposed to light for 2 hours. On the contrary, blue light caused a reduction in the amount of *madA* mRNA (Fig. S3).

We observed different thresholds for the photoactivation of




Fig. 3. Regulation of gene transcription by blue light. Total RNAs were isolated from mycelia exposed to blue light, or kept in the dark. The amount of mRNA for each gene was assayed by quantitative RT-PCR. Each fluorescent signal was first normalized to the corresponding actin signal to correct for loading errors and then was normalized to the signal obtained in the dark. The plots show the relative photoactivation in 2–15 independent experiments (average ± SEM). (A) Photoactivation of gene expression in the wild type after 30 min of blue light (2.3×10^3 J/m²). (B) Threshold determination for the photoactivation of gene expression in the wild type. Mycelia were exposed to blue light of different intensities during 30 min before mRNA extraction. 1 J/m² of blue light corresponds to 4 μmol/m² of 450 nm. (C) Photoactivation of gene expression in the wild type and *mad* mutants after 30 min of blue light (2.3×10^3 J/m²). The *madA madB* double mutant is shown as *madAB*, and the *madA madB madC* triple mutant is shown as *madABC*. For strain numbers see *SI Text*.

the *wc* genes. A low light exposure, 10^{-5} – 10^{-4} J/m² (4×10^{-5} – 10^{-4} μmol/m² for 450 nm), was sufficient to induce the expression of *wctB*, *wctD*, and *wcoA*. We observed a small but detectable photoinduction for *wctB* at $<10^{-4}$ J/m² that we did not explore further. However, the threshold for the photoactivation of *wcoB*, 1 J/m² (4 μmol/m² for 450 nm), was similar to that of *hspA* but markedly higher than that for the other *wc* genes (Fig. 3B).

The photoactivation of the *wc* genes required the MADA and MADB proteins. A reduced photoactivation was observed in *Phycomyces* strains carrying *madA* alleles that either resulted in an amino acid change in the LOV domain (strain C47), or produced aberrant mRNAs because of a splicing mutation (strain C21) (23). Similarly, the splicing mutation in the *madB* strain reduced the photoactivation of *wcoA*, *wcoB*, and *wctD*, but




Fig. 4. MADA and MADB interact physically. (A) Yeast 2-hybrid assays. The coding regions of the *madA* and *madB* genes were fused adjacent to the AD and BD segments of *S. cerevisiae* GAL4. Plasmids were cotransformed into a *S. cerevisiae* strain in which the GAL4 UAS regulates *URA3*, *HIS3*, and *lacZ* genes. Growth of strains in the absence of uracil (-ura) or histidine (-his) and increased β-galactosidase activity (β-gal ± SE, Miller units) indicate protein–protein interactions. (B) MADA and MADB form a complex when coexpressed in *E. coli*. SDS/PAGE stained with coomassie to detect proteins purified with a GST resin. Arrowheads indicate the MADA and MADB recombinant proteins (Upper). Western hybridisations with anti-MADA and anti-MADB after purification with a GST resin (Lower).

not *wctB* (Fig. 3C). However, MADB was required for *wctB* photoactivation as shown by the differences in photoactivation observed in *madA* and *madA madB* double mutants (Fig. 3C). Mutations in other *mad* genes did not change the relative accumulation of *wc* mRNAs in light-exposed mycelia. A notable exception was a *madH* strain that showed a reduced photoactivation for *wcoB* and *wctD*, similar to the phenotype observed in *madA* strains (Fig. 3C).

MADA and MADB Proteins Interact to Form a MAD Complex. To test protein–protein interactions, the *madA*, *wcoA*, *wcoB*, *madB*, *wctB* and *wctD* cDNAs were fused in frame to the activator (AD) or DNA binding (BD) domains of *S. cerevisiae* Gal4 in vectors for yeast 2-hybrid analysis. The recipient yeast strain carries *URA3*, *HIS3* and *lacZ* reporter genes controlled by the GAL4 promoter region to detect the formation of a complex by their growth on minimal agar and the detection of β-galactosidase activity. When *S. cerevisiae* cells expressed MADA and MADB fused to the AD or BD domains, the *HIS3* and *URA3* reporter genes were both induced and cells grew in the absence of uracil or histidine. In addition, the *lacZ* reporter gene was induced, further validating that MADA and MADB form a complex in yeast cells (Fig. 4A). In contrast, *S. cerevisiae* cells expressing only MADA or MADB fused either to AD or BD domains did not grow on selective media. We did not observe any effect of light on the reporter gene-dependent growth of the strains. No other positive interactions were observed in yeast 2-hybrid assays between any of the other WC proteins: WCOA, WCOB, WCTB, and WCTD (Fig. S4).

As confirmation of the MADA/MADB interaction we expressed the cDNAs for these 2 genes in *E. coli* and carried out copurification assays (Fig. 4B). The *madA* cDNA was fused in

frame to a GST domain, and the *madB* cDNA was fused in frame to a 6-His tag. The *E. coli* protein fraction that bound the GST resin was eluted, and resolved by electrophoresis. The molecular nature of each protein band was revealed after hybridization with specific polyclonal antisera against MADA or MADB. Because we observed the MADB-His tagged protein after GST purification we concluded that the GST resin purified a MADA-MADB protein complex, corroborating the yeast 2-hybrid results (Fig. 4B).

Discussion

Research on the molecular biology of *Phycomyces* sensory transduction has been limited despite the experimental advantages of a large unicellular structure capable of sensing many environmental signals. However, the isolation and characterization of phototropic *mad* mutants, initially in Delbrück's lab and later in other laboratories, allowed the identification of genetic loci that are required for reception and processing of the light signal. Among them, the *madA* and *madB* gene products are key components of the *Phycomyces* light-transduction pathway (3). The identification of the nature of *madB* as a *wc-2* gene, together with our previous discovery that *madA* is a *wc-1* gene (23), provide 2 essential elements of *Phycomyces* photoreception.

The MADA and MADB interaction suggests that this photoreceptive complex acts as its *Neurospora* counterpart by binding to the promoters of light-regulated genes. However, the role of the MAD complex in short-term phototropic photoreponses is less clear. The *Phycomyces* sporangiophore is a single cell cylinder, and phototropic bending is observed within 5 minutes after applying unilateral light (33), a short time for a light-dependent transcription factor to operate efficiently through gene transcription and protein translation. It is possible that, in addition, the MAD complex may operate through the light-dependent postranslational modification of other regulatory proteins or by other mechanisms not yet described for WC-1 and WC-2 in *Neurospora*.

The *Phycomyces* genome contains 3 *wc-1* and 4 *wc-2* genes that originated via gene duplication events that occurred early in the zygomycete evolutionary lineage, and have not been observed in other fungal groups. The absence of any additional interactions between WC proteins in yeast 2-hybrid assays suggests that either these proteins participate in weak interactions that are not sufficient for detection in yeast, that species-specific modifications are required, or they do not interact at all, a surprising result considering the presence of PAS domains in all of the *Phycomyces* WC proteins. It is possible that additional proteins, perhaps the MAD complex, serve as scaffolds for additional interactions between the remaining WC proteins.

The observation that *madA* and *madB* are not induced by light suggests that the proteins are present in the dark poised to activate gene transcription and phototropism upon light exposure. The activation by light of *wcoA*, *wcoB*, *wctB*, and *wctD* suggests that the corresponding gene products may be required after initial exposure to light, or to respond to higher light intensities. These genes may be required for a particular light response in *Phycomyces*, as has been shown for some of the *wc-1* genes in *Mucor* (24). We can propose that in the dark the MAD complex will be available for light perception. After low light exposure, the activation of *wcoA*, *wctB*, and *wctD* genes would provide additional photoreceptors that could be devoted to low light reception. Higher light intensities would activate *wcoB*, perhaps to perceive high-intensity photons. However, the differences in the thresholds for *wc* gene activation suggest that additional photoreceptor proteins, most likely WCOA or WCOB, collaborate in the dark with the MAD complex to activate gene transcription at different intensity ranges. Our observation that *madA* and *madB* are required for the photoactivation of these genes supports this hypothesis. Further

characterization of the *Phycomyces* WC proteins will allow us to establish the detailed role these proteins play in photoreception.

It is remarkable that of the remaining 8 *mad* photoreponse mutants (*madC-madJ*), none result from mutations in the remaining 5 *wc* genes. The identity of the gene products encoded by *madC-madJ* will be uncovered by meiotic mapping studies and may reveal light sensing machinery, or components that lie downstream of the primary MAD photoreceptor complex in the sensory transduction cascade. Our discovery that phototropism and other responses to light in *Phycomyces* require the activity of a complex composed of MADA and MADB proteins will help to initiate the molecular characterization of *Phycomyces* photoreception, but it is clear that much remains to be learned from this model fungal system that will advance our understanding of how organisms perceive their environment.

Materials and Methods

Strains. We used the standard wild type strain of *P. blakesleeana* (NRRL1555). Other wild type and *mad* mutant strains are described in Table S1. *Escherichia coli* DH5-alpha was used for cloning plasmids.

Cloning and Characterization of *Phycomyces wc* Genes in the Wild Type and *mad* Mutant Strains.

A fragment of the *Phycomyces wc-2* gene was amplified by PCR from genomic DNA using primers for conserved areas of the gene (Table S2). The 1 kb PCR product was cloned into plasmid pGEM-T (Promega) and sequenced. Other *wc-1* and *wc-2* genes were identified in the subsequently released *Phycomyces* genome sequence (Joint Genome Institute <http://www.jgi.doe.gov>). The predicted ORFs were confirmed or corrected after amplification and sequencing the cDNAs (see primers in Table S2), except the cDNA for *wcoB* that was obtained in 2 steps (see SI Text). *Phycomyces* genomic DNA was used to amplify and sequence the *wc* genes in *mad* mutant strains (see the list of strains in Table S1 and the list of primers in Table S2). DNA manipulations followed standard methods (34). For domain predictions, protein comparisons, and phylogenetic analysis see SI Text.

Mating Assays and Genetic Segregation. Crosses were conducted between *madB* mutant strains C109 and C111 (–) and isogenic strain A56 (+) following methods described in ref. 23. A 478-bp fragment of the *madB* gene was amplified with primers (Table S2), and cut with *RsaI* restriction enzyme. The wild type allele results in 3 fragments (217, 169 and 92 bp) and the mutant allele in 2 fragments (309 and 169 bp). Mating type for all progeny was assessed by crossing to (+) and (–) parental strains to confirm meiotic segregation.

Light Exposures, RNA Isolation and cDNA Synthesis, Quantitative RT-PCR. The *Phycomyces* dark-grown mycelia were exposed to blue light at the age of 48 h during 30 min unless otherwise stated, and used for RNA extraction as reported (32). Total RNA was used for cDNA synthesis and amplification (see primers in Table S2). Quantitative PCRs were performed to determine relative mRNA abundance using 1-step RT-PCR in a 7500 Real Time PCR System (Applied Biosystems) as described in ref. 35 (see primers in Table S2). The results for each gene were normalized to the corresponding results obtained with the actin gene (*act-1*) to correct for sampling errors. Then, the results obtained with each sample were normalized to the RNA sample from the corresponding mycelia grown in the dark. For phototropism measurements, strains were illuminated with white light of an intensity below that required for *madB* response, as determined empirically.

Yeast 2-Hybrid Experiments. cDNAs of *madA*, *wcoA*, *wcoB*, *madB*, *wctB* and *wctD* were amplified by RT-PCR, and cloned into plasmids pDESTTM32 and pDESTTM22 in the *S. cerevisiae* reporter strain MaV203 using the ProQuest™ Two-Hybrid System (Invitrogen). Double transformants were selected on media lacking leucine and tryptophan. Interactions were assessed by growth in the absence of uracil or histidine with 10 mM 3-aminotriazole, and by β -galactosidase assays.

Expression and Detection of the MADA/MADB Complex in *E. coli*. *Phycomyces* cDNAs for MADA and MADB were amplified and cloned into the expression vectors pGEX4T1 (GST-tag) (GE Healthcare) and pET28a (6XHis) (Novagen). The fusion proteins were expressed in *E. coli* strain Rosetta2 (DE3) pLysS (Novagen) transformed with each or both recombinant plasmids, and the GST proteins purified with a GST-bind resin (Novagen). Purified proteins were used for SDS/PAGE and western hybridization using an anti-MADA polyclonal antibody and an anti-MADB polyclonal antibody (Pacific Immunology) (see SI Text for details).

ACKNOWLEDGMENTS. We thank E. Cerdá-Olmedo for scientific advice, D. Perez del Camino for technical help, and G. Gutiérrez for phylogenies. This work was supported by European funds (ERDF), Spanish Grants BIO2006-14897, AGL2005-08081, P06-CVI-01650 (J. Andalucía), and GR64 (to J. Castilla-

León); and National Institutes of Health Grants AI039115 and AI073917. J.R.R. held an European Molecular Biology Organization short-term fellowship in the laboratory of J.M.C. We acknowledge access to the *Phycomyces* genome sequence (US Department of Energy, Joint Genome Institute).

1. Bahn YS, et al. (2007) Sensing the environment: Lessons from fungi. *Nat Rev Microbiol* 5:57–69.
2. Corrochano LM, Galland P (2006) In *The Mycota I Growth, Differentiation and Sexuality*, The Mycota, eds Kues U, Fischer R (Springer-Verlag, Berlin), 2nd Ed Vol I, pp 233–259.
3. Cerdá-Olmedo E (2001) *Phycomyces* and the biology of light and color. *FEMS Microbiol Rev* 25:503–512.
4. Linden H, Ballario P, Macino G (1997) Blue light regulation in *Neurospora crassa*. *Fungal Genet Biol* 22:141–150.
5. Cerdá-Olmedo E, Lipson ED eds (1987) *Phycomyces* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).
6. Bergman K, et al. (1969) *Phycomyces*. *Bacteriol Rev* 33:99–157.
7. Galland P, Lipson ED (1987) Blue-light reception in *Phycomyces* phototropism: Evidence for two photosystems operating in low- and high-intensity ranges. *Proc Natl Acad Sci USA* 84:104–108.
8. Bergman K, Eslava AP, Cerdá-Olmedo E (1973) Mutants of *Phycomyces* with abnormal phototropism. *Mol Gen Genet* 123:1–16.
9. Orejas M, Peláez MI, Alvarez MI, Eslava AP (1987) A genetic map of *Phycomyces blakesleeanus*. *Mol Gen Genet* 210:69–76.
10. Campuzano V, Galland P, Eslava AP, Alvarez MI (1995) Genetic characterization of two phototropism mutants of *Phycomyces* with defects in the genes *madl* and *madj*. *Curr Genet* 27:524–527.
11. Liu Y, He Q, Cheng P (2003) Photoreception in *Neurospora*: A tale of two White Collar proteins. *Cell Mol Life Sci* 60:2131–2138.
12. Ballario P, et al. (1996) White collar-1, a central regulator of blue light responses in *Neurospora*, is a zinc finger protein. *EMBO J* 15:1650–1657.
13. Froehlich AC, Liu Y, Loros JJ, Dunlap JC (2002) White Collar-1, a circadian blue light photoreceptor, binding to the *frequency* promoter. *Science* 297:815–819.
14. He Q, et al. (2002) White collar-1, a DNA binding transcription factor and a light sensor. *Science* 297:840–843.
15. Christie JM (2007) Phototropin blue-light receptors. *Annu Rev Plant Biol* 58:21–45.
16. Zoltowski BD, et al. (2007) Conformational switching in the fungal light sensor Vivid. *Science* 316:1054–1057.
17. Linden H, Macino G (1997) White collar 2, a partner in blue-light signal transduction, controlling expression of light-regulated genes in *Neurospora crassa*. *EMBO J* 16:98–109.
18. He Q, Liu Y (2005) Molecular mechanism of light responses in *Neurospora*: From light-induced transcription to photoadaptation. *Genes Dev* 19:2888–2899.
19. Belden WJ, Loros JJ, Dunlap JC (2007) Execution of the circadian negative feedback loop in *Neurospora* requires the ATP-dependent chromatin-remodeling enzyme CLOCKSWITCH. *Mol Cell* 25:587–600.
20. Idnurm A, Heitman J (2005) Light controls growth and development via a conserved pathway in the fungal kingdom. *PLoS Biol* 3:e95.
21. Lu YK, Sun KH, Shen WC (2005) Blue light negatively regulates the sexual filamentation via the Cwc1 and Cwc2 proteins in *Cryptococcus neoformans*. *Mol Microbiol* 56:480–491.
22. Terashima K, Yuki K, Muraguchi H, Akiyama M, Kamada T (2005) The *dst1* gene involved in mushroom photomorphogenesis of *Coprinus cinereus* encodes a putative photoreceptor for blue light. *Genetics* 171:101–108.
23. Idnurm A, et al. (2006) The *Phycomyces madA* gene encodes a blue-light photoreceptor for phototropism and other light responses. *Proc Natl Acad Sci USA* 103:4546–4551.
24. Silva F, Torres-Martínez S, Garre V (2006) Distinct white collar-1 genes control specific light responses in *Mucor circinelloides*. *Mol Microbiol* 61:1023–1037.
25. Silva F, et al. (2008) A RING-finger protein regulates carotenogenesis via proteolysis-independent ubiquitylation of a White Collar-1-like activator. *Mol Microbiol* 70:1026–1036.
26. Blumenstein A, et al. (2005) The *Aspergillus nidulans* phytochrome FphA represses sexual development in red light. *Curr Biol* 15:1833–1838.
27. Purschwitz J, et al. (2008) Functional and physical interaction of blue- and red-light sensors in *Aspergillus nidulans*. *Curr Biol* 18:255–259.
28. Bayram O, Biesemann C, Krappmann S, Galland P, Braus GH (2008) More than a repair enzyme: *Aspergillus nidulans* photolyase-like CryA is a regulator of sexual development. *Mol Biol Cell* 19:3254–3262.
29. Bayram O, et al. (2008) VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science* 320:1504–1506.
30. Corrochano LM (2007) Fungal photoreceptors: Sensory molecules for fungal development and behaviour. *Photochem Photobiol Sci* 6:725–736.
31. Herrera-Estrella A, Horwitz BA (2007) Looking through the eyes of fungi: Molecular genetics of photoreception. *Mol Microbiol* 64:5–15.
32. Rodríguez-Romero J, Corrochano LM (2004) The gene for the heat-shock protein HSP100 is induced by blue light and heat-shock in the fungus *Phycomyces blakesleeanus*. *Curr Genet* 46:295–303.
33. Galland P (1990) Phototropism of the *Phycomyces* sporangiophore: A comparison with higher plants. *Photochem Photobiol* 52:233–248.
34. Sambrook J, Russell DW (2001) *Molecular Cloning. A Laboratory Manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor).
35. Navarro-Sampedro L, Yanofsky C, Corrochano LM (2008) A genetic selection for *Neurospora crassa* mutants altered in their light regulation of transcription. *Genetics* 178:171–183.