



Published in final edited form as:

Cell Metab. 2021 June 01; 33(6): 1124–1136.e5. doi:10.1016/j.cmet.2021.03.008.

Oxylipin biosynthesis reinforces cellular senescence and allows detection of senolysis

Christopher D. Wiley^{1,2,*}, Rishi Sharma¹, Sonnet S. Davis¹, Jose Alberto Lopez-Dominguez¹, Kylie P. Mitchell¹, Samantha Wiley¹, Fatouma Alimirah¹, Dong Eun Kim¹, Therese Payne¹, Andrew Rosko¹, Eliezer Aimontche¹, Sharvari M. Deshpande¹, Francesco Neri¹, Chisaka Kuehnemann¹, Marco Demaria^{1,3}, Arvind Ramanathan^{1,4,*}, Judith Campisi^{1,5,6,*}

¹Buck Institute for Research on Aging, 8001 Redwood Blvd, Novato, CA, USA, 94945

²Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts, USA, 02111

³European Research Institute for the Biology of Aging (ERIBA), University Medical Center Groningen (UMCG), Antonius Deusinglaan 1, 9713 Groningen, Netherlands

⁴Institute for Stem Cell Science and Regenerative Medicine, GKVK - Post, Bellary Rd, Bengaluru, Karnataka 560065, India

⁵Lawrence Berkeley National Laboratory, 1 Cyclotron Rd, Berkeley, CA, USA, 94720

⁶Lead contact

Summary

Cellular senescence is a stress or damage response that causes a permanent proliferative arrest and secretion of numerous factors with potent biological activities. This senescence-associated secretory phenotype (SASP) has been characterized largely for secreted proteins that participate in embryogenesis, wound healing, inflammation and many age-related pathologies. By contrast, lipid components of the SASP are understudied. We show that senescent cells activate the biosynthesis of several oxylipins that promote segments of the SASP and reinforce the proliferative arrest. Notably, senescent cells synthesize and accumulate an unstudied intracellular prostaglandin, 1a,1b-dihomo-15-deoxy-delta-12,14-prostaglandin J2. Released 15-deoxy-delta-12,14-prostaglandin J2 is a biomarker of senolysis in culture and *in vivo*. This and other prostaglandin D2-related lipids promote the senescence arrest and SASP by activating RAS

*Correspondence: christopher.wiley@tufts.edu (C.D.W.), arvind@instem.res.in (A.R.), jcampisi@buckinstitute.org (J.C.).

Author contributions. CW and AR conceived, designed, and interpreted the lipid profiling experiments. SSD and RS designed and conducted all mass spectrometry experiments with guidance from AR. AR, RS, and AR conducted post-experiment informatics and interpretation. SSD, KM, SW, and SMD extracted lipids and conducted analyses. CW, JALD, FA, DK, and MD conducted mouse experiments. EA, FN, and CK conducted additional assays. All other experiments were designed and conducted by CW, with guidance from AR and JC. CW, AR, RS, and JC wrote the paper and addressed reviewer comments.

Declaration of Interests. JC is a founder of Unity Biotechnology, which develops senolytic therapies. CW, AR, and JC are inventors on a patent for detection of oxylipins as biomarkers of senolysis.


Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

signaling. These data identify an important aspect of cellular senescence and a method to detect senolysis.

eTOC

Senolytics, transgenic and pharmacological interventions that selectively kill senescent cells, are currently in clinical trials aiming to treat age-related degenerative pathologies. Here, Wiley et al. discover that senescent cells produce multiple signaling lipids known as oxylipins. One oxylipin, dihom-15d-PGJ2, promotes features of senescence by activating RAS and is released from cells during senolysis, serving as the first biomarker of the process in culture and *in vivo*.

Graphical Abstract



Introduction

Cellular senescence is a multifaceted response to physiological and stress-generated signals, resulting in an essentially permanent arrest of cell proliferation and several phenotypic changes, including cellular hypertrophy, nuclear and epigenetic rearrangements, and metabolic alterations (Kuilman et al., 2010; Wiley and Campisi, 2016). Cell culture and transgenic mouse studies show that senescent cells can drive several age-associated pathologies in part by secreting a myriad of biologically active molecules. These molecules, which include inflammatory cytokines and chemokines, proteases and growth factors, can have potent local – and potentially systemic – effects on tissues (Acosta et al., 2008; Coppe

et al., 2008). Senescent cells increase with age in human and mouse tissues (Campisi, 2013; Wiley and Campisi, 2016), which, at least in mice, limits both median life span and health span (Baker et al., 2016).

Thus far, the senescence-associated secretory phenotype (SASP) has been studied primarily with regard to secreted proteins (Coppe et al., 2008; Wiley et al., 2019b). Here we show that senescent cells also synthesize a number of oxylipins – a class of biologically active lipids that arise from oxygenation of polyunsaturated fatty acids (PUFAs). Oxylipins have diverse physiological effects, including inflammation, fever, vasoconstriction and vasodilation, pain, hair loss, asthma and fibrosis (Funk, 2001; Soberman and Christmas, 2003). Senescent cells are known to produce prostaglandin E2 (PGE2) (Cormenier et al., 2018; Dagouassat et al., 2013; Kabir et al., 2016; Martien et al., 2013; Yang et al., 2011; Zdanov et al., 2007), and the leukotriene-producing 5-lipoxygenase activity was shown to increase in replicative and oncogene-induced senescence, where it reinforced senescence-associated phenotypes by increasing reactive oxygen species (Catalano et al., 2005). We show that senescent human cells synthesize a much larger set of oxylipins, including PGD2 and its derivatives, 22-carbon PUFAs and monounsaturated fatty acids (MUFAs). Further, the dihomoprostaglandins – less-studied derivatives of adrenic acid – are the primary prostaglandins produced by senescent cells, and one of these derivatives, dihom-15d-PGJ2, is an intracellular biomarker of senescence and its release from cells is a hallmark of senolysis in culture and *in vivo*.

Results

Cellular senescence activates oxylipin biosynthesis

The senescence arrest and SASP entail significant changes in metabolism (Wiley and Campisi, 2016), including lipid metabolism (Buratta et al., 2017; Cadenas et al., 2012; Catalano et al., 2005; Flor et al., 2017; Quijano et al., 2012). To better understand these changes, we extracted intracellular lipids from cultured human fibroblasts (IMR-90) that were either proliferating (PRO, 10% serum), quiescent (QUI, 0.2% serum) or induced to senesce by ionizing radiation (IR), then cultured in either 10% or 0.2% serum. We also extracted lipids from mitochondrial dysfunction-associated senescent (MiDAS) cells (0.2% serum) (Wiley et al., 2016). Thus, we controlled for differences in growth state (QUI vs PRO vs SEN(IR) and MiDAS) and culture media (0.2% vs 10% serum). We measured the relative abundance of the extracted lipids by mass spectrometry.

The lipid profiles (Table S1) showed that certain subsets of lipids significantly increased or decreased upon senescence. Most notable were striking elevations in the relative abundance of oxylipins: a class of potent signaling lipids derived from 20- and 22-carbon fatty acids such as arachidonic (AA) and adrenic acid (AdA) (Figure 1A). We also observed changes in intracellular ceramides and other lipids (Table S1).

The most highly elevated senescence-associated prostaglandin was 1a,1b-dihomo-15-deoxy- Δ 12,14-prostaglandin J2 (dihomo-15d-PGJ2). However, 1a,1b-dihomo-prostaglandin D2 (PGD2), 1a,1b-dihomo-prostaglandin E2 (PGE2), and other variants of prostaglandins PGD2, PGE2 and PGF2 α also increased (Figure 1A). Additionally, senescent cells

contained higher levels of specific leukotrienes, notably leukotriene B₄ (LTB₄) and the related lipoxygenase product 5-hydroxyeicosatetraenoic acid (5-HETE) (Figure 1A). Consistent with these findings, the 20-carbon oxylipin precursors arachidonic acid (AA), eicosapentanoic acid (EPA), and dihomo-gamma-linoleic acid (DGLA) were elevated in senescent cells (Figure 1B).

The 22-carbon lipid AdA, a product of AA elongation (Jump, 2009; Ohno et al., 2010) and precursor of dihomo prostaglandins (Sprecher et al., 1982), also increased in senescent cells (Figure 1B). Further, senescent cells contained higher levels of the 22-carbon PUFAs docosahexanoic acid (DHA) and docosapentanoic acid (DPA) and the 22-carbon MUFA erucic acid, but not the 22-carbon saturated fatty acid behenic acid. Additionally, incubating senescent cells with deuterated AA resulted in an increase in deuterated AdA, confirming elongation of AA to AdA by senescent cells (Figure S1A). Thus, senescent cells show a selective and marked increase in oxylipins and elongated PUFAs and MUFAs.

To understand the mechanism by which senescent cells accumulate these lipids, we used qPCR to measure mRNA levels of several genes that function in oxylipin biosynthesis. We compared cells induced to senesce by ionizing radiation [SEN(IR)], mitochondrial dysfunction (MiDAS) and RAS oncogene activation [SEN(RAS)] (Figure 1C). Among the mRNAs that increased significantly were those encoding prostaglandin synthases *PTGS2* (COX-2), *PTGDS*, *PTGES* and *TBXAS*, but not *PTGIS*, which showed lower expression. mRNAs encoding leukotriene biosynthetic genes, including *ALOX5* (5-LO), *ALOX15*, *ALOX5AP*, *LTC4S* and to a lesser extent *LTA4H* were similarly elevated. We observed similar qPCR results for oxylipin biosynthesis genes using human umbilical cord endothelial cells induced to senesce by IR [SEN(IR) HUVECs] (Figure S1B) and replicatively senescent [SEN(Rep)] IMR-90 fibroblasts (Figure S1C).

To determine whether oxylipin synthesis is elevated in senescent cells *in vivo*, we used a mouse model in which p16^{Ink4a}-positive senescent cells can be eliminated by administration of ganciclovir (GCV) (p16-3MR mice) (Demaria et al., 2014). We treated p16-3MR mice with doxorubicin (DOXO) - an inducer of widespread senescence in mice (Demaria et al., 2017) - and measured mRNA levels of several oxylipin synthases in livers (Figure 1D). Much like cultured human cells, oxylipin synthase mRNAs increased significantly upon inducing senescence *in vivo* in mice. Further, eliminating senescent cells with GCV lowered both oxylipin synthase (Figure 1D) and p16^{Ink4a} (Figure S1D) expression, confirming that the elevation was senescence-associated. Kidneys from aged mice also showed elevated levels of oxylipin synthase mRNAs (Figure S1E).

Since many oxylipins are secreted, we collected conditioned media from IMR-90 fibroblasts that were either quiescent or induced to senesce by mitochondrial dysfunction (MiDAS), ionizing radiation [SEN(IR)] or RAS oncogene activation [SEN(Ras)]. Using ELISAs, we measured the levels of PGD₂ - a secreted prostaglandin that is also a precursor of 15d-PGJ₂ (Figure 1E). All senescent cells showed elevated PGD₂ secretion, although the ELISA does not distinguish dihomo-PGD₂ from PGD₂. The same was true for an epithelial-derived human cancer cell line (HEPG2 - Figure 1F) and primary human umbilical vein endothelial cells (HUVECs - Figure 1G) induced to senesce by IR. Notably, hydrocortisone

– a glucocorticoid that prevents AA release from intracellular membranes by activating annexin A1 (Perretti and D’Acquisto, 2009), prevented the senescence-associated release of PGD2 (Figure S1F), suggesting that the primary source of AA for prostaglandin synthesis is likely intracellular membranes. Since glucocorticoids can extend the replicative lifespan of cultured cells (Rosner and Cristofalo, 1979) and attenuate the SASP (Laberge et al., 2012), one or more AA-derived oxylipins might prevent senescence and promote the SASP.

Prostaglandins are inactivated primarily by coordinated import by a prostaglandin transporter (gene name: *SLCO2A1*) coupled to NAD⁺-dependent oxidation of the 15(S)-hydroxyl group to an inactive 15-keto-prostaglandin metabolite by 15-hydroxy-prostaglandin dehydrogenase (PGDH) (Nomura et al., 2004). Consistent with increased intracellular levels of prostaglandins (Figure 1A), *SLCO2A1* mRNA levels increased >100-fold in human fibroblasts induced to senesce by IR and mitochondrial dysfunction (Figure 1H) and >50-fold in RAS-induced senescent cells (Figure 1I). Smaller increases occurred during replicative senescence (Figure S1) and in IR-induced senescent HUVECs (Figure 1J). However, there was no commensurate increase in *PGDH* mRNA during any form of senescence (Figures 1H–1J, Figure S1H). Indeed, *PDGH* mRNA levels decreased in replicatively senescent fibroblasts and IR-induced senescent HUVECs (Figure 1J, Figure S1G). Thus, senescence entails an activation of several oxylipin-synthesis pathways, especially those that generate PGD2-related lipids (Figure S1H). This activation is likely coupled to increased import without inactivation by PGDH, resulting in intracellular accumulation of 15-deoxy-prostaglandins.

Dihomo-15d-PGJ2 is a biomarker of senolysis in culture and *in vivo*.

Since dihom-15d-PGJ2 accumulated inside senescent cells (Figure 1A) and was not substantially secreted (Figure 2A–D), we hypothesized that senolysis -- the death of senescent cells -- might release dihom-15d-PGJ2 into the media of cultured cells and plasma of mice. To test this hypothesis, we cultured quiescent (QUI), mitochondrial dysfunction-associated senescent (MiDAS) and IR-induced senescent [SEN(IR)] IMR-90 fibroblasts in the presence of DMSO or the senolytic drug ABT-263 (10 μ M) (Chang et al., 2016). After 24 hours, we collected conditioned media. At 10 μ M, ABT-263 killed at least 90% of cells, regardless of senescence or quiescence. We then measured 15d-PGJ2 levels in the media by commercial ELISA - which detects both 15d-PGJ2 and dihom-15d-PGJ2. Since the assay cannot distinguish between these lipids, we refer to the ELISA signal as “(dh)15d-PGJ2” to acknowledge that this signal may come from either lipid. Media from MiDAS cells (which have a low basal level of apoptosis) showed some (dh)15d-PGJ2, but this increased significantly when the cells were treated with ABT-263 (Figure 2A). Similarly, SEN(IR) media contained only small amounts of (dh)15d-PGJ2, but contained much more after ABT-263-mediated senolysis (Figure 2A).

To determine whether (dh)15d-PGJ2 predicted senolysis in other cell lineages, we measured its release upon ABT-263-induced senolysis in senescent HEPG2 and HUVEC cells. In each case, senolysis increased (dh)15d-PGJ2 (Figures 2B and 2C). Senescent HUVECs showed some (dh)15d-PGJ2 when vehicle-treated (Figure 2C), but, since some HUVECs detach from the plate and undergo anoikis during routine culture, this elevation was expected. Since

the ELISA antibody was raised against 15d-PGJ2, not specifically dihomom-15d-PGJ2, we used mass spectrometry to determine whether dihomom-15d-PGJ2 was released by senescent cells. We assayed conditioned media from QUI+DMSO, SEN(IR)+DMSO, QUI+ABT-263, and SEN(IR)+ABT-263 cells. In agreement with Figure 1A and the ELISA data, we detected substantially elevated dihomom-15d-PGJ2 only in SEN(IR)+ABT-263 cells (Figure 2D). By comparison, non-senescent IMR-90s did not release dihomom-15d-PGJ2 when treated with non-selective apoptosis inducers (staurosporine or TRAIL) (Figure S2A), indicating that dihomom-15d-PGJ2 is a distinct feature of senolysis in culture.

To determine whether dihomom-15d-PGJ2 could be used to detect senolysis *in vivo*, we treated mice with DOXO and allowed them to accumulate senescent cells over 6 weeks, as described (Demaria et al., 2017). We then administered ABT-263 (50 mg/kg) by gavage (Chang et al., 2016). After 3 hours, mRNA levels of *p16^{Ink4a}* and *p21^{Waf1}* declined significantly in the epididymal fat of ABT-263-treated mice (Figures 2E and 2F). Further, ELISAs showed elevated (dh)15d-PGJ2 in the plasma of DOXO + ABT-263 treated mice, but not control mice (Figure 2G). Similarly, 12 hours after ABT-263-induced senolysis, urine (dh)15d-PGJ2 was elevated relative to controls (Figure 2H). Ion chromatograms from mouse plasma from DOXO + ABT-263 mice revealed the 343m/z peak indicative of dihomom-15d-PGJ2 (Figure 2I; Figure S2B), which was absent from plasma from PBS + Vehicle, PBS + ABT-263 and DOXO + Vehicle mice (Figure 2I, Figure S2C). We did not detect a (non-dihomom)15d-PGJ2 peak in plasma from any treatment group examined (Figure S2D). We conclude that senolysis is accompanied by the release of dihomom-15d-PGJ2 from senescent cells *in vivo*.

Dihomom-15d-PGJ2 was highly elevated and abundant (~1–5 μ M) in senescent cells - but is considerably unstudied (detected in one published list of analytes) (Lee et al., 2016). To confirm its identity, we obtained a synthetic dihomom-15d-PGJ2 standard (Jubilant Chemsys; 94% purity) (Figure S3A, Figure S3B), and used it to confirm the identity of the senolysis biomarker. Dihomom-15d-PGJ2 differs from 15-PGJ2 by an additional C₂H₄ group, resulting in a mass shift of 28 Da (from 315.2 to 343.2 m/z) (Figure S2B). In addition, the MS/MS fragmentation patterns of the standard (Figure 3A) revealed peaks at 299.2 m/z (loss of CO₂), 283.3 m/z (loss of CO₂ + H₂O), 255.2 (loss of CO₂ + H₂O + C₂H₄), and 233.2 (loss of H₂O + C₇H₈). Retention times between the dihomom-15d-PGJ2 standard and samples from senolytic-treated culture or plasma samples closely matched (Figure 3B, filled peaks). Samples also showed a second peak at a later time that was not specific for senolysis (Figure 3B, unfilled peaks). Analysis of the senolysis-specific peaks by MS/MS fragmentation revealed characteristic peaks at 283.3 m/z, 255.2 m/z and 233.2 m/z (Figure 3C), as well as the 299.2 peak (presented separately in Figure 3D). Together, these peaks confirmed that dihomom-15d-PGJ2 was the senolysis-released lipid detected in our analyses.

Prostaglandins promote senescent phenotypes

To understand the consequence of senescence-associated oxylipin biosynthesis, we considered that cyclooxygenase 2 (Ptgs2) overexpression causes premature aging phenotypes in mice (Kim et al., 2016), and oxylipin synthesis inhibitors extend male mouse life span (Strong et al., 2008). To determine whether prostaglandin synthesis promotes

senescent phenotypes, we treated irradiated cells [SEN(IR)] with either of two prostaglandin synthase 2 (PTGS2/COX-2) inhibitors: CAY-10404 (CAY) or NS-398 (NS), which delays replicative senescence (Han et al., 2004). Both inhibitors suppressed secreted PGD2 levels, as determined by ELISA (Figure 4A). Notably, when counting cells for normalization (Figure 4A), SEN(IR) cells exposed to either inhibitor showed increased cell number relative to control cells (Figure S4A). This finding suggests that prostaglandin biosynthesis might reinforce the senescence proliferative arrest.

To address this possibility, we irradiated IMR-90 fibroblasts with a lower IR dose (5 Gy vs 10 Gy), which causes a senescence growth arrest in 90% vs >99% of cells, and treated them with CAY or NS. CAY and, to a greater degree NS, significantly increased EdU incorporation (Figure 4B), decreased senescence-associated beta-galactosidase (SA-Bgal) positivity (Figure 4C) and decreased both *CDKN1A* (p21) and *CDKN2A* (p16^{INK4a}) mRNA levels (Figures 4D and 4E).

Most prostaglandins and leukotrienes are secreted and many can promote cell proliferation and inflammation (Castellone et al., 2005; Dennis and Norris, 2015; Ricciotti and FitzGerald, 2011). Thus, the senescence-associated oxylipins we detected might drive the inflammatory arm of the SASP. To test this idea, we inhibited prostaglandin synthesis with CAY or NS in SEN(IR) cells and measured secreted levels of the proinflammatory SASP factor interleukin 6 (IL-6) (Figure 4F). Both compounds reduced secreted IL-6 levels (Figure 4F).

To assess effects on other SASP factors, we extracted mRNA from each treatment group and measured gene expression by qPCR. COX-2 inhibition by either compound decreased mRNA levels of all the inflammatory cytokines assayed, as well as *MMP3*, but had no effect on *VEGF*—part of a pro-angiogenic subset of SASP factors (Figure 4G). COX-2 inhibition similarly affected cell number, IL-6 secretion, EdU incorporation and colony formation in oncogenic RAS-induced senescent cells (Catalano et al., 2005; Serrano et al., 1997) (Figures S4B–S4F). Thus, prostaglandin synthesis promotes phenotypes associated with senescence, including cell cycle arrest and the expression and secretion of certain SASP factors.

Dihomo-15d-PGJ2 markedly increased in senescent cells, suggesting this lipid and/or similar cyclopentenone prostaglandins might drive senescence phenotypes. To test this idea, we treated non-senescent cells with 15d-PGJ2 or dihom-15d-PGJ2 and measured markers of senescence. Both lipids reduced EdU incorporation (Figure 4H) and *LMNB1* mRNA (Figure 4M), increased SA-Bgal activity (Figure 4I–4J) and *CDKN1A/p21* mRNA levels (Figure 4K). Only dihom-15d-PGJ2 increased *CDKN2A/p16* mRNA (Figure 4L). Further, although both lipids induced several SASP factor mRNAs, dihom-15d-PGJ2 induced higher levels of these mRNAs than 15d-PGJ2 (Figure 4N). Both lipids also induced PTGS2/COX-2, PTGES, and PTGDS mRNAs, suggesting a positive feedback loop maintains prostaglandin synthesis and senescence phenotypes. PGD2 and PGJ2 had similar effects on senescence phenotypes (Figures S4G–S4L), as did PGE2 to a lesser degree, as reported (Martien et al., 2013). In agreement with positive feedback by prostaglandins, PTGS2/COX-2 expression declined upon inhibition of COX-2 with NS or CAY (Figure S4M). Phenotypes induced by PGE2 were unlikely due to senescence, as cells treated with

PGE2 - but not PGD2 or PGJ2 - became smaller and resumed proliferation once PGE2 was removed (Figures S4N–S4O). Thus, PGD2 and related metabolites can promote senescence.

To further test this idea, we depleted prostaglandin D synthase (PTGDS) by lentiviral-delivered shRNA (shPTGDS), and measured mRNA levels of SASP factors following induction of senescence by IR. One shRNA efficiently reduced PTGDS expression (Figure S4P), and also reduced several SASP factor mRNAs compared to senescent cells transduced with a control shRNA (shGFP) (Figure 4O). Together with Figures 4H–4N, we conclude that PGD2 and related lipids promote cellular senescence phenotypes.

Extracellular PGD2 signals through two receptors, DP1 (*PTGDR*) and DP2 (*PTGDR2*) (Hata and Breyer, 2004; Ricciotti and FitzGerald, 2011); 15d-PGJ2 preferentially binds DP2 (Monneret et al., 2002; Ricciotti and FitzGerald, 2011). To determine if either receptor drives senescence, we treated non-senescent cells with PGD2, a DP1-specific agonist (BW245C) (Town et al., 1983), a DP2-specific agonist (13,14-dihydro-15-keto-PGD2) (Hirai et al., 2001), both agonists or vehicle (DMSO). The DP1 agonist increased cAMP levels, while the DP2 agonist lowered cAMP levels (Figure S5A), confirming receptor activation (Hata and Breyer, 2004; Hirai et al., 2001; Hirata et al., 1994; Ricciotti and FitzGerald, 2011). PGD2 had no effect on cAMP levels (Figure S5A). Both the DP1 agonist and PGD2 reduced EdU incorporation, but only PGD2 elevated SA-Bgal activity (Figure S5B). PGD2 also increased mRNAs encoding p21 and p16^{INK4a} (Figure S5C and D), whereas the DP1 agonist lowered these mRNA levels. The DP2 agonist had no effect. Further, although *PTGDR* mRNA levels were unchanged between senescent and non-senescent cells, *PTGDR2* mRNA was ~50-fold lower in senescent cells, and thus unlikely to affect senescence (Figure S5D). PGD2 induced many mRNAs encoding SASP factors, but the DP1 agonist did not (Figure S5F). Finally, we treated cells with either a DP1 (BW-A868C) (Giles et al., 1989) or DP2 (OC000459) (Pettipher et al., 2012) antagonist for 10 days after inducing senescence by IR. Neither antagonist altered SASP gene expression (Figure S5G). These data indicate that PGD2 and its derivatives do not drive senescence through classic prostaglandin D2 receptor signaling, consistent with requirements for μ M concentrations (these receptors typically require nM concentrations). We therefore sought to identify alternative pathways.

15d-PGJ2 is the dehydration product of PGD2 (Forman et al., 1995; Kliwer et al., 1995), and dihydro-15d-PGJ2 is the most highly elevated prostaglandin inside senescent cells (Figure 1A), we therefore tested the possibility that intracellular dihydro-15d-PGJ2 promotes senescence, independent of cell surface receptor signaling. 15d-PGJ2 is a ligand for PPAR- γ (Forman et al., 1995; Kliwer et al., 1995), which is linked to senescence (Gan et al., 2008). We transduced IMR-90 fibroblasts with a baculovirus containing a luciferase PPAR- γ reporter and measured its activity in quiescent and SEN(IR) cells cultured in DMSO or the prostaglandin synthase inhibitor NS-398. PPAR- γ reporter activity increased ~18 fold upon senescence, and NS-398 lowered this activity by ~50% (Figure S5H), indicating that PPAR- γ activation is partially dependent on prostaglandin synthesis in senescent cells. We then cultured cells with two PPAR- γ agonists -- ciglitazone (20 μ M) or troglitazone (20 mM) -- for 10 days. Unlike 15d-PGJ2, neither agonist induced senescence, as determined by EdU incorporation and SA-Bgal activity (Figure S5I). Therefore, while prostaglandin

synthesis indeed activates PPAR- γ in senescent cells, PPAR- γ does not drive senescence phenotypes.

Prostaglandins reinforce senescence via positive RAS/p53 feedback

H-RAS is activated by exogenous 15d-PGJ2 at 3–5 μ M (Oliva et al., 2003) – roughly the dihomog-15d-PGJ2 level in senescent cells, and other cyclopentenone prostaglandins activate other RAS isoforms or similar GTPases (Anta et al., 2016; Renedo et al., 2007; Wall et al., 2015). We hypothesized that constitutive RAS activation might drive senescence in response to the elevated dihomog-15d-PGJ2. Indeed, incubation of quiescent cells with 15d-PGJ2 resulted in RAS activation, as well as phosphorylation of the downstream effector kinases ERK1/2 (Figure 5A; left lanes) (Oliva et al., 2003). Consistent with elevated 15d-PGJ2 levels, SEN(IR) cells showed increased RAS activation (RAS:GTP) and phospho-ERK1/2, which was attenuated by CAY or NS (Figure 5A). Similarly, shRNA-mediated knockdown of PTGDS (shPTGDS) selectively prevented RAS activation (Figure 5A). RAS activation is also associated with loss of intracellular dNTPs due to suppression of *RRM2* expression (Aird et al., 2013). A similar loss of *RRM2* occurred in IR-induced senescence, which was prevented by CAY or NS (Figure 5B). Importantly, administration of deuterated 15d-PGJ2 resulted in roughly equimolar accumulation of deuterated 15d-PGJ2 inside senescent cells (Figure S5J), demonstrating that exogenously-administered prostaglandins can accumulate intracellularly at levels that approximate those found in senescent cells.

RAS-induced senescence is driven by a mitogen activated kinase (MAPK) cascade, resulting in p53 phosphorylation on serine 37 (p53-S37) by p38-regulated activated kinase (PRAK/MAPKAPK5) (Sun et al., 2007). Indeed, PRAK was phosphorylated on threonine 182 (PRAK-R182), p53 was phosphorylated on serine 37 and p21 protein levels increased in cells induced to senesce by 15d-PGJ2 (Figures 5C–5D). These data suggest that 15d-PGJ2 reinforces senescence through a RAS-driven pathway.

H-RAS knockdown was toxic, so we used instead a RAF inhibitor (AZ 628) (Montagut et al., 2008) to block downstream RAS signaling. AZ 628 (20, 5 or 1 μ M) strongly inhibited ERK phosphorylation in response to serum stimulation without affecting basal levels of p53 or p21 (Figure S5K). Further, 1 μ M AZ 628 did not alter SA-b-gal activity (Figure S5L–S5M) or EdU labeling (Figure S5N). We therefore used 1 μ M for further studies. In response to 15d-PGJ2, AZ 628 blocked ERK1/2 and p53 phosphorylation, thus preventing the rise in p53 and p21 proteins (Figure 5D). AZ 628 also prevented the rise in SA-Bgal and decline in EdU labeling in response to 15d-PGJ2 (Figure 5E). Finally, continuous culture of either SEN(IR) or 15d-PGJ2-treated cells with AZ 628 prevented much of the rise in proinflammatory SASP factor mRNAs, lowered levels of both *PTGS2* and *PTGDS* mRNAs, and thus antagonized the positive feedback initiated by 15d-PGJ2 (Figure 5F). AZ 628 also prevented the rise in mRNAs encoding p21 and p16^{INK4a} and reduction in *LMNB1* mRNA in response to 10 Gy IR (Figures S5O–S5Q). Thus, activation of the RAS pathway is a key step in the development of senescent phenotypes, even in the absence of activating mutations.

Since 15d-PGJ2 activated p53 (S37 phosphorylation), and COX-2/PTGS2 inhibitors lowered *PTGS2* mRNA levels in senescent cells (Figure S4M), we tested the possibility that

p53 is required for the increased expression of prostaglandin synthase enzymes. Indeed, p53 depletion by a lentiviral shRNA (shp53) completely abolished *PTGS2* and *PTGDS* expression in response to IR (Figures 5G–5H). *CDKN1A/p21* expression was similarly attenuated, as expected (Figure 55R). By comparison, depletion of the NF- κ B subunit RELA using a lentiviral shRNA (shRELA) had no effect on prostaglandin synthase expression (Figures 55S–55T), despite lowering *IL6* expression (Figure 55U), as described (Freund et al., 2011). Thus, activation of p53 or RAS results in a positive feedback loop that promotes senescence and segments of the SASP by activating prostaglandin synthesis (Figure 5I). Given that prostaglandin synthesis is required for senescence, the data indicate that prostaglandin-mediated RAS activation is a common and necessary feature of at least some forms of cellular senescence.

Discussion

Our results indicate that oxylipins are a lipid-based component of the SASP. Secretion of these lipids likely has significant physiological effects, apart from the cell-autonomous role of dihomog-15d-PGJ2. For example, chorioamnion-derived PGE2 and PGF2 α are key signals for parturition (Gibb, 1998) and senescent cells appear in the chorioamniotic membranes of women in labor (Behnia et al., 2015; Menon et al., 2016), suggesting that senescent cells may be a source of parturition-inducing prostaglandins.

Importantly, we observe intracellular accumulation of cyclopentenone prostaglandins, such as dihomog-15d-PGJ2. This accumulation was linked to increased expression of prostaglandin transporter, but not prostaglandin dehydrogenase, which might explain the accumulation of 15-deoxy prostaglandins. Increased import without a commensurate change in PGDH might promote cytoplasmic accumulation of dehydrated 15-deoxy prostaglandins, rather than 15-keto prostaglandins.

Given that elevated intracellular dihomog-15d-PGJ2 and its biosynthetic enzymes occurs in several cell types and tissues in response to multiple senescence inducers, we propose that the synthesis of dihomog-15d-PGJ2 and RAS activation are common features of senescent cells. While oncogenic RAS mutations are known to drive cells into senescence (Serrano et al., 1997), our studies identify a role for wild type RAS in senescence. We show that the RAS-RAF signaling pathway is important for many aspects of senescence, including components of the SASP, increased levels of p21 and p16^{INK4a}, and decline in lamin B1 levels.

We previously showed that p53 restrains segments of the SASP by antagonizing NF- κ B signaling (Freund et al., 2011; Wiley et al., 2018). Here, we show that p53 is required for prostaglandin biosynthesis, and inhibition of prostaglandin synthesis antagonizes the SASP. These findings explain why p53 does not completely prevent the SASP. Thus, inhibition of prostaglandin synthesis did not return inflammatory cytokine levels to those of non-senescent cells, unlike *RELA* depletion, suggesting that prostaglandins contribute to, but are not exclusively necessary for, the secretion of proinflammatory factors. The role of p53-dependent prostaglandin synthesis in development of the SASP is likely overshadowed

by the loss of NF- κ B antagonism when p53 is depleted, resulting in increased inflammatory cytokine secretion in the absence of p53, even when prostaglandin synthesis is lost.

Senescence and the SASP are dynamic phenotypes that change over time (De Cecco et al., 2019; Hernandez-Segura et al., 2017). The lipids described here likely undergo dynamic regulation as well. Indeed, we recently showed that expression of specific lipid synthases is temporally regulated in the context of pulmonary fibrosis, suggesting a transition from acute to more chronic stress phenotypes (Wiley et al., 2019a).

Finally, our finding that released dihomog-15d-PGJ2 can be used as a biomarker for senolysis has several potential applications. Senolytic drugs are increasingly being used in aging and related research and – importantly – have entered early clinical trials (Hickson et al., 2019; Justice et al., 2019). Determining that senolysis is taking place is essential for evaluating these compounds as therapeutic agents. Detection of dihomog-15d-PGJ2 in biological fluids may allow rapid evaluation of the efficacy of these compounds.

Limitations of the study.

Outside of dihomog-15d-PGJ2, most of the factors in Figure 1A and 1B are assigned by m/z, and therefore are only likely assignees. We list these factors in the figure legend as “putative” as a result. This study only features analyses from mice and human cells, and for 3 cell types as representatives of endothelial, epithelial, and mesenchymal lineages. Results may not reflect all lineages or inducers of senescence.

STAR Methods

RESOURCE AVAILABILITY.

Lead contact.—Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Judith Campisi (jcampisi@buckinstitute.org).

Material Availability.—This study did not generate unique reagents. The schematic for synthesis of dihomog-15d-PGJ2 will be provided upon request.

Data and Code Availability.—Lipid profiling data is available at the NIH Common Fund’s National Metabolomics Data Repository (NMDR) website, the Metabolomics Workbench (<https://www.metabolomicsworkbench.org>), Project ID #: PR001093. See also the Key Resources Table.

EXPERIMENTAL MODEL AND SUBJECT DETAILS.

Cell lines and strains.—All cells were purchased from the American Type Culture Collection (ATCC), see Key Resource Table. IMR-90 human female fetal lung fibroblasts were the primary cell type used in this study, and were used prior to 40 population doublings - other than replicative senescent cells, which were cultured until replicative exhaustion. HEPG2 human adolescent male hepatocellular carcinoma are immortalized, but p53-positive, and therefore no population doublings were recorded. IMR-90 and HEPG2 cells were cultured in Dulbecco’s modified eagle medium (DMEM 4.5 g/L glucose, without

sodium pyruvate - Gibco) supplemented with 10% FBS and penicillin/streptomycin (Gibco). HUVEC human female umbilical vein endothelial cells were cultured using the ATCC protocol and culture media and assayed within 40 population doublings. Quiescence was induced by replacing culture media with media containing 0.2% FBS for 72 h before analysis. All cells were cultured at 37°C and 3% O₂. All cells were mycoplasma free.

Animals.—Animal experiments were conducted using a protocol approved by the Institutional Animal Care and Use Committee of the Buck Institute. Mice were housed in alternating 12:12 hour light/dark cycles at 68–72°F. General animal health rounds were performed daily, 7 days a week by trained Vivarium staff members. Any noted illness or deaths were reported via the Health Surveillance system to the Vivarium Director and/or the Attending Veterinarian (AV). The Buck Vivarium is an SPF barrier facility and routine health screening is done facility-wide to ensure the health of all colonies. Sentinel mice that were exposed to soiled bedding and byproducts from other mice on the same rack were screened quarterly for common murine viruses as directed by the AV. Internal and external parasitology screening was also done on all sentinel mice. For DOXO treatments, 10–16 wk old male p16–3MR transgenic mice on a C57BL/6 background received one intraperitoneal (i.p.) injection of 10 mg/kg of doxorubicin hydrochloride in PBS, and treated 5 d later with GCV or vehicle. GCV was administered via daily i.p. injections for 5 consecutive days at 25 mg/kg in PBS. Control mice were injected with an equal volume of PBS. Mice were euthanized and tissues collected 10 d after DOXO challenge. For aging studies, 8 male and 8 female C57BL/6 mice were aged for 6 or 24 mo, at which point mice were euthanized and tissues collected for analysis. For the biomarker studies, 10 male C57BL/6 mice per cohort with challenged with DOXO as above, and given either vehicle or ABT-263 6 weeks after DOXO challenge. Whole blood was collected by cardiac puncture into EDTA-tubes (lavender caps – BD Biosciences) and spun at 2000g for 5 minutes. For urine collection, mice were placed in plastic containers prior to euthanasia.

METHOD DETAILS.

Gene expression.—RNA was extracted from cells or tissues using commercially available kits (Isolate II - Bioline for cells; Direct-zol - Zymo for tissues) according to the manufacturer's instructions. cDNA synthesis was performed using a High Capacity cDNA Reverse Transcription Kit (Thermo Fisher) according to the manufacturer's instructions. Quantitative PCR was performed on a LightCycler 480 II (Roche) using primers and probes designed for the Universal Probe Library. Primers and probes used for human transcripts are listed in Table S2.

Induction of senescence.—Senescence SEN(IR) was induced by either 5 or 10 Gy ionizing-irradiation (IR). Non-senescent controls (proliferating or quiescent) were placed in the irradiator for an identical period of time but without irradiation. Oncogene-induced senescence was induced using lentiviral-mediated overexpression of HRAS^{V12}, as described (Wiley et al., 2016), and the lentiviral backbone (Vector) served as a control. MiDAS was induced by depleting mitochondrial DNA in the presence of 100 ng/mL ethidium bromide, 100 ug/mL sodium pyruvate, and 50 ug/mL uridine for 2 months, followed by removal of pyruvate, as described (Wiley et al., 2016).

Generation of conditioned media.—Conditioned media (CM) were generated by culturing cells in appropriate serum-free media supplemented with penicillin/streptomycin for 24 h before harvest, followed by clarification by centrifugation. Cells were then trypsinized and counted. All quantitative assays (e.g. ELISAs) from CM were normalized to cell number.

Induction of apoptosis or senolysis.—Non-senescent or SEN(IR) cells were cultured in the presence of 10 μ M ABT-263 for 24 h. Non-senescent IMR-90 fibroblasts were treated with either 1 μ M staurosporine (Cayman) or 100 ng/mL recombinant TRAIL (Abcam) for 24 h before collection. Media were centrifuged twice at 5000xg for 15 min at 4°C for clearing of apoptotic bodies and stored at –80°C until analysis.

Senescence-associated beta-galactosidase.—SA-Bgal activity was detected as described (Dimri et al., 1995) using a commercial kit (Biovision).

Colony forming assay.—Ras-transduced senescent cells were plated at a density of 1000/well and cultured in the presence of growth media containing either DMSO, CAY-10404, or NS-398 for 7 days. Cells were washed in PBS, fixed in 95% methanol, and incubated for 30 min in 0.2% crystal violet in 2% methanol. Crystal violet was removed and wells were washed in tap water until the solution ran clear, and allowed to dry.

RAS activation assay.—Active RAS (RAS:GTP) was immunoprecipitated from culture lysates using a commercial kit (Cytoskeleton, Inc) following the manufacturer's instructions. Precipitates were analyzed by immunoblot.

Immunoblots.—Cells were lysed in 5% SDS in 10 mM Tris, pH 7.4, and protein content determined by BCA assay. 20 μ g protein was separated by electrophoresis and transferred to PVDF membranes. Membranes were blocked in TBST + 5% BSA, incubated overnight with primary antibody, washed in TBST, incubated with HRP-conjugated secondary antibody for 30 min, and visualized by chemiluminescence – combined with either film or on a BioRAD Hood II Gel Doc universal imager. Antibodies to phosphorylated p53-S37, p21, phospho-ERK1/2, total ERK1/2 and total p38MAPK were from Cell Signaling. Antibodies against LMNB1, HMGB1, and actin were from Abcam, anti-phospho-p38 was from PhosphoSolutions, and anti-PRAK was from Novus. Anti-phospho-PRAK-T182 was a kind gift from Peiqing Sun, and was described in (Yoshizuka et al., 2012). Anti-RAS was provided in the kit provided by Cytoskeleton, Inc (above).

IL-6 ELISA.— 3×10^4 cells in 12-well plates were treated as indicated, and cultured in 0.5–1 ml serum-free DMEM for 24 hr. CM were collected and clarified at 2,000x g for 10 min. Supernatants were transferred to a tube; cells were trypsinized and counted. CM (2.5 μ l) were analyzed by bead-based ELISAs (AlphaLISA, Perkin-Elmer) as instructed by the manufacturer and normalized to cell number.

PGD2-MOX ELISA.—CM were isolated as for IL-6 ELISA (above), and analyzed according the manufacturer's instructions (Cayman Chemical), and quantified as pg/cell.

15d-PGJ2 ELISA.— 1×10^5 cells in 6-well plates were treated as indicated for 10 days, and then cultured in 1 ml of serum-free medium plus DMEM or ABT-263 for 24 h. CM were then clarified by centrifugation at 2,500x g for 10 min. 15d-PGJ2 ELISA was from Enzo Life Sciences, and was performed according to the manufacturer's instructions. Cell culture supernatant was analyzed directly, and all numbers were normalized to cell number. For in vivo fluids, 250 μ L of blood plasma or 50 μ L of urine were extracted as in Liquid-liquid extraction (LLE - below) with the following modification – dehydrated lipid samples were resuspended in 250 μ L of assay buffer before analysis.

Reporter assays.—Baculoviral PPAR γ reporter constructs (Signosis) were transduced into proliferating cells. Following treatment, cells were lysed using Passive Lysis Buffer (Promega) and analyzed for luciferase activity using a commercial kit (Promega) and Perkin-Elmer VictorTMX3 luminometer.

Chemicals and standards.—Ammonium acetate was from Sigma Aldrich (St. Louis, MO). HPLC-grade solvents acetonitrile and methanol were from Fisher Scientific (Pittsburgh, PA, USA) and VWR (Radnor, PA, USA). Deionized water was generated in-house. PGA2, PGD2, PGE2, PGF2 α , PGJ2, 15d-PGJ2, 15d-PGJ2-d4, NS-398, CAY-10404, ciglitazone, troglitazone and staurosporine were from Cayman Chemical. ABT-263 was from Apex Biotechnology. The dihomio-15d-PGJ2 standard was synthesized by Jubilant Chemsys (Bangalore, India). TRAIL was from Abcam.

Liquid-liquid extraction (LLE) of lipids.—Lipid extraction was performed as reported (Bligh and Dyer, 1959; Folch et al., 1951) with some modifications. Cells were rinsed with PBS and quenched using 1mL 50% methanol; 2 μ g/mL 13C1-leucine and 5 ng/mL hexanesulfonic acid were added as internal standards. 2 mL of chloroform with 1 μ g/mL heptadecanoic acid was added to each sample and mixed for 10 min at 4° C. Samples were centrifuged at 4,000g for 15 min at 4° C to separate aqueous and organic layers. After centrifugation, 700 μ L of the aqueous layer and 1.5 mL of the organic layer were recovered and concentrated by speedvac and N₂, respectively. Both aqueous and organic fractions were reconstituted in 100 μ L 50% methanol and chloroform prior to liquid-chromatography mass spectrometry (LC-MS) analysis.

Solid phase extraction (SPE) of lipids.—Lipid extraction was performed based as reported (Harkewicz et al., 2007) with some modifications. Cells were rinsed with PBS and quenched in 1 mL methanol with 2 μ g/mL 13C1-leucine, 1 μ g/mL heptadecanoic acid, and 5 ng/mL hexanesulfonic acid added as internal standards. 1 mL PBS was added to each sample and mixed for 10 min. Samples were centrifuged at 4,000g for 15 min at 4° C. Following centrifugation, oxylipins were separated on Phenomenex Strata-X polymeric sorbent columns connected to a vacuum manifold. Columns were pre-washed with 2 mL methanol followed by 2 mL water. Samples were loaded onto columns, washed with 2 mL 90:10 water:methanol, then eluted with 1mL 100% methanol. Extracts were concentrated by speedvac and reconstituted in 100 μ L 100% methanol prior to LC-MS analysis.

High-pressure liquid chromatography quadrupole-time-of-flight mass spectrometry (HPLC-QTOF-MS).—LC-MS analyses were performed on a Agilent

6520 QTOF mass spectrometer coupled to a Agilent 1260 Infinity liquid chromatography system consisting of the following modules: u-degasser (G1322A), binary pump (G1312B), thermostated column compartment (G1330B), and HiPALS auto sampler (G1367E). Chromatographic separation of cellular extracts was performed on a Phenomenex Luna NH₂ (2.0mm × 150mm, 3.0μM) column. The mobile phase included A:20mM ammonium acetate and 5% acetonitrile, pH9.5 and B:acetonitrile. The gradient is as follows: 0 to 20min, 95–10%B, 25–30min 10%B, and 30.1–35min 95%B. LC conditions included an auto sampler temperature of 4° C, injection volume of 10μL and solvent flow-rate of 0.3mL/min. Analyses were performed using the following ionization parameters: gas temperature (TEM) 350°C; drying gas, 9L/min; Vcap, 2500V; nebulizer, 35psig; fragmentor, 125V; and skimmer, 65V. MS1 acquisition was operated in the negative ion scanning mode for a mass range of 50–1000 m/z.

LC-MS data was acquired and analyzed using Agilent MassHunter Workstation (B.05.00), Agilent MassHunter Qualitative Analysis Software (B.08.00), Mass Profiler Professional (B.12.0), and Microsoft Excel 2007 (Redmond, WA, USA). Peak areas were assigned using Agilent MassHunter Qualitative Analysis Software in combination with the Find by Formula (FBF) algorithm. Peak areas were normalized to total protein or cell number.

High-pressure liquid chromatography quadrupole ion trap mass spectrometry (HPLC-QTRAP-MS).—HPLC was performed using a Shimadzu UFLC prominence

system fitted with following modules: CBM-20A (Communication bus module), DGU-A₃(degasser), two LC-20AD (liquid chromatography, binary pump), SIL-20AC HT (auto sampler) and connected to a Phenomenex Luna NH₂ (2.0mm × 150mm, 3.0μM) column. The solvent system was A=20mM ammonium acetate pH 9.5 with 5% acetonitrile and B=acetonitrile. The starting gradient conditions were 95% B at a flow rate of 0.3mL/min. The following gradient program was used: 0 to 20min, 95–10%B, 25–30min 10%B, and 30.1–35min 95%B. Samples were kept at +4°C, and the injection volume was 10μL. Mass spectrometric analysis was conducted using negative ion electrospray ionization in the multiple reaction monitoring mode (MRM) on an AbSciex 4000 QTRAP (Foster City, CA, USA) mass spectrometer fitted with a TurboV™ ion source. The ionization parameters were set as follows: curtain gas (CAD); 20psi; collision gas: medium; ion spray voltage (IS): –4500V; Temperature (TEM): 550°C; Ion source Gas 1 (GS1); 60psi; and Ion source Gas 2 (GS2): 50psi. The compound parameters were established using appropriate standards. The compound parameters were set as follows: entrance potential (EP): –10.0V; and collision cell exit potential (CXP): –5V. ABSciex Analyst®v1.6.1 was used for all data acquisition and an in-depth analysis of the HPLC-MS data, specifically for calculating the peak areas for the identified features from cellular extracts. Peak areas were normalized to total protein.

Reverse-phase LC-MS method for detection of 15d-PGJ₂.—The LC-MS analysis was performed using Agilent 6520 accurate mass quadrupole time of flight (Q-ToF) mass spectrometer coupled with Agilent 1260 Infinity Pump (Agilent Technologies, Santa Clara, CA). Ionization was performed in negative mode. Nitrogen was used as a desolvation and collision gas. The separation was carried out using Kinetex EVO C18 100Å 150×2.1, 5 micron (serial #H15173269) from Phenomenex. The autosampler was set at 5°C. The

solvent system was composed of Solvent A- 0.1% acetic acid in 95:5 (water: ACN) containing 10 mM ammonium acetate and Solvent B- 0.1% acetic acid in 95:5 (ACN: water) containing 10 mM ammonium acetate. The flow rate was maintained at 0.3 mL/min, and the initial solvent conditions started with 5% solvent B. At 2 min, the percentage of B was increased to 70% B over 14 min, followed by further increase of % B to 95% for 10 mins. At 24 mins % B was maintained at 95% B for 5 mins and then reduced to 5%B and maintained at 5% B for 5 mins. The run time was 35 mins. Drying gas flow and temperature was set at 9.0 L/min and 350 °C, respectively and nebulizer gas pressure was set at 35 psig. The applied capillary voltage was 3000 V. Full scan spectra was acquired from 100–1000 m/z. The instrument was operated with Agilent MassHunter Work station LC/MS Data Acquisition version 05.01, and chromatograms were processed with MassHunter Workstation qualitative software version B.08.00.

Quantitation of intracellular dihom-15d-PGJ2 concentration.—A standard curve was generated using 100, 50, 25, and 10 nM dihom-15d-PGJ2, and peak areas were quantified by mass spectrometry. Based upon an average measured concentration of 69 nM in 1 mL of cell lysate representing 5×10^6 senescent cells, and assuming an average volume of 7.5 pL per senescent cell, we estimated a mean intracellular concentration of 1.8 μ M by mass spectrometry, or 5.8 μ M by ELISA.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data Representation and Statistical Analysis.—All data are presented as means $\pm$ SEM, with the exception of box plots, which follow the standard median/quartile structure. All culture datasets represent means of at least 3 experiments, and mouse datasets reflect a minimum of 6 mice. Comparisons between groups were performed using a 2-tailed student t-test, 1- or 2-way ANOVA, as appropriate. No methods were used to determine whether the data met the assumptions of the statistical approaches. Heat maps use $p < 0.05$ for all entities. Significance is indicated by * = $p < 0.05$, ** = $p < 0.01$, and *** = $p < 0.001$. Statistical parameters can be found in Figure legends.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements.

The authors thank Peiqing Sun for the gift of the phospho-PRAK antibody, and Pierre-Yves Desprez for critical reading of this manuscript. This work was supported by grants from the National Institutes of Health (AG051729, AG057353, AG0172442, U2C-DK119886, and T32-AG000266) and the Science and Engineering Research Board of India (DST- SERB SPR/2019/001265).

References

- Acosta JC, O'Loughlen A, Banito A, Guisjarro MV, Augert A, Raguz S, Fumagalli M, Da Costa M, Brown C, Popov N, et al. (2008). Chemokine signaling via the CXCR2 receptor reinforces senescence. *Cell* 133, 1006–1018. [PubMed: 18555777]

- Aird KM, Zhang G, Li H, Tu Z, Bitler BG, Garipov A, Wu H, Wei Z, Wagner SN, Herlyn M, et al. (2013). Suppression of nucleotide metabolism underlies the establishment and maintenance of oncogene-induced senescence. *Cell Rep* 3, 1252–1265. [PubMed: 23562156]
- Anta B, Perez-Rodriguez A, Castro J, Garcia-Dominguez CA, Ibiza S, Martinez N, Dura LM, Hernandez S, Gragera T, Pena-Jimenez D, et al. (2016). PGA1-induced apoptosis involves specific activation of H-Ras and N-Ras in cellular endomembranes. *Cell Death Dis* 7, e2311. [PubMed: 27468687]
- Baker DJ, Childs BG, Durik M, Wijers ME, Sieben CJ, Zhong J, Saltness RA, Jeganathan KB, Verzosa GC, Pezeshki A, et al. (2016). Naturally occurring p16(Ink4a)-positive cells shorten healthy lifespan. *Nature* 530, 184–189. [PubMed: 26840489]
- Behnia F, Taylor BD, Woodson M, Kaceroovsky M, Hawkins H, Fortunato SJ, Saade GR, and Menon R (2015). Chorionic membrane senescence: a signal for parturition? *Am J Obstet Gynecol* 213, 359 e351–316. [PubMed: 26025293]
- Bligh EG, and Dyer WJ (1959). A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37, 911–917. [PubMed: 13671378]
- Buratta S, Urbanelli L, Sagini K, Giovagnoli S, Caponi S, Fioretto D, Mitro N, Caruso D, and Emiliani C (2017). Extracellular vesicles released by fibroblasts undergoing H-Ras induced senescence show changes in lipid profile. *PLoS One* 12, e0188840. [PubMed: 29182668]
- Cadenas C, Vosbeck S, Hein EM, Hellwig B, Langer A, Hayen H, Franckenstein D, Buttner B, Hammad S, Marchan R, et al. (2012). Glycerophospholipid profile in oncogene-induced senescence. *Biochim Biophys Acta* 1821, 1256–1268. [PubMed: 22178194]
- Campisi J (2013). Aging, cellular senescence, and cancer. *Annu Rev Physiol* 75, 685–705. [PubMed: 23140366]
- Castellone MD, Teramoto H, Williams BO, Druey KM, and Gutkind JS (2005). Prostaglandin E2 promotes colon cancer cell growth through a Gs-axin-beta-catenin signaling axis. *Science* 310, 1504–1510. [PubMed: 16293724]
- Catalano A, Rodilossi S, Caprari P, Coppola V, and Procopio A (2005). 5-Lipoxygenase regulates senescence-like growth arrest by promoting ROS-dependent p53 activation. *EMBO J* 24, 170–179. [PubMed: 15616590]
- Chang J, Wang Y, Shao L, Laberge RM, Demaria M, Campisi J, Janakiraman K, Sharpless NE, Ding S, Feng W, et al. (2016). Clearance of senescent cells by ABT263 rejuvenates aged hematopoietic stem cells in mice. *Nat Med* 22, 78–83. [PubMed: 26657143]
- Coppe JP, Patil CK, Rodier F, Sun Y, Munoz DP, Goldstein J, Nelson PS, Desprez PY, and Campisi J (2008). Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol* 6, 2853–2868. [PubMed: 19053174]
- Cormenier J, Martin N, Desle J, Salazar-Cardozo C, Pourtier A, Abbadie C, and Pluquet O (2018). The ATF6alpha arm of the Unfolded Protein Response mediates replicative senescence in human fibroblasts through a COX2/prostaglandin E2 intracrine pathway. *Mech Ageing Dev* 170, 82–91. [PubMed: 28803844]
- Dagouassat M, Gagliolo JM, Chrusciel S, Bourin MC, Duprez C, Caramelle P, Boyer L, Hue S, Stern JB, Validire P, et al. (2013). The cyclooxygenase-2-prostaglandin E2 pathway maintains senescence of chronic obstructive pulmonary disease fibroblasts. *Am J Respir Crit Care Med* 187, 703–714. [PubMed: 23328527]
- De Cecco M, Ito T, Petrashen AP, Elias AE, Skvir NJ, Criscione SW, Caligiana A, Broccoli G, Adney EM, Boeke JD, et al. (2019). L1 drives IFN in senescent cells and promotes age-associated inflammation. *Nature* 566, 73–78. [PubMed: 30728521]
- Demaria M, O'Leary MN, Chang J, Shao L, Liu S, Alimirah F, Koenig K, Le C, Mitin N, Deal AM, et al. (2017). Cellular Senescence Promotes Adverse Effects of Chemotherapy and Cancer Relapse. *Cancer Discov* 7, 165–176. [PubMed: 27979832]
- Demaria M, Ohtani N, Youssef SA, Rodier F, Toussaint W, Mitchell JR, Laberge RM, Vijg J, Van Steeg H, Dolle ME, et al. (2014). An essential role for senescent cells in optimal wound healing through secretion of PDGF-AA. *Dev Cell* 31, 722–733. [PubMed: 25499914]
- Dennis EA, and Norris PC (2015). Eicosanoid storm in infection and inflammation. *Nat Rev Immunol* 15, 511–523. [PubMed: 26139350]

- Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I, Pereira-Smith O, et al. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A* 92, 9363–9367. [PubMed: 7568133]
- Flor AC, Wolfgeher D, Wu D, and Kron SJ (2017). A signature of enhanced lipid metabolism, lipid peroxidation and aldehyde stress in therapy-induced senescence. *Cell Death Discov* 3, 17075. [PubMed: 29090099]
- Folch J, Ascoli I, Lees M, Meath JA, and Le BN (1951). Preparation of lipide extracts from brain tissue. *J Biol Chem* 191, 833–841. [PubMed: 14861228]
- Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, and Evans RM (1995). 15-Deoxy-delta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 83, 803–812. [PubMed: 8521497]
- Freund A, Patil CK, and Campisi J (2011). p38MAPK is a novel DNA damage response-independent regulator of the senescence-associated secretory phenotype. *EMBO J* 30, 1536–1548. [PubMed: 21399611]
- Funk CD (2001). Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science* 294, 1871–1875. [PubMed: 11729303]
- Gan Q, Huang J, Zhou R, Niu J, Zhu X, Wang J, Zhang Z, and Tong T (2008). PPAR{gamma} accelerates cellular senescence by inducing p16INK4{alpha} expression in human diploid fibroblasts. *J Cell Sci* 121, 2235–2245. [PubMed: 18544633]
- Gibb W (1998). The role of prostaglandins in human parturition. *Ann Med* 30, 235–241. [PubMed: 9677008]
- Giles H, Leff P, Bolofo ML, Kelly MG, and Robertson AD (1989). The classification of prostaglandin DP-receptors in platelets and vasculature using BW A868C, a novel, selective and potent competitive antagonist. *Br J Pharmacol* 96, 291–300. [PubMed: 2924081]
- Han JH, Roh MS, Park CH, Park KC, Cho KH, Kim KH, Eun HC, and Chung JH (2004). Selective COX-2 inhibitor, NS-398, inhibits the replicative senescence of cultured dermal fibroblasts. *Mech Ageing Dev* 125, 359–366. [PubMed: 15130753]
- Harkewicz R, Fahy E, Andreyev A, and Dennis EA (2007). Arachidonate-derived dihomoprostaglandin production observed in endotoxin-stimulated macrophage-like cells. *J Biol Chem* 282, 2899–2910. [PubMed: 17135246]
- Hata AN, and Breyer RM (2004). Pharmacology and signaling of prostaglandin receptors: multiple roles in inflammation and immune modulation. *Pharmacol Ther* 103, 147–166. [PubMed: 15369681]
- Hernandez-Segura A, de Jong TV, Melov S, Guryev V, Campisi J, and Demaria M (2017). Unmasking Transcriptional Heterogeneity in Senescent Cells. *Curr Biol* 27, 2652–2660 e2654. [PubMed: 28844647]
- Hickson LJ, Langhi Prata LGP, Bobart SA, Evans TK, Giorgadze N, Hashmi SK, Herrmann SM, Jensen MD, Jia Q, Jordan KL, et al. (2019). Senolytics decrease senescent cells in humans: Preliminary report from a clinical trial of Dasatinib plus Quercetin in individuals with diabetic kidney disease. *EBioMedicine* 47, 446–456. [PubMed: 31542391]
- Hirai H, Tanaka K, Yoshie O, Ogawa K, Kenmotsu K, Takamori Y, Ichimasa M, Sugamura K, Nakamura M, Takano S, et al. (2001). Prostaglandin D2 selectively induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH2. *J Exp Med* 193, 255–261. [PubMed: 11208866]
- Hirata M, Kakizuka A, Aizawa M, Ushikubi F, and Narumiya S (1994). Molecular characterization of a mouse prostaglandin D receptor and functional expression of the cloned gene. *Proc Natl Acad Sci U S A* 91, 11192–11196. [PubMed: 7972033]
- Jump DB (2009). Mammalian fatty acid elongases. *Methods Mol Biol* 579, 375–389. [PubMed: 19763486]
- Justice JN, Nambiar AM, Tchkonja T, LeBrasseur NK, Pascual R, Hashmi SK, Prata L, Masternak MM, Kritchevsky SB, Musi N, et al. (2019). Senolytics in idiopathic pulmonary fibrosis: Results from a first-in-human, open-label, pilot study. *EBioMedicine*.
- Kabir TD, Leigh RJ, Tasena H, Mellone M, Coletta RD, Parkinson EK, Prime SS, Thomas GJ, Paterson IC, Zhou D, et al. (2016). A miR-335/COX-2/PTEN axis regulates the secretory

phenotype of senescent cancer-associated fibroblasts. *Aging* (Albany NY) 8, 1608–1635. [PubMed: 27385366]

- Kim J, Vaish V, Feng M, Field K, Chatzistamou I, and Shim M (2016). Transgenic expression of cyclooxygenase-2 (COX2) causes premature aging phenotypes in mice. *Aging* (Albany NY) 8, 2392–2406. [PubMed: 27750221]
- Kliwer SA, Lenhard JM, Willson TM, Patel I, Morris DC, and Lehmann JM (1995). A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 83, 813–819. [PubMed: 8521498]
- Kuilman T, Michaloglou C, Mooi WJ, and Peeper DS (2010). The essence of senescence. *Genes Dev* 24, 2463–2479. [PubMed: 21078816]
- Laberge RM, Zhou L, Sarantos MR, Rodier F, Freund A, de Keizer PL, Liu S, Demaria M, Cong YS, Kapahi P, et al. (2012). Glucocorticoids suppress selected components of the senescence-associated secretory phenotype. *Aging Cell* 11, 569–578. [PubMed: 22404905]
- Lee JW, Mok HJ, Lee DY, Park SC, Ban MS, Choi J, Park CG, Ahn YS, Kim KP, and Kim HD (2016). UPLC-MS/MS-Based Profiling of Eicosanoids in RAW264.7 Cells Treated with Lipopolysaccharide. *Int J Mol Sci* 17, 508. [PubMed: 27058537]
- Martien S, Pluquet O, Vercamer C, Malaquin N, Martin N, Gosselin K, Pourtier A, and Abbadie C (2013). Cellular senescence involves an intracrine prostaglandin E2 pathway in human fibroblasts. *Biochim Biophys Acta* 1831, 1217–1227. [PubMed: 24046862]
- Menon R, Behnia F, Poletini J, Saade GR, Campisi J, and Velarde M (2016). Placental membrane aging and HMGB1 signaling associated with human parturition. *Aging* (Albany NY) 8, 216–230. [PubMed: 26851389]
- Monneret G, Li H, Vasilescu J, Rokach J, and Powell WS (2002). 15-Deoxy-delta 12,14-prostaglandins D2 and J2 are potent activators of human eosinophils. *J Immunol* 168, 3563–3569. [PubMed: 11907120]
- Montagut C, Sharma SV, Shioda T, McDermott U, Ulman M, Ulkus LE, Dias-Santagata D, Stubbs H, Lee DY, Singh A, et al. (2008). Elevated CRAF as a potential mechanism of acquired resistance to BRAF inhibition in melanoma. *Cancer Res* 68, 4853–4861. [PubMed: 18559533]
- Nomura T, Lu R, Pucci ML, and Schuster VL (2004). The two-step model of prostaglandin signal termination: in vitro reconstitution with the prostaglandin transporter and prostaglandin 15 dehydrogenase. *Mol Pharmacol* 65, 973–978. [PubMed: 15044627]
- Ohno Y, Suto S, Yamanaka M, Mizutani Y, Mitsutake S, Igarashi Y, Sassa T, and Kihara A (2010). ELOVL1 production of C24 acyl-CoAs is linked to C24 sphingolipid synthesis. *Proc Natl Acad Sci U S A* 107, 18439–18444. [PubMed: 20937905]
- Oliva JL, Perez-Sala D, Castrillo A, Martinez N, Canada FJ, Bosca L, and Rojas JM (2003). The cyclopentenone 15-deoxy-delta 12,14-prostaglandin J2 binds to and activates H-Ras. *Proc Natl Acad Sci U S A* 100, 4772–4777. [PubMed: 12684535]
- Perretti M, and D'Acquisto F (2009). Annexin A1 and glucocorticoids as effectors of the resolution of inflammation. *Nat Rev Immunol* 9, 62–70. [PubMed: 19104500]
- Pettipher R, Vinall SL, Xue L, Speight G, Townsend ER, Gazi L, Whelan CJ, Armer RE, Payton MA, and Hunter MG (2012). Pharmacologic profile of OC000459, a potent, selective, and orally active D prostanoid receptor 2 antagonist that inhibits mast cell-dependent activation of T helper 2 lymphocytes and eosinophils. *J Pharmacol Exp Ther* 340, 473–482. [PubMed: 22106101]
- Quijano C, Cao L, Fergusson MM, Romero H, Liu J, Gutkind S, Rovira II, Mohny RP, Karoly ED, and Finkel T (2012). Oncogene-induced senescence results in marked metabolic and bioenergetic alterations. *Cell Cycle* 11, 1383–1392. [PubMed: 22421146]
- Renedo M, Gayarre J, Garcia-Dominguez CA, Perez-Rodriguez A, Prieto A, Canada FJ, Rojas JM, and Perez-Sala D (2007). Modification and activation of Ras proteins by electrophilic prostanoids with different structure are site-selective. *Biochemistry* 46, 6607–6616. [PubMed: 17489560]
- Ricciotti E, and FitzGerald GA (2011). Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol* 31, 986–1000. [PubMed: 21508345]
- Rosner BA, and Cristofalo VJ (1979). Hydrocortisone: a specific modulator of in vitro cell proliferation and aging. *Mech Ageing Dev* 9, 485–496. [PubMed: 459571]

- Serrano M, Lin AW, McCurrach ME, Beach D, and Lowe SW (1997). Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell* 88, 593–602. [PubMed: 9054499]
- Soberman RJ, and Christmas P (2003). The organization and consequences of eicosanoid signaling. *J Clin Invest* 111, 1107–1113. [PubMed: 12697726]
- Sprecher H, VanRollins M, Sun F, Wyche A, and Needleman P (1982). Dihomo-prostaglandins and -thromboxane. A prostaglandin family from adrenic acid that may be preferentially synthesized in the kidney. *J Biol Chem* 257, 3912–3918. [PubMed: 6801054]
- Strong R, Miller RA, Astle CM, Floyd RA, Flurkey K, Hensley KL, Javors MA, Leeuwenburgh C, Nelson JF, Ongini E, et al. (2008). Nordihydroguaiaretic acid and aspirin increase lifespan of genetically heterogeneous male mice. *Aging Cell* 7, 641–650. [PubMed: 18631321]
- Sun P, Yoshizuka N, New L, Moser BA, Li Y, Liao R, Xie C, Chen J, Deng Q, Yamout M, et al. (2007). PRAK is essential for ras-induced senescence and tumor suppression. *Cell* 128, 295–308. [PubMed: 17254968]
- Town MH, Casals-Stenzel J, and Schillinger E (1983). Pharmacological and cardiovascular properties of a hydantoin derivative, BW 245 C, with high affinity and selectivity for PGD2 receptors. *Prostaglandins* 25, 13–28. [PubMed: 6302737]
- Wall SB, Oh JY, Mitchell L, Laube AH, Campbell SL, Renfrow MB, and Landar A (2015). Rac1 modification by an electrophilic 15-deoxy Delta(12,14)-prostaglandin J2 analog. *Redox Biol* 4, 346–354. [PubMed: 25677088]
- Wiley CD, Brumwell AN, Davis SS, Jackson JR, Valdovinos A, Calhoun C, Alimirah F, Castellanos CA, Ruan R, Wei Y, et al. (2019a). Secretion of leukotrienes by senescent lung fibroblasts promotes pulmonary fibrosis. *JCI Insight* 4.
- Wiley CD, and Campisi J (2016). From Ancient Pathways to Aging Cells-Connecting Metabolism and Cellular Senescence. *Cell Metab* 23, 1013–1021. [PubMed: 27304503]
- Wiley CD, Liu S, Limbad C, Zawadzka AM, Beck J, Demaria M, Artwood R, Alimirah F, Lopez-Dominguez JA, Kuehnemann C, et al. (2019b). SILAC Analysis Reveals Increased Secretion of Hemostasis-Related Factors by Senescent Cells. *Cell Rep* 28, 3329–3337 e3325. [PubMed: 31553904]
- Wiley CD, Schaum N, Alimirah F, Lopez-Dominguez JA, Orjalo AV, Scott G, Desprez PY, Benz C, Davalos AR, and Campisi J (2018). Small-molecule MDM2 antagonists attenuate the senescence-associated secretory phenotype. *Sci Rep* 8, 2410. [PubMed: 29402901]
- Wiley CD, Velarde MC, Lecot P, Liu S, Sarnoski EA, Freund A, Shirakawa K, Lim HW, Davis SS, Ramanathan A, et al. (2016). Mitochondrial Dysfunction Induces Senescence with a Distinct Secretory Phenotype. *Cell Metab* 23, 303–314. [PubMed: 26686024]
- Yang HH, Kim C, Jung B, Kim KS, and Kim JR (2011). Involvement of IGF binding protein 5 in prostaglandin E(2)-induced cellular senescence in human fibroblasts. *Biogerontology* 12, 239–252. [PubMed: 21191810]
- Yoshizuka N, Chen RM, Xu Z, Liao R, Hong L, Hu WY, Yu G, Han J, Chen L, and Sun P (2012). A novel function of p38-regulated/activated kinase in endothelial cell migration and tumor angiogenesis. *Mol Cell Biol* 32, 606–618. [PubMed: 22124154]
- Zdanov S, Bernard D, Debacq-Chainiaux F, Martien S, Gosselin K, Vercamer C, Chelli F, Toussaint O, and Abbadie C (2007). Normal or stress-induced fibroblast senescence involves COX-2 activity. *Exp Cell Res* 313, 3046–3056. [PubMed: 17560572]

Highlights

- Senescent cells make several oxylipins, dihomoprostaglandins, and leukotrienes
- Dihomo-15d-PGJ2 is intracellular during senescence and is released during senolysis
- Dihomo-15d-PGJ2 activates RAS, promoting senescence and the SASP
- Positive feedback between prostaglandins, RAS, and p53 maintains senescence




Figure 1. Senescent cells synthesize oxylipins.

A-B. IMR-90 cells were proliferative in 10% serum, made quiescent by incubation for 3 days in 0.2% serum, or made senescent by treatment with 10 Gy X-rays (IR) or ethidium bromide to induce mitochondrial-dysfunction induced senescence (MiDAS). Lipids were extracted from proliferating (PRO – 10%), quiescent (QUI – 0.2%), IR-induced senescent (SEN(IR) – 10% or 0.2% serum), or mitochondrial dysfunction-associated senescent (MiDAS – 0.2%) cells and analyzed by liquid chromatography combined with mass spectrometry (LC-MS). Putative oxylipins (A) and lipid precursors (B) were detected in control and senescent cells. Heat maps indicate the averages of at least 3 experiments (* = $p < 0.05$, 1-way ANOVA). C. RNA was isolated from QUI, SEN(IR), MiDAS and vector or RAS-V12 expressing [SEN(RAS)] cells, reverse transcribed, and mRNAs encoding oxylipin synthesis genes were measured by quantitative (q) PCR. Heat maps indicate the averages of 3 experiments (* = $p < 0.05$, 1-way ANOVA). D. p16–3MR mice were given a single dose (10 mg/kg) of doxorubicin (DOXO) or phosphate-buffered saline (PBS) by intraperitoneal (i.p.) injection. After 5 days, mice were given GCV (25 mg/kg) or vehicle by i.p. injection for 5 consecutive days. Livers were harvested on day 10 and analyzed for oxylipin synthase mRNAs by qPCR. Heat maps indicate averages of at least 5 mice (* = $p < 0.05$, 1-way ANOVA). E. IMR-90 fibroblasts were induced to senesce by MiDAS, IR, or RAS-V12 overexpression. 10 days after induction of senescence, conditioned media were collected and analyzed for PGD2 by ELISA. F-G. HEPG2 (F) and HUVEC (G) cells were induced to senesce by IR. Conditioned media were collected 10 days after IR and analyzed for PGD2

by ELISA. Data show means of at least 3 experiments (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, unpaired, 2-tailed t test). H-I. RNA from (C) was analyzed for expression of SLCO2A1 or PGDH by qPCR. J. HUVECs were induced to senesce by IR, and analyzed at Day 10 for SLCO2A1 and PGDH as in H and I. See also Figure S1 and Table S1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. (Dihomo)15d-PGJ2 is a biomarker of senolysis.

A. IMR-90 fibroblasts were induced to senesce by MiDAS or IR. 10 days after induction, cells were treated with DMSO or 10 μ M ABT-263, and (dihomo)15d-PGJ2 [(dh)-15d-PGJ2] was measured in conditioned media by ELISA. Data reflect means of at least 3 experiments. B-C. HEPG2 (B) or HUVEC (C) cells were induced to senesce by IR, and conditioned media from DMSO or ABT-263 treated cells were generated as in (A). (dh)15d-PGJ2 was measured by ELISA. D. Representative overlay of dihomom-15d-PGJ2 peaks from extracted ion chromatograms of lipids from media treated as in A. E-I. C57BL/6 mice were injected intraperitoneally with doxorubicin (DOXO; 10 mg/kg) or PBS. 6 weeks later, ABT-263 (50 mg/kg) or vehicle were administered by gavage. Data are presented as box plots for at least 6 mice per condition. E-F. Epididymal fat pads were harvested 3 hours after ABT-263 administration and analyzed for p16^{INK4a} (E) or p21^{WAF1} (F) mRNA levels by qPCR. G. 3 hours after gavage, blood was collected by cardiac puncture, and (dh)15d-PGJ2 was measured by ELISA. H. 12 hours after gavage, urine was collected, and (dh)15d-PGJ2 was measured by ELISA. I. Representative overlay of dihomom-15d-PGJ2 peaks from extracted ion chromatograms of lipids extracted from plasma of the animals from (G). * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, 1-way ANOVA. See also Figure S2.




Figure 3. Validation of dihomom-15d-PGJ2 as a biomarker of senolysis.

A. A synthesized dihomom-15d-PGJ2 standard was analyzed at 3 different voltages for fragmentation patterns. Key identification peaks included 299.2 m/z ($C_{21}H_{31}O^-$: green box), 283.2 m/z ($C_{21}H_{31}^-$: blue box), 255.2 ($C_{19}H_{27}^-$: red box), 233.2 ($C_{15}H_{21}O_2^-$: purple box). B. Extracted ion chromatograms from a dihomom-15d-PGJ2 standard (top), plasma from a DOXO+ABT-263 mouse (middle), and senescent cell extracts (bottom). Filled peaks appeared only in senolysis (middle) or senescence (bottom). C. Fragmentation patterns of filled peaks from B indicating a dihomom-15d-PGJ2 standard (top), plasma

from a DOXO+ABT-263 mouse (middle), or senescent cell extracts (bottom). D. Zoomed chromatograms for the 299.2 m/z peak (zoomed area indicated by green boxes on Figure 3C) in DOXO + ABT-263 mouse plasma (upper) or senescent cell extracts (lower). Note that relative abundance of a non-specific peak (unfilled in Figure 3B) required earlier fragmentation in SEN(IR) extracts. See also Figure S3.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. Prostaglandins promote cellular senescence phenotypes.

A-F. IMR-90 fibroblasts were induced to senescence by IR and treated with DMSO or COX-2 inhibitors CAY-10404 (CAY) or NS-398 (NS) for 10 days. A. Conditioned media were harvested and secreted PGD2 was measured by ELISA. B. 24 h EdU incorporation for irradiated cells in each treatment group. C. SA-Bgal positivity for irradiated cells in each treatment group. D. p21 mRNA levels in each treatment group. E. p16 mRNA levels from each treatment group. F. Conditioned media from (A) were analyzed for IL-6 secretion by ELISA. G. mRNA levels of genes encoding SASP factors from all treatment groups were measured by qPCR. H-N. Proliferating cells were cultured for 10 days in the presence of DMSO or 5 μ M 15d-PGJ2 or dihom-15d-PGJ2. H. 24 h EdU labeling indices for each treatment. I. Example of SA-Bgal positive cells after each treatment. Panels are cropped zooms to larger images, and red saturation has been lowered for all images in order to allow visualization of the blue x-gal staining. J. SA-Bgal positivity for each treatment. K. p21 mRNA levels in extracts from prostaglandin- and DMSO-treated cells. L. *p16^{INK4a}*

mRNA levels in extracts from prostaglandin- and DMSO-treated cells. M. *LMNB1* mRNA levels in extracts from prostaglandin- and DMSO-treated cells. N. SASP factor PTGS2, PTGDS and PTGES mRNA levels were measured by qPCR. Normalized to actin. O. Cells were transduced with lentiviruses expressing shRNA targeting either GFP (shGFP; control) or PTGDS (shPTGDS), irradiated as in (A), and mRNA extracted 10 d after irradiation. SASP gene expression was measured by qPCR. Bar graphs indicate averages of at least 3 experiments (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, unpaired, 2-tailed t test). Heat maps indicate averages of 3 experiments (* = $p < 0.05$, 2-way ANOVA). See also Figure S4.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 5. 15d-PGJ2 promotes senescent phenotypes by activating RAS.

A. Extracts from IMR-90 fibroblasts were analyzed for activated RAS (Ras:GTP), total Ras, phospho-ERK1/2, total ERK1/2, and actin by immunoblot. (Left) Quiescent cells were treated with DMSO or 5 μ M 15d-PGJ2 for 15 minutes before harvest. (Right) Cells were induced to senesce by 10 Gy IR, and treated with DMSO, CAY-10404, NS-398, or lentiviral shRNAs to GFP (shGFP) or PTGDS (shPTGDS). Cells were cultured for 10 days prior to harvest, and in 0.2% FBS for the last 48 hours. B. Cells were induced to senesce by 5 Gy IR, and cultured in the presence of vehicle (DMSO), CAY or NS for 10 d. RNA was extracted, reverse-transcribed, and analyzed for RRM2 mRNA relative to actin mRNA by qPCR. C. Immunoblot of cells treated with DMSO or 10 μ M 15d-PGJ2 for 10 days. Blots were probed for phospho-PRAK-T182, total PRAK, phospho-p53-S37, and total p53, p21 and actin. D. Cells were treated with vehicle or 15d-PGJ2, and DMSO or AZ 628 for 10 d. Protein was extracted and analyzed by immunoblot for phospho-ERK1/2, total ERK1/2, phospho-p53-S37, and total p53, p21 and actin. E. Cells treated as in D were analyzed for SA-Bgal or 24 h EdU incorporation. F. Cells were induced to senesce by 10

Gy IR or continuous culture with 10 μ M 15d-PGJ2 for 10 days. DMSO or AZ 628 were concurrently added from Day 0 and present until day 10. Quiescent cells cultured in DMSO served as a control. RNA was then extracted and analyzed by qPCR for mRNAs encoding SASP factors, PTGS2 and PTGDS. G-H. Cells were transduced with lentiviral shRNAs to either GFP (shGFP) or p53 (shp53), mock irradiated or given 10 Gy IR and cultured for 10 days (shGFP) or 3 days (shp53) before RNA extraction and analysis for (G) PTGS2 or (H) PTGDS mRNAs. I. Model for a feedback loop involving 15d-PGJ2, RAS and p53 that reinforces senescent phenotypes. Chemiluminescent western images were captured with either film or by Bio-rad Gel-Doc. Image formats may therefore be inconsistent between lanes within panels. Bar graphs indicate averages of at least 3 experiments (***) = $p < 0.001$, unpaired, 2-tailed t test). Heat maps indicate averages of 3 experiments (* = $p < 0.05$, 2-way ANOVA). See Also Figure S5.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Phospho-PRAK-T182	Peiqing Sun - Wake Forest Univeristy	
Phospho-p53 (Ser37) Antibody	Cell Signaling Technology	#9289
p21 Waf1/Cip1 (12D1) Rabbit mAb	Cell Signaling Technology	#2947
Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Antibody	Cell Signaling Technology	#9101
p44/42 MAPK (Erk1/2) Antibody	Cell Signaling Technology	#9102
p38 MAPK Antibody	Cell Signaling Technology	#9212
Anti-Lamin B1 antibody	Abcam	ab16048
Anti-HMGB1 antibody	Abcam	ab18256
Monoclonal Anti- β -Actin antibody	Sigma	A2228
p53 Antibody (DO-1)	Santa Cruz Biotechnology	sc-126
Bacterial and Virus Strains		
shGFP lentivirus	Campisi Lab	
shP53 lentivirus	Campisi Lab	
shPTGDS lentivirus	Sigma-Aldrich	SHCLNV-NM_000954
Chemicals, Peptides, and Recombinant Proteins		
NS-398	Cayman Chemical Company	70590
CAY-10404	Cayman Chemical Company	70210
Ciglitazone	Cayman Chemical Company	71730
Troglitazone	Cayman Chemical Company	71750
Staurosporine	Cayman Chemical Company	81590
Recombinant Human TRAIL protein	Abcam	ab78818
Arachidonic Acid-d ₁₁	Cayman Chemical Company	10006758
Prostaglandin A ₂	Cayman Chemical Company	10210
Prostaglandin D ₂	Cayman Chemical Company	12010
Prostaglandin E ₂	Cayman Chemical Company	14010
Prostaglandin F _{2α}	Cayman Chemical Company	16010
Prostaglandin J ₂	Cayman Chemical Company	18500
15-deoxy- $\Delta^{12,14}$ -Prostaglandin J ₂	Cayman Chemical Company	18570
1a,1b-dihomo-15-deoxy- $\Delta^{12,14}$ -Prostaglandin J ₂	Jubilant Chemsys	P1171
13,14-dihydro-15-keto Prostaglandin D ₂	Cayman Chemical Company	12610
BW A868C	Cayman Chemical Company	12060
OC000459	Cayman Chemical Company	12027
BW 245C	Cayman Chemical Company	12050
Hydrocortisone	Cayman Chemical Company	20739
Acetic Acid, glacial	VWR	VW0125-3
Acetonitrile	VWR/BDH	BDH83639.400

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Ammonium acetate	Sigma Aldrich	73594–25G-F
LCMS Water	Honeywell Burdick & Jackson	LC365–4
Methanol	Honeywell Burdick & Jackson	LC230–4
Chloroform	JT Baker	9175–2
Critical Commercial Assays		
Baculoviral PPAR-gamma reporter assay	Signosis	BT-0014
Senescence Detection Kit	BioVision	K320
15-deoxy- Δ 12,14-PGJ2 ELISA Kit	ENZO	ADI-900–023
Prostaglandin D2-MOX Express ELISA Kit	Cayman Chemical Company	500151
IL-6 AlphaLISA	PerkinElmer	AL223F
Ras Pull-down Activation Assay Biochem Kit	Cytoskeleton, Inc	BK008-S
Experimental Models: Cell Lines		
HEPG2	ATCC	HB-8065
IMR-90	ATCC	CCL-186
HUVEC-C	ATCC	CRL-1730
Experimental Models: Organisms/Strains		
C57BL/6J	The Jackson Laboratory	000664
p16–3MR	Campisi Laboratory	
Teklad global 18% protein	Envigo Teklad	2918
Oligonucleotides		
See Table S2		
Software and Algorithms		
MassHunter	Agilent	
Other		
Kinetex 5 μ m EVO C18 100Å 150x2.1mm	Phenomenex	00F-4633-AN
Data accessibility		
Lipid Profiling Dataset	National Metabolomics Data Repository (NMDR)	PR001093